

BIBLIOGRAPHY

South Baffin Island Study Area

GENERAL REFERENCES

ECOLOGICAL REFERENCES

WILDLIFE REFERENCE

MARINE MAMMAL REFERENCES

HISTORICAL INFORMATION

ADDITIONAL REFERENCES

COASTAL CLASSIFICATION
ILLUIS AREA 13 South Baffin Island

CODE	CLASS AND DESCRIPTION	DIAGRAMS (NOT TO SCALE)
Bc	CONTEMPORARY BEACH COMPLEX - High energy depositional environments, which are usually extensive. Spits, bars, lagoons, former beach ridges and storm beaches all common, as are gravel forerlands and raised beaches inland. Coarse sands and gravels are typical materials.	
Br	RAISED BEACH SPURRING - Series of beach ridges occurring on slopes steeper than 5° to 10° extending up to 2 km inland. Surface materials are gravel. Narrow contemporary beaches of low energy are ubiquitous. Wider beaches are classed as Bc, and strandlines on generally level plains are classed as Fg.	
Ca	CLIFFS WITH NO TALLS. Precipitous rock slopes, that range from 65° to vertical or overhanging. The cliff face may include prominent rock ledges. This class has no beach and occurrence is very limited.	
Cb	CLIFFS WITH LESS THAN 1/3 COVERED BY BLOCKY, RUBBLY COLLUVIUM. As class Ca but with up to one third of the height covered by colluvial deposits of various thickness and extent. Colluvium is normally barren to sparsely vegetated. Narrow (1-3m) coarse gravel to boulder beaches are common.	
Cd	CLIFFS WITH 1/3 TO 2/3 COVERED BY BLOCKY, RUBBLY COLLUVIUM. As class Cb, but with more extensive colluvium.	
Ce	CLIFFS WITH MORE THAN TWO-THIRDS TALLS. As class Cc, but mostly talus covered. As talus attains complete cover, this class grades into Bc.	
G	GLACIERS - glacier ice which reaches the sea and results in calving. Unit includes associated lateral moraines and bluffs of till with small gravel beaches. Classes Cc, Cd and Ce are normally found juxtaposed. Where the ice has retreated to leave a plain, class Fg or Fz takes precedence.	
Hb	HUFFS - ERODING HILLS. Steep slopes over 10m high of unconsolidated materials, generally free of talus, resulting from past or present erosion at the base. Gullies and a narrow beach at low tide are common.	
Hc	COLLUVIAL HILLSLOPES. Smooth sloping colluvium, typically undergoing sheet wash or solifluction extending up narrow, gravel beaches. This class occurs mostly in sheltered waters; exposure to significant wave action would quickly erode these materials and create beach or foreland conditions. This class grades into Cc with increasing slope.	
Hr	ROCKY HILLS. Mainly bedrock controlled slopes, over 5-10° which extend from over 10m elevation. Pockets of colluvium small pocket in crevices, and sections of precipitous rock slopes are common. Stretches of narrow, coarse textured beaches may extend along the shoreline. Islands and reefs are common offshore.	
Hs	SLOPES. Steep over 10 m. high slopes of unconsolidated, or weakly consolidated materials, which display rotational or planar slumping. Pool scars, and flows, extent and narrow tidal beaches are common.	
Pb	ERODED PLAINS - CUTNEASES. Coastal plains of unconsolidated, cohesive materials such as clay - till or laminated sands, which are eroded at the shoreline to produce low bluff backbeaches and typically narrow beaches. Similar coasts in bedrock are classified Fz.	
Pc	ROCKY PLAINS WITH LOW CLIFFS. These normally occur as low rock forerlands in front of rocky hills or high cliffs with talus. These rock forerlands range in width from 10's of metres to 1+2 km and are typically less than 10 m high. Near vertical cliffs are the seaward end are common. Beaches of any kind are rare.	
Pf	FLUVIAL PLAINS - SANDFLATS - Extensive deposits (up to 100 km ²) of unconsolidated alluvial sand and/or silt resulting from rapid terrestrial erosion inland, and coinciding with low beach development and free nearshore conditions; significant such as channels and beach ridges are quickly obliterated by wind action. Shorelines may be marked by push ridges, but it is also common to find no beach forms whatever, just a smooth progression from land to sea.	
PI	IRREGULAR PLAIN (formerly till plain, Pt). Similar to Pt in broad relief - plains of less than 5-10° elevation near the coast and slopes typically less than 5° to 10°, covered with fine beach development and shallow nearshore. Medium to coarse sediments and irregular terrain prevail, such as related to normal plain or strandlines. Strandlines on progressively inclined surface are classified as Br.	
PI	INCLINED PLAIN: Plains of less than 10° elevation near the coast and slopes typically less than 5-10°, covered with limited beach development and shallow nearshore. Fine beach development, organic materials and smooth terrain prevail, such as related to emergent marine deposits or pediments. Local drainage is typically parallel and perpendicular to the coast.	
PI	LEVEL PLAIN. Extensive deposits of marine and/or fluvial deposits of fine textured materials coinciding with wave and current free nearshore conditions. Backshore is typically an almost zero raised plain with narrow shallow ponds, and possibly widely spaced low strandlines. Shorelines may be indicated by beach ridges, but it is also common to find no beach forms whatever, just a smooth progression from land to sea. Typically, the plain extends offshore with wide near-shore or tidal flats.	
Po	ORGANIC PLAINS. A very low, often featureless plain or fringe of organic materials, usually associated with very sheltered waters. Shores are low banks of organic accumulations or progressive retreating of open water and emergent vegetation. Usually recorded as the minor component in a shoreline rating.	
Pr	ROCKY PLAINS. Rocky slopes of low elevation usually less than 10 m elevation generally less than 5-10°. Patchy colluvial or moraine veneer and pocket beaches and fans and intermittently to scattered narrow gravel beaches occur. Islands, reefs and shoals are common offshore.	
Ps	PLAINS WITH STEEP BACKSHORE (other than rock controlled). Forelands or plains consisting of unconsolidated sediments, usually as raised fans or raised marine platforms covered by beach deposits past or present. The plain extends abruptly to steep backshore usually 20-35° with a narrow contemporary beach.	
Rf	RIVER MOUTH AND FLUVIAL COMPLEXES - Includes features associated with river mouths: bars, deltas, estuaries, tidal flats, marshes, baymouth bars, spits, etc. Smaller fans grade into Fz where isostatic rebound exceeds the stream's debris supply, causing the fan to be raised and the stream to incise.	
U	UNDIFFERENTIATED CLIFFS, HILLS OR PLAINS. In cases where imagery is poor and no field checks were made, it may be impossible to determine the sub-class.	

NTING AND TRAPPING