

BIBLIOGRAPHY

South Victoria Island - Northeast Keewatin District Study Area
WEST FALD
GENERAL REFERENCES
WILDLIFE

Stefanov, V. 1911. THE FRIENDLY ARCTIC. THE STORY OF THE EXPEDITION TO INVESTIGATE THE POSSIBILITIES OF THE INTERIOR AND MEXICO INDUSTRIES IN THE NORTHERN CANADA.
Swatman, Gordon K. 1951. THE BARN SWALLOW AT CAMBRIDGE BAY, N.W.T.
Tamer, J.R. 1958. THE DISTRIBUTION OF MUSKELIN IN CANADA.
Jalabach, R.D., and D.R. Carruthers. 1980. CARIBOU AND MUSKELIN ON VICTORIA ISLAND, N.W.T.

Mitchell, R. and R.B. Rowley. 1981. Geology and cumulative effects of initial population size of cetaceans in the eastern Canadian Arctic.
Hunter, J. G., S. Leach, D. B. McAllister, W. Steigerwald. 1983. Distribution atlas of the narwhal (Monodon monoceros) in the eastern Canadian Arctic.
Jacobson, R. 1980. Land use for resource harvesting on Victoria Island, Northwest Territories.
Jacobson, R. 1980. Land use for resource harvesting on Victoria Island, Northwest Territories.

Stewart, B. and L. R. J. Bernier. 1982. An aquatic resource survey of King William and Victoria Islands, District of Franklin, Northwest Territories.
Stewart, B. and L. R. J. Bernier. 1983. An aquatic resource survey of King William and Victoria Islands, District of Franklin, and the northeastern District of Keewatin, Northwest Territories.
Jacobson, R. 1980. Land use for resource harvesting on Victoria Island, Northwest Territories.

COASTAL CLASSIFICATION
NATIVE LAND USE INFORMATION
CLIFFS WITH MORE THAN ONE THIRD TALUS COVER AS CLASS Cc
CLIFFS WITH ONE THIRD TO TWO THIRDS TALUS COVER AS CLASS Cb
GLACIERS - glacier ice which reaches the sea and results in past or present erosion at the base.
BLUFFS - ERODING HILLS. Steep slopes over 10m high of unconsolidated materials, generally free of talus.