

Manitoba Land Resource Unit
Brandon Research Centre
Agriculture and Agri-Food Canada

Rural Municipality of Lorne
Information Bulletin 97-15

Soils and Terrain

An introduction
to the land resource



Rural Municipality of Lorne

Information Bulletin 97-15

Prepared by:

Manitoba Land Resource Unit,
Brandon Research Centre, Research Branch,
Agriculture and Agri-Food Canada.

Department of Soil Science, University of Manitoba.

Manitoba Soil Resource Section,
Soils and Crops Branch, Manitoba Agriculture.

Printed March, 1998

PREFACE

This is one of a new series of information bulletins for individual rural municipalities of Manitoba. They serve to introduce the newly developed digital soil databases and illustrate several typical derived and interpretive map products for agricultural land use planning applications. The bulletins will also be available in diskette format for each rural municipality.

Information contained in this bulletin may be quoted and utilized with appropriate reference to the originating agencies. The authors and originating agencies assume no responsibility for the misuse, alteration, re-packaging, or re-interpretation of the information.

This information bulletin serves as an introduction to the land resource information available for the municipality. More detailed information, including copies of the primary soil and terrain maps at larger scales, may be obtained by contacting

Manitoba Land Resource Unit
Room 360 Ellis Bldg, University of Manitoba,
Winnipeg, Manitoba R3T 2N2
Phone: 204-474-6118 FAX: 204-474-7633.

CITATION

Manitoba Land Resource Unit, 1997. Soils and Terrain. An Introduction to the Land Resource. Rural Municipality of Lorne. Information Bulletin 97-15, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.

ACKNOWLEDGEMENTS

This project was financially supported in part by the Canada-Manitoba Agreement on Agricultural Sustainability, Prairie Farm Rehabilitation Administration (PFRA), and Agriculture and Agri-Food Canada.

The following individuals and agencies contributed significantly to the compilation, interpretation, and derivation of the information contained in this report.

Managerial and administrative support was provided by:

R.G. Eilers, Head, Manitoba Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.

G.J. Racz, Head, Dept. of Soil Science, University of Manitoba.

F. Wilson, Manager, Manitoba Land and Soil Programs, PFRA, Agriculture and Agri-Food Canada

K.S. McGill, Manager, Soil Resource Section, Soils and Crops Branch, Manitoba Agriculture.

Technical support was provided by:

G.W. Lelyk, Manitoba Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.

J. Fitzmaurice, N. Lindberg, A. Waddell, M. Fitzgerald and S. Grift, Dept. of Soil Science, University of Manitoba.

J. Griffiths, C. Aglugub, Soil Resource Section, Soils and Crops Branch, Manitoba Agriculture.

R. Lewis, PFRA, Agriculture and Agri-Food Canada.

G.F. Mills, P.Ag, Winnipeg, Manitoba

Professional expertise for data conversion, correlation, and interpretation was provided by:

W.R. Fraser and W. Michalyna, Manitoba Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.

P. Haluschak and G. Podolsky, Soil Resource Section, Soils and Crops Branch, Manitoba Agriculture.

LAND RESOURCE DATA

The soil and terrain information presented in this bulletin was compiled as part of a larger project to provide a uniform level of land resource information for agricultural and regional planning purposes throughout Agro-Manitoba. This information was compiled and analysed in two distinct layers as shown in Figure 2.




Figure 2. Soil and Base Map data.

Base Layer

Digital base map information includes the municipality and township boundaries, along with major streams, roads and highways. Major rivers and lakes from the base layer were also used as common boundaries for the soil map layer. Water bodies larger than 25 ha in size were digitized as separate polygons.

Soil Layer

The most detailed soil information currently available was selected as the data source for the digital soil layer for each rural municipality.

Comprehensive detailed soil maps (1:20 000 to 1:50 000 scale) have been published for many rural municipalities. Where they were available, the individual soil map sheets were digitized and compiled as a single georeferenced layer to match the digital RM base. Map polygons have one or more soil series components, as well as slope and stoniness classes. Soil database information was produced for each polygon, to meet national standards (MacDonald and Valentine, 1992). Slope length classes were also added, based on photo-interpretation.

Older, reconnaissance scale soil maps (1:126 720 scale) represented the only available soil data source for many rural municipalities. These maps were compiled on a **soil association** basis, in which soil landscape patterns were identified with unique surficial geological deposits and textures. Each soil association consists of a range of different soils ("associates") each of which occurs in a repetitive position in the landscape. Modern soil series that best represent the soil association were identified for each soil polygon. The soil and modifier codes provide a link to additional databases of soil properties. In this way, both detailed and reconnaissance soil map polygons were related to soil drainage, surface texture, and other soil properties to produce various interpretive maps. Slope length classes were also added, based on photo-interpretation.

LAND RESOURCE OVERVIEW

The Rural Municipality (RM) of Lorne covers an area of 10 townships (approximately 97 600 hectares) of land in south-central Manitoba (page 3). The towns of Notre Dame de Lourdes, Somerset, Swan Lake, and Mariapolis are the largest population and service centres in the municipality.

The climate in the municipality can be related to weather data from Somerset. The mean annual temperature is 1.8°C and the mean annual precipitation is 547 mm (Environment Canada, 1982). The average frost-free period is 124 days and degree-days above 5°C average 1543 (Ash, 1991). The calculated seasonal moisture deficit for the period between May and September is 200 to 250 mm. The estimated effective growing degree days (EGDD) above 5°C accumulated from May to September is 1400 to 1500 (Agronomic Interpretations Working Group, 1995). These parameters provide an indication of moisture and heat energy available for crop growth.

Physiographically, the RM of Lorne is located within the Saskatchewan Plain (Canada-Manitoba Soil Survey, 1980). The majority of the RM occurs in the Pembina Hills Upland whereas the northwest corner of the municipality is in the Tiger Hills Upland. A small part of the Pembina Escarpment crosses the northeast corner of the RM. The large meltwater channel of the Pembina Valley and occurs along the southern edge of the municipality. Elevations within the RM range from a high of 525 m asl in the northwest near Bruxelles to 330 m asl in the Pembina Escarpment to the east. The RM is characterized by a dominantly hummocky to rolling land surface interspersed with areas of nearly level terrain with slopes of less than 2 percent (page 9). Local relief ranges from 3 to 9 m with moderate slopes of 5 to 9 percent. Scattered areas of higher relief with 15 to 30 percent slopes occur throughout the RM. Greatest local relief is associated with the Pembina Valley with a depth of 60 m and very steeply sloping areas (slopes in excess of 30 percent) along the valley sides. High relief also occurs in eroded channels in the Pembina Escarpment and along portions of the Cypress River.

The most extensive soil materials in the RM are loamy textured glacial tills. Loamy and fine loamy lacustrine sediments overlie the till in gently sloping landscapes. Sandy to coarse loamy textured glacio fluvial deposits occur at scattered locations throughout the municipality and underlie extensive areas of coarse loamy lacustrine deposits north of the Pembina Channel. Prominent ridges and knolls throughout the area are commonly underlain by weathered shale or bedrock at shallow depth (page 11).

Soils in the municipality have been mapped at a semi-detailed level (1:50 000 scale) and published in the report, Soils of the Rural Municipality of Lorne, Soil Report No. D70. (Langman, 1986). According to the Canadian System of Soil Classification, (Expert Committee on Soil Survey, 1987), the soils in the municipality are classified as Dark Gray and Black Chernozems and Humic Gleysols. Regosolic soils occur on stratified minor stream deposits and on steeply sloping areas of eroded slopes. A more detailed and complete description of the type, distribution and textural variability of soils in the municipality is provided in the soil report.

Surface drainage for the largest portion of the area is via the Cypress River. Drainage for the eastern part of the RM occurs to the east from the Pembina Escarpment and a few intermittent creeks provide drainage to the south into the Pembina Valley. Surface drainage of hummocky terrain in the RM is largely local in nature. The majority of soils in the RM are well drained with significant areas of imperfect drainage on lower slopes and local areas of poor drainage (page 13). Surface runoff collects in poorly drained depressions and potholes, many of which contain shallow ponds and small lakes. Soils bordering the Pembina Channel and along the Pembina Escarpment are well to rapidly drained due in part to deep channels eroded into the land surface.

Minor areas of scattered soil salinity (page 15) occur in association with imperfectly drained soils in drainage channels and bordering poorly drained depressions. Salinity is more extensive in the Tiger Hills Upland in the northwestern portion of the municipality. In addition to salinity, other management considerations are primarily

related to topography and wetness (page 17). Sandy textures are of local importance throughout the area and there are no significant bedrock outcrops. Local areas of slightly stony soils occur in till landscapes while areas of lacustrine soil are stone-free.

Approximately two-thirds of the land in the RM is rated as **Class 1, 2 and 3** for agriculture capability (page 19) and three-quarters of the soils are classified as **Good to Fair** for irrigation suitability (page 21). Topography and excess water (wetness) are the main limitations for agriculture. Well drained soils in gently sloping landscapes are rated in **Class 1 or 2** for agriculture and **Good** for irrigation. Very poorly drained soils are rated in **Class 6** and steeply sloping land is rated in **Class 7** for agriculture and **Poor** for irrigation. Salinity affected soils in the RM which are rated in **Class 3 or Class 5** for agriculture depending on the degree of wetness and **Poor** for irrigation.

A major issue currently receiving considerable attention is the sustainability of agricultural practices and their potential impact on the soil and groundwater environment. To assist in highlighting this concern to land planners and agricultural producers, an assessment of potential environmental impact (EI) under irrigation has been included in this bulletin (page 23). Approximately 50 percent of the soils in the RM are at **Low** risk of degradation. However, areas of sandy soil, coarse sandy and gravelly subsoil, very poorly drained soil and steeply sloping soils are rated as having a **High** potential for environmental impact under irrigation. The coarse textures increase the risk for deep leaching of potential contaminants on the soil surface and the steep slopes increase the rapid runoff into adjacent wetlands or water bodies. This EI map is intended to be used in association with the irrigation suitability map.

Another issue of concern to producers and soil conservation and land use specialists is soil erosion caused by agricultural cropping and tillage practices. To highlight areas with potential for water erosion, a risk map has been included to show where special practices should be adopted to mitigate this risk (page 25). As shown, over 30 percent of the land in the RM is at **Severe** risk to

degradation from water erosion, primarily due to steeply sloping topography. A **High** risk for erosion occurs on 12 percent of the land while nearly 20 percent of the area is at **Moderate** risk to degradation. Management practices focus primarily on maintaining adequate crop residues to provide sufficient surface cover. However, protection of the steeper sloping lands most at risk to water erosion may require a shift in land use away from annual cultivation to production of perennial forages and pasture or permanent tree cover.

An assessment of the status of land use in the RM of Lorne in 1994 was obtained through an analysis of satellite imagery. It showed that annual crops occupied about 67% of the land in the RM, while the remaining areas were in grassland (11.9%), forest (8.3%) and forage production (3.7%). Wetlands and small water bodies occupy 5.3% of the RM. Woodland and grassland areas provide native pasture for livestock. Various non-agricultural uses such as recreation and infrastructure for urban areas and transportation occupy about 3.4 % of the RM (page 27).

While most of the soils in the RM of Lorne have moderate to moderately severe limitations for arable agriculture, management of steeply sloping soils and coarser textured loamy and sandy soils requires careful choice of crops and maintenance of adequate surface cover. This includes leaving adequate crop residues on the surface to provide sufficient trash cover during the early spring period. The provision of shelter belts, minimum tillage practices, and crop rotations including forage will help to reduce the risk of soil degradation, maintain productivity and help insure that agriculture and use is sustainable over the long-term.

DERIVED AND INTERPRETIVE MAPS

A large variety of computer derived and interpretive maps can be generated from the digital soil and landscape databases. These maps are based on selected combinations of database values and assumptions.

Derived maps show information that is given in one or more columns in the computer map legend (such as soil drainage, soil salinity, or slope class).

Interpretive maps portray more complex land evaluations based on a combination of soil and landscape information. Interpretations are based on soil and landscape conditions in each polygon. Interpretative maps typically show land capabilities, suitabilities, or risks related to sustainability.

Several examples of derived and interpretive maps are included in this information bulletin:

Derived Maps

Slope

Surface Texture

Drainage

Salinity

Management Considerations

Interpretative Maps

Agricultural Capability

Irrigation Suitability

Potential Environmental Impact

Water Erosion Risk

Land Use.

Digital databases derived from recent detailed soil inventories contain additional detailed information about significant inclusions of differing soil and slope conditions in each map polygon. This information can be portrayed at larger map scale than shown in this bulletin.

Information concerning particular interpretive maps, and the primary soil and terrain map data, can be obtained by contacting the Manitoba Soil Resource Section of Manitoba Agriculture, the local PFRA office, or the Manitoba Land Resource Unit.

The maps have all been reduced in size and generalized (simplified) in order to portray conditions for an entire rural municipality on one page. These generalized maps provide a useful overview of conditions within a municipality, but are not intended to apply to site specific land parcels. On-site evaluations are recommended for localized site specific land use suitability requirements.

Slope Map.


Slope describes the steepness of the landscape surface. The slope classes shown on this map are derived from the digital soil layer database. Specific colours are used to indicate the dominant slope class for each soil polygon in the RM. Additional slope classes may occur in each polygon area, but cannot be portrayed at this reduced map scale.

Table 1. Slope Classes¹

Slope Class	Area (ha)	Percent of RM
0 - 2 %	35748	36.6
2 - 5 %	29328	30.0
5 - 9 %	11468	11.7
9 - 15 %	10870	11.1
15 - 30 %	2983	3.1
> 30 %	4430	4.5
Unclassified	0	0.0
Water	2805	2.9
Total	97631	100.0


¹ Area has been assigned to the dominant slope class in each soil polygon.

Slope Map



Slope Classes

Dark Green	0 - 2 %
Green	2 - 5 %
Light Green	5 - 9 %
Yellow	9 - 15 %
Orange	15 - 30 %
Red	> 30 %
Light Blue	Water
Grey	Unclassified



Land Resource Unit
Brandon Research Centre
April 1998

Universal Transverse Mercator
(UTM27) Projection

Surface Texture Map.


The soil textural class for the upper most soil horizon of the dominant soil series within a soil polygon was utilized for classification. Texture may vary from that shown with soil depth and location within the polygon.

Table 2. Surface Texture¹


Surface Texture	Area (ha)	Percent of RM
Organics	3004	3.1
Coarse Sands	472	0.5
Sands	1681	1.7
Coarse Loamy	125	0.1
Loamy	84436	86.5
Clayey	678	0.7
Eroded Slopes	4430	4.5
Marsh	0	0.0
Unclassified	0	0.0
Water	2805	2.9
Total	97631	100.0

¹ Based on the **dominant** soil series for each soil polygon.

Surface Texture Map



Surface Textures	
Light Green	Organic
Pink	Coarse Sands
Yellow	Sands
Orange	Coarse Loamy
Brown	Loamy
Dark Brown	Clayey
Grey	Unclassified
Purple	Marsh
Red	Eroded slopes
Cyan	Water



Universal Transverse Mercator
(UTM) Projection

Lane Resource Unit
Brandon Research Centre
April 1998

Soil Drainage Map.

Drainage is described on the basis of actual moisture content in excess of field capacity, and the length of the saturation period within the plant root zone. Six drainage classes plus four land classes are shown on this map.

Very Poor - Water is removed from the soil so slowly that the water table remains at or on the soil surface for the greater part of the time the soil is not frozen. Excess water is present in the soil throughout most of the year.

Poor - Water is removed so slowly in relation to supply that the soil remains wet for a large part of the time the soil is not frozen. Excess water is available within the soil for a large part of the time.

Imperfect - Water is removed from the soil sufficiently slowly in relation to supply to keep the soil wet for a significant part of the growing season. Excess water moves slowly down the profile if precipitation is the major source.

Well - Water is removed from the soil readily but not rapidly. Excess water flows downward readily into underlying materials or laterally as subsurface flow.

Rapid - Water is removed from the soil rapidly in relation to supply. Excess water flows downward if underlying material is pervious. Subsurface flow may occur on steep slopes during heavy rainfall.


Drainage classification is based on the dominant soil series within each individual soil polygon.

Table 3. Drainage Classes¹

Drainage Class	Area (ha)	Percent of RM
Very Poor	2931	3.0
Poor	7168	7.3
Imperfect	14502	14.9
Well	63643	65.2
Rapid	6583	6.7
Marsh	0	0.0
Unclassified	0	0.0
Water	2805	2.9
Total	97631	100.0


¹ Area has been assigned to the dominant drainage class for each soil polygon.

Soil Drainage Map



Drainage Classes

- Rapid
- Well
- Imperfect
- Poor
- Very poor
- Unclassified
- Marsh
- Water



Land Resource Unit
Brandon Research Centre
April 1998

Universal Transverse Mercator
(NAD83) Projection

Soil Salinity Map.

A saline soil contains soluble salts in such quantities that they interfere with the growth of most crops. Soil salinity is determined by the electrical conductivity of the saturation extract in decisiemens per metre (dS/m). Approximate limits of salinity classes are:

non-saline	< 4 dS/m
weakly saline	4 to 8 dS/m
moderately saline	8 to 15 dS/m
strongly saline	> 15 dS/m.


The salinity classification of each individual soil polygon was determined by the most severe salinity classification present within that polygon.

Table 4. Salinity Classes¹

Salinity Class	Area (ha)	Percent of RM
Non Saline	83785	85.8
Weakly Saline	6156	6.3
Moderately Saline	0	0.0
Strongly Saline	455	0.5
Eroded Slopes	4430	4.5
Marsh	0	0.0
Unclassified	0	0.0
Water	2805	2.9
Total	97631	100.0


¹ Area has been assigned to the most severe salinity class for each soil polygon.

Soil Salinity Map



Salinity Classes

- Strongly saline (>15 dS/m)
- Moderately Saline (8-15 dS/m.)
- Weakly saline(4-8 dS/m)
- Non saline (<4 dS/m.)
- Unclassified
- Marsh
- Eroded Slopes
- Water



Land Resource Unit
Brandon Research Centre
April 1998

Universal Transverse Mercator
(NAD27) Projection

Management Considerations Map.

Management consideration maps are provided to focus on awareness of land resource characteristics important to land use. This map does not presume a specific land use. Rather it portrays the most common and wide spread attributes that apply to most soil landscapes in the province.

These maps **highlight attributes** of soil-landscapes that the land manager must consider for any intended land use.

- Fine texture
- Medium texture
- Coarse texture
- Topography
- Wetness
- Organic
- Bedrock

F = Fine texture - soil landscapes with **fine textured soils (clays and silty clays)**, have low infiltration and internal permeability rates. These require special considerations to mitigate surface ponding (water logging), runoff, and trafficability. Timing and type of tillage practices used may be restricted.

M = Medium texture - soil landscapes with medium to moderately fine textures (**loams to clay loams**), and good water and nutrient retention properties. Good management and cropping practices are required to minimize leaching and the risk of erosion.

C = Coarse texture - soil landscapes with **coarse to very coarse textured soils (loamy sands, sands and gravels)** have a high permeability throughout the profile, and require special management practices related to application of agricultural chemicals, animal wastes, and municipal effluent to protect and sustain the long term quality of the soil and water resources. The risk of soil erosion can be minimized through the use of shelterbelts and maintenance of crop residues.

T = Topography - soil landscapes with **slopes greater than 5 %** are steep enough to require special management practices to minimize the risk of erosion.

W = Wetness - soil landscapes that have **poorly drained soils and/or >50 % wetlands** (due to seasonal and annual flooding, surface ponding, permanent water bodies (sloughs), and/or high water tables), require special management practices to mitigate adverse impact on water quality, protect subsurface aquifers, and sustain crop production during periods of high risk of water logging.

O = Organic - soil landscapes with organic soils, requiring special management considerations of drainage, tillage, and cropping to sustain productivity and minimize subsidence and erosion.


R = Bedrock - soil landscapes that have **shallow depth to bedrock (< 50 cm) and/or exposed bedrock** which may prevent the use of some or all tillage practices as well as the range of potential crops. They require special cropping and management practices to sustain agricultural production.

Table 5. Management Considerations¹

Land Resource Characteristics	Area (ha)	Percent of RM
Fine Texture	678	0.7
Fine Texture and Wetness	0	0.0
Fine Texture and Topography	0	0.0
Medium Texture	54036	55.3
Coarse Texture	262	0.3
Coarse Texture and Wetness	0	0.0
Coarse Texture and Topography	1891	1.9
Topography	27860	28.5
Topography and Bedrock	0	0.0
Wetness	9041	9.3
Wetness and Topography	0	0.0
Bedrock	0	0.0
Organic	1058	1.1
Marsh	0	0.0
Unclassified	0	0.0
Water	2805	2.9
Total	97631	100.0


¹ Based on **dominant** soil series for each soil polygon.

Management Considerations Map



Land Resource Characteristics

- | | |
|--|--|
|  Medium Texture |  Organic |
|  Coarse Texture |  Marsh |
|  Coarse texture and topography |  Wetness |
|  Coarse Texture and Wetness |  Unclassified |
|  Topography |  Water |
|  Fine texture and topography | |
|  Fine Texture | |
|  Fine texture and Wetness | |



Land Resource Unit
Brandon Research Centre
April 1998

Universal Transverse Mercator
(UTM) Projection

Agricultural Capability Map.

This evaluation utilizes the 7 class Canada Land Inventory system (CLI, 1965). Classes 1 to 3 represent the prime agricultural land, class 4 land is marginal for sustained cultivation, class 5 land is capable of perennial forages and improvement is feasible, class 6 land is capable of producing native forages and pasture but improvement is not feasible, and class 7 land is considered unsuitable for dryland agriculture. Subclass modifiers include structure and/or permeability (D), erosion (E), inundation (I), moisture limitation (M), salinity (N), stoniness (P), consolidated bedrock (R), topography (T), excess water (W) and cumulative minor adverse characteristics (X).

This generalized interpretive map is based on the dominant soil series and phases for each soil polygon. The CLI subclass limitations cannot be portrayed at this generalized map scale.

Table 6. Agricultural Capability¹


Class Subclass	Area (ha)	Percent of RM
1	8342	8.6
2	38774	39.9
2M	275	0.3
2T	24407	25.1
2TE	224	0.2
2TP	180	0.2
2TW	545	0.6
2W	11182	11.5
2WE	53	0.1
2X	1909	2.0
3	17021	17.5
3ET	141	0.1
3EW	80	0.1
3I	165	0.2
3M	5001	5.1

Table 6. Agricultural Capability¹(cont)

Class Subclass	Area (ha)	Percent of RM
3MT	1501	1.5
3N	1510	1.6
3T	7242	7.5
3TE	1167	1.2
3TP	215	0.2
4	11063	11.4
4D	687	0.7
4ET	466	0.5
4T	6125	6.3
4TE	3785	3.9
5	11552	11.9
5ET	195	0.2
5M	1601	1.6
5ME	193	0.2
5MT	130	0.1
5T	973	1.0
5TE	1377	1.4
5W	7082	7.3
6	6978	7.2
6EM	225	0.2
6ET	364	0.4
6W	4440	1.9
6WI	73	0.1
Water	2367	2.4
Organic	1052	1.1
Total	97149	100.0

¹ Based on **dominant** soil, slope gradient, and slope length of each soil polygon.


Agriculture Capability Map



Canada Land Inventory Classes

-  Class 1
-  Class 2
-  Class 3
-  Class 4
-  Class 5
-  Class 6
-  Class 7
-  Organic
-  Unclassified
-  Water

Land Resource Unit
Winnipeg Manitoba
June 2003



Universal Transverse Mercator
(NAD 83) Projection

Irrigation Suitability Map.

Irrigation ratings are based on an assessment of the most limiting combination of soil and landscape conditions. Soils in the same class have a similar relative suitability or degree of limitation for irrigation use, although the specific limiting factors may differ. These limiting factors are described by subclass symbols at detailed map scales. The irrigation rating system does not consider water availability, method of application, water quality, or economics of irrigated land use.

Irrigation suitability is a four class rating system. Areas with no or slight soil and/or landscape limitations are rated **Excellent** to **Good** and can be considered irrigable. Areas with moderate soil and/or landscape limitations are rated as **Fair** and considered marginal for irrigation providing adequate management exists so that the soil and adjacent areas are not adversely affected by water application. Soil and landscape areas rated as **Poor** have severe limitations for irrigation.


This generalized interpretive map is based on the dominant soil series for each soil polygon, in combination with the dominant slope class. The nature of the subclass limitations and the classification of subdominant components is not shown at this generalized map scale.

Table 7. Irrigation Suitability¹

Class	Area (ha)	Percent of RM
Excellent	0	0.0
Good	40619	41.6
Fair	34130	35.0
Poor	19019	19.5
Organic	1058	1.1
Unclassified	0	0.0
Water	2805	2.9
Total	97631	100.0


¹ Based on **dominant** soil, slope gradient, and slope length of each soil polygon.

Irrigation Suitability Map



R 12 R 11 R 10 R 9 R 8 W.
Irrigation Suitability Classes

-  Excellent
-  Good
-  Fair
-  Poor
-  Organic
-  Unclassified
-  Water



Land Resource Unit
Brandon Research Centre
April 1998

Universal Transverse Mercator
(NAD83) Projection

Potential Environmental Impact Under Irrigation Map.

A major environmental concern for land under irrigated crop production is the possibility that surface and/or ground water may be impacted. The potential environmental impact assessment provides a relative rating of land into 4 classes (minimal, low, moderate and high) based on an evaluation of specific soil factors and landscape conditions that determine the impact potential.

Soil factors considered are those properties that determine water retention and movement through the soil; topographic features are those that affect runoff and redistribution of moisture in the landscape. Several factors are specifically considered: soil texture, hydraulic conductivity, salinity, geological uniformity, depth to water table and topography. The risk of altering surface and subsurface soil drainage regimes, soil salinity, potential for runoff, erosion and flooding is determined by specific criteria for each property.

Use of this rating is intended to serve as a warning of potential environmental concern. It may be possible to design and/or give special consideration to soil-water-crop management practices that will mitigate any adverse impact.


This generalized interpretive map is based on the dominant soil series and slope class for each soil polygon. The nature of the subclass limitations, and the classification of subdominant components is not shown at this generalized map scale.

Table 8. Potential Environmental Impact Under Irrigation¹

Class	Area (ha)	Percent of RM
Minimal	5422	5.6
Low	52214	53.5
Moderate	12387	12.7
High	23746	24.3
Organic	1058	1.1
Unclassified	0	0.0
Water	2805	2.9
Total	97631	100.0

¹ Based on **dominant** soil, slope gradient, and slope length of each soil polygon.


Potential Environmental Impact Under Irrigation



Potential Impact Classes

- Minima
- Low
- Moderate
- High
- Organic
- Unclassified
- Water

Land Resource Unit
Brandon Research Centre
April 1998



Universal Transverse Mercator
(14°D27) Projection

Water Erosion Risk Map.

The risk of water erosion was estimated using the universal soil loss equation (USLE) developed by Wischmeier and Smith (1965). The USLE predicted soil loss (tons/hectare/year) is calculated for each soil component in each soil map polygon. Erosion risk classes are assigned based on the weighted average soil loss for each map polygon. The map shows 5 classes of soil erosion risk based on bare unprotected soil:


negligible
low
moderate
high
severe.

Cropping and residue management practices will significantly reduce this risk depending on crop rotation program, soil type, and landscape features.

Table 9. Water Erosion Risk


Class	Area (ha)	Percent of RM
Negligible	11439	11.7
Low	21985	22.5
Moderate	19347	19.8
High	11823	12.1
Severe	30232	31.0
Unclassified	0	0.0
Water	2805	2.9
Total	97631	100.0

Water Erosion Risk Map



Mean Risk Values

- Negligible
- Low
- Moderate
- High
- Severe
- Water



Land Resource Unit
Brandon Research Centre
April 1998

Universal Transverse Mercator
(NAD83) Projection

Land Use Map.

The land use classification of the RM has been interpreted from LANDSAT satellite imagery, using supervised computer classification techniques. Many individual spectral signatures were classified and grouped into the seven general land use classes shown here. Although land use changes over time, and some land use practices on individual parcels may occasionally result in similar spectral signatures, this map provides a general representation of the current land use in the RM.

The following is a brief description of the land use classes:

Annual Crop Land - land that is normally cultivated on an annual basis.

Forage - perennial forages, generally alfalfa or clover with blends of tame grasses.

Grasslands - areas of native or tame grasses, may contain scattered stands of shrubs.

Trees - lands that are primarily in tree cover.

Wetlands - areas that are wet, often with sedges, cattails, and rushes.

Water - open water - lakes, rivers streams, ponds, and lagoons.

Urban and Transportation - towns, roads, railways, quarries.

Table 10. Land Use¹

Class	Area (ha)	Percent of RM
Annual Crop Land	66303	67.4
Forage	3618	3.7
Grasslands	11715	11.9
Trees	8114	8.3
Wetlands	2973	3.0
Water	2264	2.3
Urban and Transportation	3362	3.4
Total	98349	100.0

¹ Land use information (1995) and map supplied by Prairie Farm Rehabilitation Administration. Areas may vary from previous maps due to differences in analytical procedures.


RURAL MUNICIPALITY OF LORNE LAND USE

Reference:

Satellite imagery obtained from RSI
1995.
Landsat TM (30m pixel resolution)
Date of image: Annual Crop Land
and Forage May 24, 1994. All
other classes: May 9, 1988.
Classification from Manitoba Remote
Sensing Centre.

LAND USE

- Annual Crop Land
- Timber
- Water
- Grassland
- Wetlands
- Forage
- Urban & Transportation
- Municipal Boundary



REFERENCES

Agronomic Interpretations Working Group. 1995. Land Suitability Rating System for Agricultural Crops: 1. Spring-seeded Small Grains. Edited by W.W. Pettapiece. Tech. Bull. 1995-6E. Centre for Land and Biological Resources Research, Agriculture and Agri-Food Canada, Ottawa. 90 pages, 2 maps.

Ash, G.H.B. 1991. An Agroclimatic Risk Assessment of Southern Manitoba and Southeastern Saskatchewan. M.A. Thesis. Department of Geography, University of Manitoba, Winnipeg.

Canada Land Inventory. 1965. Soil Capability Classification for Agriculture. Canada Land Inventory Report No. 2. ARDA, Dept. of Forestry, Canada, Ottawa.

Canada-Manitoba Soil Survey. 1980. Physiographic Regions of Manitoba. Ellis Bldg., University of Manitoba, Winnipeg. Revised. Unpublished Report.

Canada-Manitoba Soil Survey. 1979. Ecological Regions and Subregions in Manitoba. Ellis Bldg., University of Manitoba, Winnipeg. Revised. Unpublished Report.

Environment Canada. 1982. Canadian Climatic Normals 1951-1980. Frost, Vol. 6; Atmospheric Environment, Downsview, Ontario.

Environment Canada. 1993. Canadian Climatic Normals 1961-1990. Prairie Provinces. Atmospheric Environment, Downsview, Ontario.

Expert Committee on Soil Survey. 1987. The Canadian System of Soil Classification. Second Edition. Publ. No. 1646. Research Branch, Agriculture Canada.

Irrigation Suitability Classification Working Group. 1987. An Irrigation Suitability Classification System for the Canadian Prairies. LRRC contribution no. 87-83, Land Resource Research Centre, Research Branch, Agriculture Canada, Ottawa

Langman, M.N., 1986. Soils of the Rural Municipality of Lorne. Report No. D70. Canada-Manitoba Soil Survey. Winnipeg.

MacDonald, K.B., and Valentine, K.W.G. 1992. CanSIS Manual 1 CanSIS/NSDB: A General Description. Land Resource Division, Centre for Land and Biological Resources Research, Research Branch, Agriculture Canada, Ottawa.

Manitoba Land Resource Unit. 1998. Soil and Terrain Classification System Manual. In preparation. Ellis Bldg. University of Manitoba. Winnipeg.

Wischmeier, W.H. and Smith, D.D. 1965. Predicting Rainfall-erosion Loss from Cropland East of the Rocky Mountains. U.S. Department of Agriculture, Agriculture Handbook No. 282, U.S. Government Printing Office, Washington, D.C.