


THE SOILS OF
PLEASANTDALE

RURAL MUNICIPALITY No. 398

SASKATCHEWAN



Saskatchewan Centre for Soil Research
Publication SM398

Centre For Land and Biological
Resources Research
Land Resources Division,
Agriculture Canada
Publication 91-49

THE SOILS OF
PLEASANTDALE
RURAL MUNICIPALITY No. 398
SASKATCHEWAN

MAY 1, 1995

by

Staff, Saskatchewan Soil Survey
SASKATCHEWAN CENTRE FOR SOIL RESEARCH
University of Saskatchewan



Saskatchewan
Agriculture
And Food

Agriculture
Development
Fund



Canada-Saskatchewan
Agreement
On Soil Conservation



Agriculture
Canada

Research
Branch

STAFF, SASKATCHEWAN SOIL SURVEY

Map and Report Compilation: M.D. Bock

Program Directors: D.W. Anderson, Director, H.P.W. Rostad, Associate Director, Saskatchewan
Centre for Soil Research, Saskatoon

Soil Correlation: H.B. Stonehouse

Project Supervisors: H.B. Stonehouse and L.M. Kozak

Soil Mapping: M.D. Bock, D.D. Cerkowniak, L.G. Fuller, M. Samadi, C.T. Stushnoff, R. D. Stushnoff,
and F. Youatt

Secretarial: S.S.M. Wood

Drafting: P.M. Sardinha

Geographical Information System: S.R.E. Johnson

Laboratory: B.G. Goetz and J.D. Key

Students: R. Gares, D. Horn, R. McBride, and Y. Morrissette

TABLE OF CONTENTS

1. INTRODUCTION.....	1-1
1.1 Using the Soil Map and Report	1-1
2. INTRODUCTION TO SOILS	2-1
2.1 The Soil Profile	2-1
2.2 The Soil Map	2-1
2.3 Surface Deposits	2-1
2.4 Surface Forms	2-2
2.5 Surface Texture	2-3
3. DESCRIPTION OF SOILS.....	3-1
4. SOIL INTERPRETATIONS	4-1
4.1 Salinity	4-1
4.2 Irrigation Suitability	4-3
4.3 Stones	4-5
4.4 Soil Capability for Agriculture	4-6
4.5 Surface pH.....	4-8
4.6 Wetlands Classification	4-9
4.7 Wind Erosion	4-10
4.8 Water Erosion	4-11
4.9 Past Wind and Water Erosion	4-12
4.10 Sand and Gravel	4-13
4.11 Soil Moisture and Yield	4-14
5. ACREAGE FACTS	5-1
6. INTERPRETIVE DATA TABLES	6-1
7. GLOSSARY	7-1
8. FURTHER INFORMATION	8-1
KEY MAP	8-2
MAP (in pocket)	

1. INTRODUCTION

This publication continues the series of soil survey reports for Saskatchewan initiated on an R.M. basis in 1984. This series of publications is a continuation of the basic soil survey program in the Province, initiated in 1958; however, the publication format has been substantially changed to include more interpretive information on an R.M. basis.

The main purpose of a soil survey is to inventory the soil resources of an area, providing a description of the soils and showing their extent and distribution. It has become increasingly apparent, however, that many users require additional interpretive information for the resolution of production, conservation and other problems related to a particular set of soil conditions. To that end, this report also presents a number of interpretations based on the soil inventory information.

In order to gain the most information about any particular area within the municipality, both the soil map and report must be used together.

1.1 USING THE SOIL MAP AND REPORT

Each delineation on the soil map contains a map symbol and a unique number which are described and illustrated below.

The Soil Map Symbol

The map symbol is made up of a soil association code, a map unit number, and in some cases a substrate modifier code, along with a surface texture code in the numerator and a code composed of numbers and letters indicating the slope class and surface form of the landscape in the denominator. A brief explanation of each of these map symbol components is provided in the legend on the side of the map. The legend describes the general type of soil development, the geologic material in which each soil has developed, and in complex areas, where each geologic material occurs in the landscape, as well as the kinds of soils comprising each map unit.


Delineation Number and Soil Interpretations

Each map delineation contains a unique number which is used to reference additional soils and interpretive information in Section 6 of the report. This section provides a tabular listing of interpretive symbols for each delineation. An explanation of these symbols is provided under the appropriate subsection in Section 4 entitled "Soil Interpretations."

Example

To determine the agricultural capability classification for area 12 (used in the example below), turn to Section 6 and look up the number 12 listed in the left-hand column under the heading, "Area No.". Next, read across to the symbols listed in the column headed, "Agricultural Capability". These symbols are explained in Subsection 4.4, entitled, "Soil Capability for Agriculture".

Figure 1. Sequence of Symbols.



2. INTRODUCTION TO SOILS

The nature and agriculturally important properties of the soils of the area are described in succeeding sections of this report. The present section, largely adapted from H.C. Moss, in *A Guide to Understanding Saskatchewan Soils*, deals mainly with features common to most prairie soils.

2.1 THE SOIL PROFILE

A soil is a natural body that occupies a relatively thin section (usually less than a meter) of the earth's surface and consists of several layers or horizons which differ in appearance and composition from the underlying material. Its formation from the original geological deposit involves various physical, chemical and biological processes which result in the formation of individual layers or horizons, extending from the surface downwards, that have specific characteristics. The whole succession of layers down to and including the original geological deposit is called the **soil profile**. Each individual layer is called a **soil horizon**. A particular soil is recognized and separated from other soils by identifying the various layers or horizons which make up its profile. The recognition of soil profiles forms the basis of soil classification and mapping.

The soils of Saskatchewan are classified according to a national system of soil classification and the names given to the soils are derived, in part, from this system. For example, an orthic profile is a soil whose characteristics are defined as an Orthic Chernozemic soil of the National system.

In profiles of mineral soils, three main horizons are recognized. From the surface downward, these are designated by the letters A, B, and C. The A horizon forms all or part of the surface soil. It may be dark colored representing an accumulation of humus, or it may be a light-colored horizon from which clay, humus and other materials have been removed. The B horizon occurs immediately below the A horizon. It may have an accumulation of clay and may have been altered to give a change in color or structure. The C horizon occupies the lower portion of the soil profile and usually represents the parent material. It is relatively unaffected by soil forming processes operative in the A and B horizon.

2.2 THE SOIL MAP

Ideally, the area represented by each soil profile should be shown on the map. This, however, is only possible where large, uniform areas of a single soil occur, or in detailed soil surveys where small areas can be separated on the map. Since, on the semi-detailed maps, it is rarely possible to delineate areas of a single soil, it is almost always necessary

to combine small areas of several soils into a larger area. These larger areas are represented on the map by a map unit that identifies the kinds and distribution of the component soil profiles.

The **soil association** is used to show the relationship between map units that have formed on a similar geological deposit within a particular soil zone. The **Oxbow Association**, for instance, is the name given to a group of soil profiles formed on loamy glacial till occurring in the black soil zone. The various **map units** of the Oxbow Association reflect variations in the kind and distribution of Oxbow soils from one area to the next.

Where two geological deposits occur within a delineated area on the map, two associations are used. As an example, Meota-Oxbow is the name given to a group of soils of the Meota and Oxbow soil associations. Different map units of this complex are used to reflect variations in the kind and distribution of Meota and Oxbow soils from one area to the next. As an exception, areas in which several geological deposits occur in a somewhat chaotic and unpredictable pattern throughout the landscape are often given a single association name. For example, Keppel is the name given to soils formed in a highly complex mixture of loamy glacial till, silty water-modified glacial till and silty glaciolacustrine materials.

It is possible also to find soils reflecting the characteristics of two soil zones within a local area. Under these circumstances, two associations are used to reflect these different soil properties. For example, Black and Dark Gray soils that occur together are mapped in the Oxbow-Whitewood complex, the Oxbow referring to the Black soils and Whitewood to the Dark Gray soils.

The soil map, then, attempts to portray the kinds and distribution of various soil profiles throughout the municipality. The symbols on the map identify the soil map unit, the soil texture, the slope class and surface form. The map legend provides a brief description of these features. More complete descriptions of individual soil associations and their component soil types are provided in the Description of Soils section of the report. The types of geological deposits which comprise the parent materials of the various soil associations and the surface forms or shape of the land are described below.

2.3 SURFACE DEPOSITS

Alluvial Deposits - Alluvial deposits are materials laid down by streams and rivers, in valley bottoms and collection basins, since glaciation. These deposits are stratified and often contain beds or layers that are oblique to the main planes of stratification, indicative of their river or stream origin.

Eolian Deposits - Eolian deposits are sandy or silty deposits that have been moved and redeposited by the wind, often in the form of sand dunes or silty loessial veneers or blankets. Eolian deposits are well-sorted, poorly compacted and may contain beds or layers.

Fluvial Deposits - Fluvial deposits are materials laid down in rivers and streams carrying glacial meltwater. They are usually sandy or gravelly and, like the alluvial deposits described above, may contain beds or layers that are inclined or oblique to the main planes of stratification. These deposits are usually thick but may be thin, like a veneer, and underlain by glacial till. Materials laid down in direct contact with the glacier are termed *glaciofluvial*.

Lacustrine Deposits - Lacustrine deposits are materials laid down in a glacial lake. These deposits are often stratified and characterized by dark- and light-colored beds or layers reflecting summer and winter depositional cycles in a glacial lake. Lacustrine deposits usually have a high content of very fine sand-, silt- or clay-sized particles. Those dominated by sand-sized particles are termed *loamy lacustrine* while those dominated by silt- and clay-sized particles are termed *silty* and *clayey lacustrine*, respectively. They are usually thick but may be thin, like a veneer, and underlain by glacial till or gravel. Materials laid down in close contact with the glacier are termed *glaciolacustrine* deposits.

Morainal Deposits - Morainal deposits, often referred to as glacial till, are materials laid down by the glacial ice. These deposits are generally comprised of stones and gravels embedded in a matrix of sand-, silt- and clay-sized materials. When this matrix contains nearly equal amounts of sand, silt and clay they are called *loamy morainal* deposits. When there is a preponderance of sand or silt, they are referred to as *sandy morainal* or *silty morainal* deposits, respectively. Usually, there are fewer stones and gravels present in silty morainal deposits than in sandy or loamy types. Morainal deposits characterized by an abundance of surface stones are called *bouldery morainal* deposits.

Organic Deposits - Organic deposits are materials laid down by the accumulation of plant remains. They are generally 40 cm thick or greater and are comprised of either the remains of mosses or sedges and grasses and often have inclusions of woody materials. When the organic materials are largely undecomposed, so that there is a large amount of well-preserved fiber that is readily identifiable as to botanical origin, they are called *fibric organic* deposits. When the organic materials are in an intermediate stage of decomposition, so that there is an intermediate amount of fiber that is identifiable as to botanical origin, they are called *mesic organic* deposits. Highly decomposed materials, which have a small amount of fiber that can be identified as to botanical origin, are called *humic organic* deposits.

Undifferentiated Deposits - Areas where the origin of the materials for the purpose of mapping has not been specified are termed undifferentiated deposits. These deposits, usually consisting of several materials (morainal, fluvial, lacustrine, or others) occur in areas of steeply sloping land such as coulees and valley sides.

2.4 SURFACE FORMS

Aprons and Fans - A fan is a gently sloping fan-shaped area, usually occurring at the base of a valley wall, resulting from the accumulation of sediments brought down by a stream descending through a steep ravine. A series of adjacent, coalescing fans is called an apron.

Hummocky - Landscapes with a complex pattern of generally short, steep slopes extending from prominent knolls to somewhat rounded depressions or kettles are termed hummocky. They are called *hummocky dissected* where shallow gullies join one low area or kettle to the next and *hummocky gullied* where numerous, parallel or subparallel, narrow ravines interrupt the hummocky features of the landscape. Occasionally, areas have a complex of ridged and hummocky features. They are called *hummocky-ridged*.

Inclined - Landscapes in which the general slope is in one direction only are called inclined. Where shallow gullies occur along the slope, the areas are called *inclined dissected*; where a series of parallel or subparallel, deep gullies or ravines occur, they are called *inclined gullied*.

Level - Landscapes that are flat or have very gently sloping surfaces are said to be level. Along flood plains of rivers and streams where the level surface is broken by abandoned river channels they are called *level channelled*.

Ridged - Landscapes that have a linear pattern, usually of short and straight parallel ridges but sometimes a single, sinuous ridge or a series of intersecting ridges are termed ridged.

Rolling - Landscapes that are characterized by a sequence of long (often 1.6 km or greater), moderate to strong slopes extending from rounded, sometimes confined depressions to broad, rounded knolls, that impart a wave-like pattern to the land surface are called rolling. They are called *dissected rolling* where shallow gullies join one low area or kettle to the next.

Terraced - Areas, usually along a valley, that have a steep, short scarp slope and a horizontal or gently inclined surface above it are called terraced.

Undulating - Landscapes that are characterized by a sequence of gentle slopes extending from smooth rises to gentle hollows, that impart a wavelike pattern to the land surface are called undulating. Where shallow gullies extend from one low area to the next in these landscapes they are called *undulating dissected* and where the undulating surface is broken by abandoned river channels they are called *undulating channelled*.

2.5 SURFACE TEXTURE

Mineral soil is a mixture of various-sized mineral particles, decaying organic matter, air and water. The mineral particles, exclusive of stones and gravel, may be grouped into three particle-size fractions: sands (soil particles between 0.05 and 2 mm in diameter), silts (soil particles between 0.002 and 0.05 mm in diameter), and clays (soil particles less than 0.002 mm in diameter). The relative proportions of these particle-size fractions in a soil determine its texture. The textural triangle (Figure 2) is used to illustrate the proportion of sand, silt and clay in the main textural classes. The vertical axis is percent clay, the horizontal axis is percent sand, while the remainder of each class is percent silt. Thus, when sand is dominant, it yields a sandy- or coarse-textured soil, whereas a fine-textured soil is made up largely of silt and clay. The terms "light" and "heavy" are often used to refer to sandy- and clayey-textured soils respectively, and are actually a measure of the power required to till the soil. These terms have nothing to do with the actual weight of soil, as a given volume of dry sand actually weighs slightly more than that of clay.

Textural class names such as sandy loam, clay loam, heavy clay, etc., are given to soils based upon the relative proportions of sand, silt and clay. Three broad, fundamental textural groups are recognized: sands, loams and clays.

SANDS - The sand group includes soils in which the sand particles make up at least 70% of the material by weight. Two main classes are recognized: sand and loamy sand. Sands are further broken down into different sand sizes such as fine sand or coarse sand. A description of these is found under "Sand" in the glossary.


LOAMS - The loam group is intermediate in texture between the coarse-textured sands and the fine-textured clays, and these soils usually contain a significant proportion of each particle-size fraction. Class names include: sandy loam, silt loam, silty clay loam, sandy clay loam, clay loam and loam.

CLAYS - The clay group includes soils that contain at least 35% clay-size particles, and in most cases, more than 40%. Class names are: sandy clay, silty clay, clay and heavy clay. Soils of this group are often referred to as "gumbo".

Table 1. Soil texture classes.

Symbol	Soil Texture Class
Coarse-Textured	
gs	Gravelly sand
s	Sand
fs	Fine sand
gls	Gravelly loamy sand
ls	Loamy sand
lfs	Loamy fine sand
Moderately Coarse-Textured	
gsl	Gravelly sandy loam
gl	Gravelly loam
sl	Sandy loam
fl	Fine sandy loam
vl	Very fine sandy loam
Medium-Textured	
scl	Sandy clay loam
fcl	Fine sandy clay loam
vcl	Very fine sandy clay loam
l	Loam
Moderately Fine-Textured	
sil	Silt loam
cl	Clay loam
sicl	Silty clay loam
Fine-Textured	
c	Clay
sic	Silty clay
hc	Heavy clay
Miscellaneous	
o	Organic
U	Unclassified

Table 1 lists the surface textures and symbols that may be used in this report, grouped into particle-size categories. The miscellaneous category contains two non-texture entries. The "o" or organic soil texture class is used for organic soils. By definition, these soils do not contain any mineral component and, therefore, do not have a surface texture as defined and described above. The symbol "o" merely identifies the surface as being organic. The "U" or unclassified class is used for areas in which surface texture has not been determined. These include areas that have been greatly altered (such as gravel pits or mines), most wetlands and lakes, areas that have not been examined (such as towns and cities), and areas of extremely variable texture (such as some Hillwash or Runway delineations).



3. DESCRIPTION OF SOILS

ALLUVIUM (Ay) SOILS

Alluvium soils are a mixture of soils formed in variable-textured alluvial materials, associated with stream floodplains and drainage channels. These soils have formed in materials derived from a variety of sources and thus vary markedly in color, texture, and composition. Surface textures range from sand to clay.

Alluvium soils are usually stone free, but may be underlain by stony deposits. Eroded stream beds within Alluvium areas, for example, may be very stony. Alluvium soils usually occur on level or undulating landscapes with very gentle to gentle slopes.

Alluvium soils occur in complex with soils of other associations and, in most of these complexes, the Alluvium soils occur on the lower slopes in the landscape.

Kinds of Alluvium Soils

Poorly Drained Alluvium - Poorly drained Alluvium soils represent a variety of wet soils. They occur in undrained depressional areas that are subject to flooding and in undrained areas associated with abandoned stream meanders. They often have thick, dark-colored A horizons and drab subsurface colors that are dotted with reddish spots and streaks. Some of these soils have a very light-gray leached horizon below the dark A horizon. Peaty poorly drained Alluvium soils have a layer of peaty material, 15 to 40 cm thick, overlying mineral materials. These soils are frequently carbonated to the surface (carbonated poorly drained) and, in many cases, also saline (saline poorly drained), with soluble salts generally present within 50 cm of the surface. They are generally wet for all or a significant portion of the growing season and are often flooded. Most poorly drained Alluvium soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. In some areas, these soils may be saline and/or carbonated.

Agricultural Properties of Alluvium Soils

The agricultural capability of Alluvium soils ranges from class 2, good agricultural soils, to class 7, soils with no capability for arable agriculture or permanent pasture. This wide range in agricultural capability is mainly the result of varying degrees of salinity and excess wetness. Where salinity and wetness are not a problem and where the soil texture provides an adequate water-holding capacity, they are often good agricultural soils. Alluvium soils are often downrated based on soil and landscape limitations (i.e. salinity, excess wetness, erosion, etc.) that are peculiar to individual delineations. Limitations for each delineation are

listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Alluvium soils generally have a moderate amount of organic matter in the A horizon, resulting in reasonable fertility and good tilth. Because of the association of these soils with lower portions of the landscape and their occurrence in narrow bands in valley bottoms, wind erosion is not usually a serious problem. They are, however, susceptible to water erosion and to flooding because of their association with low landscape positions and drainage channels which receive runoff water in the spring or during periods of intense rainfall. They usually occur on favorable topography but areas are often small and irregular and cut by erosion channels, making cultivation difficult or impractical.

Stones are not generally a problem, however, annual clearing may be required along eroded stream beds. Areas dominated by saline or poorly drained Alluvium soils generally have little potential for crop land and are suitable mainly for forage production or pasture. If drainage can be improved and if salinity and accessibility do not pose serious limitations, some of these areas can be brought into agricultural production.

BLAINE LAKE (Bb) SOILS

Blaine Lake soils are Black soils that have formed in moderately calcareous, silty lacustrine materials. Surface textures are usually loam to silty clay loam.

Blaine Lake soils are typically stone free, however, some stones may occur where the lacustrine sediments are shallow (less than 1 m thick) and underlain by glacial till or gravel. Blaine Lake soils occur extensively on undulating landscapes with nearly level to gentle slopes and also on hummocky landscapes with gentle to moderate slopes.

Blaine Lake soils frequently occur in complex with soils of other associations. In most of these complexes, the Blaine Lake soils tend to occur on the mid- and lower slope positions.

Kinds of Blaine Lake Soils

Orthic Blaine Lake - The orthic Blaine Lake soil occurs on mid- to lower slopes in most Blaine Lake landscapes, but may extend to upper slopes on landscapes with gentle slopes. It is a well-drained soil characterized by a black A horizon, 13 to 18 cm thick, underlain by a brownish-colored, lime-free B horizon and a grayish-colored, moderately calcareous C horizon.

Carbonated Blaine Lake - The carbonated Blaine Lake soil occurs on mid- to lower slopes, frequently surrounding sloughs or poorly drained depressions, and on imperfectly poorly drained flat or depressional areas. It is an imperfectly drained soil with a black, usually calcareous A horizon, underlain by a highly calcareous B or C horizon. The B and C horizons often have dull colors and reddish spots and stains, indicative of imperfect soil drainage. Streaks of carbonate frequently occur throughout the upper horizons.

Saline Blaine Lake - The saline Blaine Lake soil occurs on mid- to lower slopes, often surrounding sloughs or poorly drained depressional areas, and along drainage channels or gullies. It is characterized by the presence of soluble salts, usually within 50 cm of the surface. The salts occur as a white surface crust or as small, white specks within the soil, although salts may not always be visible. Dull colors and reddish spots and stains, indicative of imperfect soil drainage, are often present in the subsoil. It frequently occurs intermixed with carbonated Blaine Lake soils on lower slopes surrounding sloughs.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Blaine Lake Soils

The Blaine Lake soils with clay loam or silty clay loam surface textures are very good agricultural soils of capability class 1. Most Blaine Lake soils have loam or silt loam surface textures. These are considered to be good agricultural soils of capability class 2. A slight moisture deficit, imparted by the subhumid regional climate and a moderate water-holding capacity, is the main limitation to the production of common field crops. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Blaine Lake soils have a high amount of organic matter in the A horizon. They are low in available phosphorus and high in available potassium. Crops grown on these soils usually respond to additions of nitrogen and phosphorus fertilizers. Under favorable moisture conditions, the majority of Blaine Lake soils often result in some of the best crops produced in the area. Wet conditions may be prevalent on the imperfectly drained soils in the spring, therefore, cultivation and seeding operations may be delayed. Drying of the surface of carbonated Blaine Lake soils in the spring can produce a light and fluffy consistency, which may make preparation of a good seed bed difficult. Blaine Lake soils that occur in landscapes with gentle slopes have a low susceptibility to wind and water erosion. Stones are gener-

ally not a problem, but where the soils are shallow and underlain by eroded glacial till, occasional clearing may be required.

BODMIN (Bd) SOILS

Bodmin soils are Gray soils that have formed in gravelly fluvial materials where wooded vegetation has had a strong influence on soil formation. Soils that have formed under these conditions are usually strongly leached, resulting in low organic matter levels and, hence, a grayish-colored surface when cultivated. Surface textures range from loamy sand to loam or gravelly loamy sand to gravelly loam, depending on the nature of the parent material.

Bodmin soils are usually slightly to moderately stony, however, where the gravelly deposit is thin and underlain or mixed with glacial till, surface stones are often more numerous. These soils occur on a variety of landscapes. Undulating landscapes with very gentle to gentle slopes and hummocky landscapes with gentle to moderate slopes are two of the more common landforms.

Bodmin soils frequently occur in complex with soils of other associations. In most of these complexes, the Bodmin soils occur randomly throughout the landscape.

Kinds of Bodmin Soils

Gray Wooded Bodmin - The gray wooded Bodmin soil usually occurs on upper slopes and knolls, but may extend to lower slopes in some landscapes. It is a well- to rapidly drained soil that, when cultivated, has a very light grayish-colored A horizon, underlain by a dense, brownish-colored B horizon with prismatic structure that breaks to angular blocky or coarse angular aggregates when dry. The B horizon, in turn, is underlain by a grayish-brown, weakly calcareous C horizon. In the native state, these soils usually occur under aspen forest, where the uppermost horizons consist of a layer of leaf litter and a very thin, dark-colored A horizon overlying a very light grayish-colored, leached horizon with platy structure. Upon cultivation, these horizons become mixed, resulting in the very light grayish-colored surface.

Dark Gray Wooded Bodmin - The dark gray wooded Bodmin soil usually occurs on mid- and upper slope positions, but may extend to lower slopes in some landscapes. It is a well- to rapidly drained soil with a grayish-colored A horizon, 8 to 13 cm thick, underlain by a dense dark-brown to reddish-brown B horizon with a weak structure that breaks to rough, irregular, cloddy aggregates when dry. The B horizon, in turn, is underlain by a grayish-brown, weakly calcareous C horizon. Most of these soils, and in particular those that occur under native vegetation, have a light-gray horizon with platy structure separating the surface horizon from the B horizon. Part of this horizon may remain intact below the plow layer following cultivation.

Agricultural Properties of Bodmin Soils

Bodmin soils are poor agricultural soils of capability class 4. Their coarse textures, resulting in a very low to low water-holding capacity and, consequently, poor drought resistance, as well as their low inherent fertility, are their main limitations. These soils may be downrated further based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. For example, some Bodmin soils have been rated as class 5 (unsuitable for sustained agriculture) because of their coarse surface textures and excessive amounts of stone. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

These soils are low in organic matter. They usually require applications of nitrogen and phosphorus fertilizers and may also need additions of potassium and sulphur, depending on the crops grown. Stones are usually present, however, amounts vary considerably. Some areas are almost stone free, while others require periodic removal if the soils are to be cultivated. Due to their coarse textures and low organic matter contents, wind erosion may present a further problem, especially on the steeper slopes and knolls. Conservation practices that maintain a continual crop or trash cover should be used to protect these soils. Although forage crops have been grown successfully on some of the Bodmin soils, they are generally considered to be marginal to unsuitable for annual crop production.

BREDBURY (Bu) SOILS

Bredbury soils are Thick Black soils that have formed in loamy lacustrine materials in moist areas of the Black soil zone, where soils with thicker-than-normal surface horizons have formed. These soils are distinguished from Hamlin soils (thin Black soils formed in loamy lacustrine materials) by having a surface horizon that is greater than 20 cm thick. Surface textures range from sandy loam to loam.

Bredbury soils are usually stone free. However, some stones may occur where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel, or where the lacustrine materials occur in complex with glacial till or gravel. Bredbury soils usually occur on undulating landscapes with very gentle to gentle slopes, however, they can also occur on hummocky landscapes with gentle to moderate slopes, particularly where they occur in complex with soils formed in glacial till.

The Bredbury soils frequently occur in complex with soils of other associations. In most of these complexes, the Bredbury soils occur on the mid- and lower slopes.

Kinds of Bredbury Soils

Orthic Bredbury - The orthic Bredbury soil occurs on mid- and lower slopes, however, it can extend onto upper slopes in some landscapes. It is a well-drained soil

characterized by a black A horizon, 21 to 33 cm thick, underlain by a brownish-colored B horizon and a grayish-brown, moderately calcareous C horizon.

Gleyed Orthic Bredbury - The gleyed orthic Bredbury soil occurs on mid- to lower slopes in some landscapes. It is a moderately well- to imperfectly drained soil characterized by a black A horizon, 23 to 41 cm thick, underlain by B and C horizons that often have dull colors and reddish spots and stains, indicative of restricted soil drainage.

Carbonated Bredbury - The carbonated Bredbury soil usually occurs on lower slopes, however, it can occur on all slope positions in some landscapes, and is affected to varying degrees by imperfect soil drainage. It is characterized by a highly calcareous A horizon underlain by a highly calcareous B or C horizon. The B and C horizons often have drab colors and reddish spots and stains, indicative of imperfect soil drainage. In addition, the subsoil is often affected by salinity to some degree.

Saline Bredbury - The saline Bredbury soil occurs on lower slopes, often surrounding sloughs or poorly drained depressional areas, and along drainage channels and gullies, however, it can occur on all slope positions in some landscapes. It is characterized by the presence of soluble salts, usually within 50 cm of the surface. The salts occur as a white surface crust or as small, white specks within the soil, although salts may not always be visible. Dull colors and reddish spots and stains, indicative of imperfect soil drainage, are often present in the subsoil. It frequently occurs intermixed with carbonated Bredbury soils.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Bredbury Soils

The best Bredbury soils are those with a loam to fine sandy loam surface texture. These soils have an agricultural capability rating of class 2 (good); a moderate moisture deficit, imparted by a moderate water-holding capacity and the subhumid regional climate, is their main limitation. Bredbury soils with sandy loam surface textures have been rated as capability class 3 (fair) due primarily to their lower water-holding capacity. Although many of the Bredbury soils have few agricultural limitations, some have been downrated based on other soil and landscape limitations (i.e.

salinity, topography, stones, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Most Bredenbury soils have a moderately high organic matter content in the A horizon, resulting in soils of reasonably good tilth. They are reasonably fertile soils, although additions of phosphorus and nitrogen are usually required to obtain optimum yields. Bredenbury soils have a weak to moderate cloddy structure that breaks to fine granular and single grain, making them moderately susceptible to wind erosion. Relatively high infiltration rates, coupled with very gentle to moderate slopes, result in a relatively low susceptibility to water erosion. It is recommended that soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, inclusions of forages in crop rotations and shelterbelts, be utilized wherever possible to control soil erosion and maintain or enhance soil organic matter content and aggregate stability. Stones are not generally a problem, however, periodic clearing may be required where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel.

CUT KNIFE (Cf) SOILS

Cut Knife soils are Black soils that have formed in shallow (less than 1 m thick), silty lacustrine materials underlain by glacial till. Surface textures are predominantly loam and silt loam, but can range to silty clay loam and clay loam.

Cut Knife soils are usually slightly stony to stone free, but may be moderately stony in areas where the lacustrine materials are very shallow, or where they occur in complex with glacial till or gravel. Cut Knife soils are typically associated with undulating or undulating dissected landscapes with very gentle to gentle slopes, but may also occur on hummocky landscapes with stronger slopes, particularly where they occur in complex with soils formed in glacial till.

Cut Knife soils frequently occur in complex with soils of other associations. In most of these complexes, the Cut Knife soils occur on the mid- and lower slope positions.

Kinds of Cut Knife Soils

Orthic Cut Knife - The orthic Cut Knife soil may occupy all slope positions. It is a well-drained soil characterized by a black A horizon, 12 to 18 cm thick, underlain by a brownish-colored, lime-free B horizon and a grayish-colored, moderately calcareous C horizon.

Calcareous Cut Knife - The calcareous Cut Knife soil usually occurs on upper slopes and knolls. It is a well-drained soil characterized by a black, often calcareous A horizon, 12 to 18 cm thick, underlain by a calcareous B horizon and a grayish-colored, moderately calcareous C horizon.

Agricultural Properties of Cut Knife Soils

The Cut Knife soils with clay loam and silty clay loam surface textures are very good agricultural soils of capability class 1. Cut Knife soils with silt loam and loam surface textures are rated as agricultural capability class 2 (good). A slight moisture deficit, imparted by a moderate water-holding capacity and the subhumid regional climate, is their main limitation. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. Ratings for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Most Cut Knife soils have a high amount of organic matter in the A horizon, resulting in reasonably fertile soils of good tilth. These soils are low in available phosphorus and high in available potassium. Crops grown on these soils usually respond to additions of nitrogen and phosphorus fertilizers.

Stones are seldom a problem on most Cut Knife soils, however, due to the presence of glacial till near the surface, occasional clearing may be required. Similarly, water erosion is seldom a problem, but can become serious during periods of intense rainfall due to the relatively low infiltration rate of these soils. The potential for water erosion is greatest in areas with long slopes and management practices, such as cultivation across slopes and grassing of runways, should be followed as much as possible. Wind erosion can be a serious problem unless conservation practices are followed. Such practices include the maintenance of crop residues through reduced tillage or leaving stubble standing, continuous cropping, or the establishment of forages in seriously affected areas.

CARROT RIVER (Cr) SOILS

Carrot River soils are Gleyed Dark Gray soils that have formed in sandy fluvial materials, in areas of mixed grassland and forest, where wooded vegetation and restricted soil drainage have had an influence on soil formation. Soils formed under wooded vegetation are usually slightly leached. They have lower organic matter levels than soils occurring in the Black soil zone, resulting in a dark-gray surface horizon. Although cultivation may have improved surface and internal drainage, soil characteristics indicative of restricted soil drainage are still present. Surface textures are predominantly loamy sand and sandy loam, but can range from sand to fine sandy loam.

Carrot River soils are usually stone free. Some stones may occur where the fluvial materials are shallow (less than 1 m thick) and underlain by glacial till or gravel. These soils usually occur on undulating or hummocky landscapes with very gentle to gentle slopes.

Carrot River soils frequently occur in complex with soils of other associations. In most of these complexes, the Carrot River soils occur on mid- and lower slopes.

Kinds of Carrot River Soils

Gleyed Orthic Carrot River - The gleyed orthic Carrot River soil usually occurs on mid- and lower slopes. It is a moderately well- to imperfectly drained soil characterized by a dark-gray A horizon, 10 to 19 cm thick, underlain by a brownish-colored B horizon and a grayish to yellowish-brown, moderately calcareous C horizon. There may be a thin, grayish-colored layer with slightly platy structure between the A and B horizons which, upon cultivation, is often incorporated into the plow layer. The B and C horizons often have drab colors and reddish spots and stains, indicative of restricted soil drainage. A peaty layer, less than 40 cm thick, may occur on the surface in some areas.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Carrot River Soils

The best Carrot River soils are those with a fine sandy loam to sandy loam surface texture. These are fair agricultural soils of capability class 3; a moderate moisture deficit, imparted by a low water-holding capacity and the subhumid regional climate, is their main limitation. Carrot River soils with loamy sand and sand surface textures have been rated as capability class 4. These are poor agricultural soils due primarily to their very low water-holding capacity. Carrot River soils may be further downrated based on other soil and landscape limitations (i.e. excessive wetness, erosion, topography, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

As a result of their low to moderate organic matter content and sandy nature, most Carrot River soils have a low natural fertility and are not easily kept in good tilth. They are generally low in potassium, sulphur, nitrogen, and phosphorus. Carrot River soils have a high susceptibility to water erosion and a low susceptibility to wind erosion. Their sandy texture, weak structure and low organic matter content

make it difficult to maintain a cloddy surface that is resistant to wind erosion. It is recommended that soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, and frequent use of legumes and grasses in rotations, be utilized wherever possible to control erosion and maintain or enhance the organic matter content, fertility and aggregate stability of these soils.

Stones are not generally a problem on Carrot River soils, however, periodic clearing may be required where the fluvial materials are shallow (less than 1 m thick) and underlain by glacial till or gravel. Carrot River soils have formed under conditions of restricted drainage and, in most cases, it is necessary to provide artificial drainage before they can be cultivated. In most instances, drainage must be continued on a permanent basis to ensure sustained cultivation.

GLENBUSH (Gb) SOILS

Glenbush soils are Dark Gray soils that have formed in gravelly fluvial materials, in areas of mixed grassland and forest, where wooded vegetation has had some influence on soil formation. Surface textures usually range from loamy sand to sandy loam or loam. Gravelly phases of these textures also occur in some areas.

These soils are usually slightly to moderately stony, however, stoniness can be more severe, especially in areas where the gravelly parent materials are shallow (less than 1 m thick) and overlie glacial till. Glenbush soils are often associated with hummocky landscapes having gentle to moderate slopes. Undulating landscapes with very gentle to gentle slopes and hummocky landscapes with strong or steep slopes are also common in some areas.

Glenbush soils frequently occur in complex with soils of other associations. In many areas of these complexes, the Glenbush soils occur randomly throughout the landscape, whereas, in other areas, they occur in a more regular pattern, often occupying the upper slopes and knolls.

Kinds of Glenbush Soils

Orthic Glenbush - The orthic Glenbush soil can occur on all slope positions. It is a well- to rapidly drained soil with a dark- gray A horizon, 10 to 17 cm thick, underlain by a brownish-colored B horizon of variable thickness and a gravelly, grayish-colored, weakly calcareous C horizon.

Peaty Poorly Drained Soils - Peaty poorly drained soils occur mainly in sloughs and low-lying depressional areas. They are characterized by a shallow layer of peaty material, from 15 to 40 cm thick, that overlies a dark-colored A horizon. The underlying horizons have drab colors that often include reddish spots and stains, indicative of formation under poorly drained conditions. When the organic material is primarily fibric peat, the organic layer may be up to 60 cm thick. Unless artificially drained, most of these soils are frequently wet for all or a significant portion of the growing season and are often flooded.

Agricultural Properties of Glenbush Soils

The Glenbush soils are fair agricultural soils of capability class 3. Their loam to sandy loam textures, coupled with a gravelly subsoil, result in a low water-holding capacity. The coarser-textured Glenbush soils, those with gravelly sandy loam or loamy sand surface textures, are considered to be poor agricultural soils of capability class 4. These soils may be downrated further based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to an individual delineation. For example, Glenbush soils occurring in eroded channels may be so stony that extensive clearing may be required before they can be cultivated. These soils are usually rated as capability class 5. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Glenbush soils generally have moderate amounts of organic matter in the A horizon but, nevertheless, tend to be relatively infertile. In addition to applications of nitrogen and phosphorus, additions of potassium and sulphur may benefit some crops. Due to the sandy nature of these soils, they are often susceptible to wind erosion; thus, stubble conservation and the maintenance of a rough surface may help to slow soil degradation due to wind erosion. Management practices that include forage crops in rotations will help maintain or increase organic matter content and also help prevent wind erosion. Although stones are usually present, amounts vary considerably; some areas are almost stone free, while others require periodic removal if the soils are to be cultivated.

GRONLID (G₀) SOILS

Gronlid soils are Gleyed Dark Gray soils that have formed in loamy lacustrine materials, in areas of mixed grassland and forest, where wooded vegetation and restricted soil drainage have had an influence on soil formation. Soils formed under wooded vegetation are usually slightly leached. They have lower organic matter levels than soils occurring in the Black soil zone, resulting in a dark-gray surface horizon. Although cultivation may have improved surface and internal drainage, soil characteristics indicative of restricted soil drainage are still present. Surface textures range from loam to sandy loam.

Gronlid soils are usually stone free, however, some stones may occur where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel. These soils usually occur on undulating landscapes with very gentle to gentle slopes, however, they can occur on hummocky landscapes with gentle and moderate to strong slopes.

Gronlid soils frequently occur in complex with soils of other associations. In most of these complexes, the Gronlid soils occur on mid- and lower slopes.

Kinds of Gronlid Soils

Gleyed Orthic Gronlid - The gleyed orthic Gronlid soil usually occurs on mid- and upper slopes, however, it can extend onto lower slopes in some landscapes. It is a moderately well- to imperfectly drained soil characterized by a dark-gray A horizon, 16 to 26 cm thick, underlain by a brownish or reddish-brown B horizon and a grayish to yellowish-brown, moderately calcareous C horizon. There may be a thin, grayish-colored layer with platy structure between the A and B horizons which, upon cultivation, is often incorporated into the plow layer. The B and C horizons often have drab colors and reddish spots and stains, indicative of restricted soil drainage. A peaty layer, less than 40 cm thick, may occur on the surface in some areas.

Agricultural Properties of Gronlid Soils

The best Gronlid soils are those with a loam to very fine sandy loam surface texture. These are good agricultural soils of capability class 2; a moderate moisture deficit, imparted by a moderate water-holding capacity and the subhumid regional climate, is their main limitation. Gronlid soils with fine sandy loam to sandy loam surface textures are fair agricultural soils of capability class 3, due primarily to their lower water-holding capacity. Gronlid soils may be further downrated based on other soil and landscape limitations (i.e. excess moisture, erosion, topography, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Because of their moderate organic matter content and sandy nature, most Gronlid soils have moderate natural fertility and tilth. Gronlid soils have a weak to moderate cloddy structure that breaks to fine granular and single grain, making them moderately susceptible to wind erosion. Relatively high infiltration rates, coupled with very gentle to gentle slopes, often result in a low susceptibility to water erosion. Gronlid soils occurring on landscapes with moderate to strong slopes, however, are more susceptible to water erosion. It is recommended that soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, inclusions of forages in crop rotations and shelterbelts, be utilized wherever possible to control soil erosion and maintain or enhance soil organic matter content and aggregate stability.

Stones are not generally a problem on Gronlid soils, however, periodic clearing may be required where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel. Gronlid soils have formed under conditions of restricted drainage and, in most cases, it is necessary to provide artificial drainage before they can be cultivated. In most instances, drainage must be continued on a permanent basis to ensure sustained cultivation.

HOEY (Hh) SOILS

Hoey soils are Thick Black soils that have formed in moderately calcareous, silty lacustrine sediments. They have surface horizons that are usually greater than 20 cm thick. Surface textures range from loam and silt loam to clay loam.

Hoey soils are usually stone free, however, some stones may occur where the lacustrine deposits are shallow (less than 1 m thick) and underlain by gravel or eroded glacial till. Hoey soils are usually associated with undulating landscapes having very gentle to gentle slopes and hummocky landscapes with gentle to moderate slopes.

Hoey soils often occur in complex with soils of other associations. In most of these complexes, the Hoey soils tend to occur on the mid- and lower slope positions.

Kinds of Hoey Soils

Orthic Hoey - The orthic Hoey soil usually occupies lower slope positions, however, it can extend throughout the landscape in some areas. It is a well-drained soil with a black A horizon, 20 to 35 cm thick, underlain by a lime-free B horizon and a grayish-colored, moderately calcareous C horizon.

Agricultural Properties of Hoey Soils

Hoey soils are usually very good agricultural soils of capability class 1. Many of these soils, however, have been downrated based on soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. Some of the Hoey soils have been downrated to agricultural capability class 2 due to the accumulative effect of two or more adverse characteristics of the soil and landscape which, singly, are not serious enough to affect the class rating. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

The thick surface horizon, coupled with high organic matter content, give rise to soils that are reasonably easy to keep in good tilth. These soils are reasonably fertile; however, crops grown in these soils will respond to additions of nitrogen and phosphorus fertilizers in most years. Stones are seldom a problem, although some clearing may be required where these soils are shallow and underlain by gravel or eroded till. Hoey soils are moderately susceptible to wind erosion. Relatively low infiltration rates, when coupled with long slopes, can result in high susceptibility to water erosion. It is recommended that soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, grassing water runways, cultivation across slopes and the establishment of forages, be utilized in order to control soil erosion where necessary.

HAMLIN (Hm) SOILS

Hamlin soils are Black soils that have formed in loamy lacustrine materials. Surface textures range from sandy loam to loam.

Hamlin soils are usually stone free, however, stones may occur where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel, or where the lacustrine materials occur in complex with glacial till or gravel. Hamlin soils usually occur on undulating landscapes with very gentle to gentle slopes, however, they can occur on hummocky landscapes with moderate to strong slopes, particularly where they occur in complex with soils formed in glacial till.

The Hamlin soils commonly occur in complex with soils of other associations. In most of these complexes, the Hamlin soils occur on mid- and lower slopes, however, where they occur in complex with Dark Gray or thick Black soils formed in loamy lacustrine materials, they often occur on upper slope positions.

Kinds of Hamlin Soils

Orthic Hamlin - The orthic Hamlin soil usually occurs on mid- and lower slopes, however, it can extend onto upper slopes in landscapes where slopes are gentle. It is a well-drained soil characterized by a black A horizon, 12 to 18 cm thick, underlain by a brownish-colored B horizon and grayish-brown, moderately calcareous C horizon.

Gleyed Orthic Hamlin - The gleyed orthic Hamlin soil usually occurs on mid- and lower slopes. It is a moderately well-drained soil with a black A horizon, 10 to 20 cm thick, underlain by a brownish-colored B horizon and a drab-colored, moderately calcareous C horizon. While the gleyed orthic Hamlin soil is usually dry by the middle of the growing season, the soil profile contains mottles caused by seasonally high water tables. The mottles appear as rusty- and grayish-colored areas in the soil profile.

Calcareous Hamlin - The calcareous Hamlin soil usually occurs on locally dry upper slopes and knolls, but may extend onto mid- and lower slopes in some landscapes. It is a well-drained soil characterized by a black, usually calcareous A horizon which is usually underlain by a thin, brownish-colored, calcareous B horizon and grayish-brown, moderately calcareous C horizon.

Carbonated Hamlin - The carbonated Hamlin soil usually occurs on mid- and lower slopes, but it may extend onto upper slopes in some landscapes. It is affected to varying degrees by imperfect soil drainage. It is characterized by a highly calcareous A horizon underlain by a highly calcareous B or C horizon. The B and C horizons often have drab colors and reddish spots and stains, indicative of imperfect soil drainage. In addition, the subsoil is often affected by salinity to some degree.

Saline Hamlin - The saline Hamlin soil occurs on lower slopes, often surrounding sloughs or poorly drained depressional areas, and along drainage channels and gullies. It is characterized by the presence of soluble salts, usually within 50 cm of the surface. The salts occur as a white surface crust or as small, white specks within the soil, although salts may not always be visible. Dull colors and reddish spots and stains, indicative of imperfect soil drainage, are often present in the subsoil. It frequently occurs intermixed with carbonated Hamlin soils.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, there may be a thin peaty layer 15 to 40 cm on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Hamlin Soils

The best Hamlin soils are those with a loam or very fine sandy loam surface texture. These are good agricultural soils of capability class 2; a moderate moisture deficit, imparted by the subhumid regional climate and a moderate water-holding capacity, is their main limitation. Hamlin soils with fine sandy loam to sandy loam surface textures are fair agricultural soils of capability class 3 due primarily to their lower water-holding capacity. Although many of the Hamlin soils have few agricultural limitations, some have been downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Most Hamlin soils have a moderate amount of organic matter in the A horizon, resulting in soils of reasonably good tilth. They are low in available phosphorus and high in available potassium. Hamlin soils have a weak to moderate cloddy surface that breaks to fine granular and single grain, making them moderately susceptible to wind erosion. Relatively high infiltration rates, coupled with very gentle to gentle slopes, often result in a low susceptibility to water erosion. Hamlin soils occurring on landscapes with moderate to strong slopes, however, are more susceptible to water erosion. The eroded Hamlin soils on knolls are locally dry because of the rapid runoff associated with these slopes. They have low nutrient reserves and are susceptible to wind and water erosion. It is recommended that soil conservation

practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, strip cropping, inclusion of forages in crop rotations, and shelterbelts, be utilized wherever possible to control soil erosion and maintain or enhance soil organic matter content and aggregate stability. Stones are generally not a problem, however, periodic clearing may be required where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel.

HILLWASH (Hw) SOILS

Hillwash soils are formed in various deposits associated with the steep and eroding sides of escarpments and valleys of rivers, creeks and tributaries. They are a group of shallow, eroded and weakly developed soils. Surface textures and stoniness are extremely variable because of the variable nature of the parent material and the association of these soils with steeply sloping, eroded landscapes.

Agricultural Properties of Hillwash Soils

Hillwash soils are primarily nonarable due to the nature of the landscape on which they occur. Steepness of slope and susceptibility to erosion are the main limitations. They do, however, have some value as pasture land, depending upon steepness of slopes, density of tree cover and availability of water. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

KAMSACK (Ka) SOILS

Kamsack soils are Dark Gray soils that have formed in silty lacustrine materials, where wooded vegetation has had an influence on soil formation. Surface textures are usually silt loam, but loam and clay loam textures can occur.

Kamsack soils are usually stone free, however, a few stones can occur in areas where these soils are shallow (less than 1 m thick) and underlain by a mixture of fluvial gravels and glacial till or eroded glacial till. These soils commonly occur on undulating landscapes with very gentle to gentle slopes and on hummocky landscapes with slopes ranging from gentle to strong.

Kamsack soils frequently occur in complex with soils of other associations. In most of these complexes, the Kamsack soils tend to occur on mid- and lower slope positions.

Kinds of Kamsack Soils

Orthic Kamsack - The orthic Kamsack soil usually occurs on mid- to upper slopes, but can extend onto lower slopes in some landscapes. It is a well-drained soil with a dark grayish-brown A horizon, 11 to 20 cm thick, underlain by a brownish-colored B horizon and a grayish-colored, moderately to strongly calcareous C horizon.

Agricultural Properties of Kamsack Soils

Kamsack soils are generally very good to good agricultural soils of capability classes 1 and 2. A slight moisture deficit, imparted by the subhumid regional climate and a moderate water-holding capacity, is their main limitation. Kamsack soils are often downrated based on other soil and landscape limitations (i.e. salinity, wetness, stones, etc.) that are peculiar to individual delineations. Steepness of slope, for example, may reduce the agricultural capability of these soils to classes 3 or 4 in some areas. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Kamsack soils have a moderate amount of organic matter in the A horizon. Stones are generally not present in these soils, but where the lacustrine materials are shallow and underlain by gravel or eroded till, a few stones may occur and occasional clearing may be required. Wet conditions may be encountered on the imperfectly drained Kamsack soils in the spring, delaying cultivation and seeding operations.

These soils have a low susceptibility to wind erosion. Relatively low infiltration rates, when coupled with long slopes, however, can result in a high susceptibility to water erosion. It is recommended that soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, grassing runways, cultivation across slopes and establishment of forages, be utilized to control soil erosion.

La CORNE (Lc) SOILS

La Corne soils are Gray soils that have formed in loamy lacustrine materials in areas where wooded vegetation has had a strong influence on soil formation. Soils that have formed under these conditions are usually strongly leached, resulting in low organic matter levels and, hence, have a dark-gray to light grayish-colored surface upon cultivation. Surface textures are predominantly fine sandy loam to very fine sandy loam.

La Corne soils are usually stone free, however, some stones may occur where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel. La Corne soils usually occur on undulating landscapes with very gentle to gentle slopes, but can also occur on hummocky landscapes with gentle to moderate slopes, particularly where they occur in complex with soils formed in glacial till.

La Corne soils frequently occur in complex with soils of other associations. In complexes with soils formed in finer-textured lacustrine materials, the La Corne soils tend to occur on mid- and upper slope positions; whereas, in complex with soils formed in coarser-textured sandy fluvial materials or glacial till, they usually occur on the mid- and lower slope positions of the landscape.

Kinds of La Corne Soils

Gray Wooded La Corne - The gray wooded La Corne soil usually occurs on mid- and upper slopes and knolls but in some landscapes, it may be restricted to lower slopes and depressions. It is a well-drained soil which, under forested conditions, is characterized by the presence of a very thin, dark-colored surface horizon below the forest litter, underlain by a strongly leached, gray to grayish-brown, platy horizon of variable thickness. Upon cultivation, part or all of this leached horizon is incorporated into the plow layer, producing a light-gray surface. In some areas, a portion of the strongly leached, grayish-colored horizon will remain below the cultivated surface layer. Below these horizons is a relatively thick, dark-brown to dark grayish-brown, B horizon that usually has a strong (hard), angular blocky to prismatic structure, and a dark grayish-brown, moderately calcareous C horizon.

Dark Gray Wooded La Corne - The dark gray wooded La Corne soil can occur on all slope positions in most La Corne landscapes. It is a well-drained soil which, under forested conditions, is characterized by the presence of a dark-colored A horizon below the forest litter, underlain by a strongly leached, gray to grayish-brown horizon with platy structure. These horizon are underlain by a brownish or reddish-brown B horizon having a strong, angular blocky structure and a dark grayish-brown, moderately calcareous C horizon. Upon cultivation, the forest litter, A horizon and part of the underlying leached horizon are incorporated together, producing a dark-gray surface horizon.

Agricultural Properties of La Corne Soils

La Corne soils are fair agricultural soils of capability class 3 due mainly to their moderate water-holding capacity. Although the majority of the La Corne soils are considered to be class 3 soils, some of them may have been downrated further based on various soil and landscape limitations (i.e. topography, wetness etc.) that are peculiar to individual delineations. La Corne soils occurring on strongly sloping hummocky landscapes, for example, are poor agricultural soils of capability class 4, due to topographic limitations. Ratings for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

The sandy nature and low organic matter content of the La Corne soils result in soils of only moderate fertility and tilth. The low organic matter content in some of these soils results in a structure that makes seedbed preparation difficult and also makes the soils susceptible to crusting after heavy rains, resulting in poor seedling emergence, especially for small-seeded crops. To ensure productivity, management practices should be followed that will increase the organic

matter content, maintain or increase fertility levels and prevent water and wind erosion. Such practices should include using forages in crop rotations, addition of manure, tillage methods that will conserve crop residues, strip cropping and shelterbelts.

MOSS PEAT (MP) SOILS

Moss peat soils are organic soils that have formed in accumulations of vegetative materials, primarily mosses and forest vegetation. These partly decomposed plant residues have accumulated in wet lowland areas. These areas are usually treed with white spruce, black spruce, or tamarack. The ground cover is mainly composed of various types of mosses but may include other herbaceous plants and shrubs. The deposits are usually less than 2 m thick, but greater than 0.6 m thick, with the thickest portion occurring towards the center of the area. The surface is generally level with strong, hummocky microrelief.

The Moss Peat soils may occur in complex with other organic and mineral soil associations. In complexes with Sedge Peat soils, Moss Peat soils occur on the higher landscape positions. In complexes with mineral soils, Moss Peat soils typically occur on the lower landscape positions.

Kinds of Moss Peat Soils

Mesic Moss Peat - The mesic Moss Peat soil is characterized by an accumulation of moderately decomposed plant residues that are derived mainly from the accumulation of mosses. Decomposition of this material has occurred to a moderate extent, such that plant residues are difficult to identify as to their origin.

Agricultural Properties of Moss Peat Soils

Moss Peat soils are not classified for agricultural capability unless they are being used for agricultural purposes. The rating for each delineation is listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report. Areas that are not classified are given the symbol O.

Moss Peat soils are usually very wet and in their natural state, are suitable only for native grazing, or forestry purposes if treed. If these areas are to be developed for agricultural use, they will require drainage and, usually, clearing before breaking. At best, most of these efforts only result in poor agricultural soils. Due to their location in depressional areas, these soils are susceptible to cold air drainage and, consequently, to late spring and early fall frosts. The organic mat is an excellent insulator and, thus, these soils are also slow to warm up in the spring. This may delay crop development and shorten the growing season to the extent that crops may not have sufficient time to mature. Also, once the vegetative mat is disturbed by cultivation and the surface dries, these soils become very susceptible to wind erosion. If drainage is provided, the growing of perennial forage crops

may be more feasible than annual cropping. However, considering the cost of drainage and clearing of these lands, the possible requirement of specialized agricultural equipment, and the potential need for soil amendments such as fertilizer and lime, it may not be economically feasible to grow annual crops on these soils.

MEOTA (Me) SOILS

Meota soils are Black soils that have formed in sandy fluvial materials. Surface textures range from very fine sandy loam to loamy sand.

Meota soils are usually stone free, however, some stones may occur where the sandy deposits are shallow (less than 1 m thick) and underlain by glacial till or gravel. Meota soils frequently occur on undulating landscapes with very gentle to gentle slopes, but in some areas, these soils commonly occur on hummocky landscapes with gentle to moderate slopes.

Meota soils frequently occur in complex with soils of other associations. In most of these complexes, the Meota soils tend to occupy the mid- and lower slope positions. However, in complex with thick sandy soils or some of the Dark Gray soils, the Meota soils often occur on upper slope positions. In other complexes, the Meota soils may occur randomly throughout the landscape.

Kinds of Meota Soils

Orthic Meota - The orthic Meota soil usually occurs on mid- and upper slopes, but may extend onto lower slopes in some Meota landscapes. It is a well- to rapidly drained soil characterized by a black A horizon, 11 to 17 cm thick, underlain by a brownish-colored B horizon and a light-brown, weakly calcareous C horizon.

Gleyed Orthic Meota - The gleyed orthic Meota soil usually occurs on mid- and lower slope positions in the landscape. It is a moderately well- to imperfectly drained soil with a black A horizon overlying B and C horizons, which have dull colors and reddish spots and stains indicative of formation under conditions of restricted drainage.

Eroded Meota - The eroded Meota soil, as the name implies, is a Meota soil whose topsoil has been partially or totally removed by wind erosion. It occurs most often on upper slopes and knolls, but can occur almost anywhere in the landscape. It can be easily recognized in cultivated fields by its light-brown surface color, as compared to the darker-colored Meota soils on mid- and lower slopes.

Carbonated Meota - The carbonated Meota soil occurs on lower slopes, often surrounding sloughs. It is a moderately well- to imperfectly drained soil with a black, usually calcareous A horizon underlain by a highly calcareous B or C horizon. The B and C horizons often have dull colors and reddish spots and stains, indicative of imperfect soil drainage.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and,

occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, there may be a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Meota Soils

The best Meota soils, those with very fine sandy loam surface textures, are good agricultural soils of capability class 2; a moderate moisture deficit, imparted by the sub-humid regional climate and a moderate water-holding capacity, is their main limitation. Most Meota soils, however, have fine sandy loam to sandy loam surface textures, and are, at best, only fair agricultural soils of capability class 3; a low water-holding capacity is their main limitation. Meota soils with coarser textures (loamy sands) have an even lower water-holding capacity and are more droughty. These soils are considered to be poor agricultural soils of capability class 4. Some Meota soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. For example, Meota soils that have been severely eroded may be rated as capability class 4 or 5, depending on the severity of the erosion and its effect on crop production. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

The moderate organic matter content of these soils, coupled with their sandy textures, results in soils of low fertility that are not easily kept in good tilth. The potential for wind erosion is high on most Meota soils due to their sandy nature and weak structure, which makes it difficult to keep a rough surface that is resistant to wind erosion. Management practices that maintain a trash cover and maintain or increase the organic matter content of these soils are recommended. Such practices include reduction of tillage, leaving stubble standing, frequent inclusion of forages in crop rotations and addition of manure. Water erosion is generally not a problem because of their high infiltration rates. Likewise, stones are seldom a problem on these soils.

MARSH (Mh) SOILS

Marsh soils are poorly drained soils that have formed in variable-textured alluvial sediments and are typically associated with low-lying depressional basins and the margins of shallow lakes. In many cases, these soils have formed in

material derived from a variety of sources, thus, they can vary markedly in color, texture and composition. These soils are frequently carbonated to the surface and are often saline. Surface textures are primarily silty clay loam but may be more variable along the edges of some lakes, ranging from sandy loam to clay loam.

Marsh soils are usually stone free, however, some stones may occur in areas associated with the edges of shallow lakes. Marsh soils seldom form complexes with soils of other associations. The notable exception is Sedge Peat. In these areas, the Marsh soils occur on upper slope positions where the thickness of the peat is less than 40 cm.

Kinds of Marsh Soils

Poorly Drained Marsh - The poorly drained Marsh soils occur along the margins of shallow lakes and in large depressional areas or drained lake bottoms. Unless artificially drained, these soils are frequently wet for all or a significant portion of the growing season and are often flooded. The majority of these soils have thick, dark-colored A horizons, 10 to 25 cm thick, and drab subsurface colors that often include reddish spots and streaks. In some areas, there may be a shallow, peaty layer on the surface. These soils are frequently carbonated to the surface (carbonated poorly drained) and, in most cases, are also saline (saline poorly drained), with soluble salts generally present within 50 cm of the surface.

Agricultural Properties of Marsh Soils

Marsh soils are usually wet for all or a significant portion of the growing season, limiting their use for arable agriculture. Improved drainage, however, has permitted cultivation in several areas. These areas are, at best, fair to poor agricultural soils having an agricultural capability rating of class 3 or 4. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Excess wetness and salinity are the main factors affecting the agricultural productivity of Marsh soils. Without improved drainage, Marsh soils are usually only suitable for improved pasture (class 5) or native grazing (class 6). The organic matter content of the surface horizon is usually high, resulting in reasonable fertility and good tilth. Stones are rarely a problem, except in areas where glacial till occurs at or near the surface. Due to their association with lower landscape positions, which receive runoff water in the spring or during periods of intense rainfall, they are susceptible to water erosion and to flooding, often making cultivation difficult. Wind erosion and topography seldom pose serious problems. Areas dominated by very poorly drained soils, especially those with peaty surfaces, or saline poorly drained soils have limited potential for arable agriculture and are best suited for forage production or pasture.

MEADOW (Mw) SOILS

Meadow soils are poorly drained soils that have formed in a mixture of variable-textured alluvial sediments typically associated with low-lying depressional basins. Surface textures are variable but usually range from loam to clay.

Meadow soils are usually stone free, although some stones may occur where the alluvial materials are shallow (less than 1 m thick) and underlain by glacial till. Meadow soils usually occur on nearly level to very gently sloping landscapes.

Meadow soils may occur in complex with soils of other associations. If they are in complex with organic soils, they usually occur on upper slope positions or along the margins of the organic deposit. Occasionally, Meadow soils may occur in complex with well-drained soils. In these cases, the Meadow soils occur in lower landscape positions.

Kinds of Meadow Soils

Poorly Drained Meadow - The poorly drained Meadow soils occur in depressional areas that are subject to flooding. They are wet for all or a significant portion of the growing season. They may have a relatively thick, dark-colored A horizon and drab-colored B and C horizons that are dotted with reddish spots and streaks. The weakly developed poorly drained soil has a thinner A horizon directly overlying the C horizon. In some areas, the poorly drained soils are almost entirely carbonated. In these areas, they have a highly calcareous A horizon underlain by a highly calcareous B or C horizon. In other areas, the soils are covered by a shallow layer of peaty material, from 15 to 40 cm thick. Where the organic material is primarily fibric peat, the organic layer may be up to 60 cm thick. In some cases, the poorly drained soils are also saline. In these areas, soluble salts are usually present within 50 cm of the surface. Saline poorly drained soils often occur intermixed with carbonated poorly drained soils.

Agricultural Properties of Meadow Soils

Meadow soils are fair to poor agricultural soils of capability classes 3 to 6. The wide range in agricultural capability is mainly a result of varying degrees of wetness and salinity. Many of these soils are wet for all or a significant portion of the growing season, thus limiting their use for arable agriculture. Improved drainage, however, has permitted cultivation in some areas and may result in fair agricultural soils. While textures vary, they usually range from loam to clay; consequently, water-holding capacity is often adequate. The organic matter content of the surface horizon is usually high, resulting in reasonable fertility and good tilth. Stones are rarely a problem in Meadow soils. Because of the association of these soils with lower portions of the landscape, wind erosion is not a serious problem. On the other hand, they do receive runoff water in the spring or

during periods of intense rainfall and, consequently, are susceptible to water erosion and to flooding. Areas with peaty surfaces and saline poorly drained soils have little potential for arable agriculture and are best suited for forage production or pasture. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

NAICAM (Nc) SOILS

Naicam soils are Thick Black soils that have formed in shallow (less than 1 m thick), moderately to highly calcareous, silty lacustrine materials underlain by glacial till. These soils have a surface horizon greater than 20 cm thick. Surface textures are predominantly loam to clay loam but can range to silt loam and silty clay loam.

Naicam soils are typically slightly stony to stone free, but may be moderately stony in areas where the lacustrine materials are very shallow or where they occur in complex with glacial till or gravel. Naicam soils are usually associated with undulating or undulating dissected landscapes with very gentle to gentle slopes, but also occur on hummocky landscapes with gentle to moderate slopes, particularly where they occur in complex with soils formed in glacial till.

Naicam soils frequently occur in complex with soils of other associations. In most of these complexes, the Naicam soils occur on the mid- and lower slope positions.

Kinds of Naicam Soils

Orthic Naicam - The orthic Naicam soil usually occupies mid- to lower slopes, however, it may extend onto upper slope positions in some landscapes. It is a well-drained soil characterized by a black A horizon, 20 to 32 cm thick, underlain by a distinct, brownish-colored, lime-free B horizon and a grayish-colored, moderately to strongly calcareous C horizon.

Calcareous Naicam - The calcareous Naicam soil usually occurs on mid- and lower slopes but may also occur upper slopes in some landscapes. It is a well-drained soil characterized by a black, usually calcareous A horizon, 22 to 34 cm thick, usually underlain by a calcareous B horizon, and a grayish-colored, moderately to strongly calcareous C horizon.

Weakly Developed Naicam - The weakly developed Naicam soil usually occurs on mid- and lower slopes, but can also occur on upper slope positions in some landscapes. It is a well-drained soil characterized by a black, usually calcareous A horizon, greater than 20 cm thick, underlain by a grayish-colored, moderately to strongly calcareous C horizon.

Carbonated Naicam - The carbonated Naicam soil occurs on mid- to lower slopes, frequently surrounding sloughs or poorly drained depressions, and in imperfectly to poorly drained depressional areas. It is an imperfectly drained soil characterized by a highly calcareous A horizon, 22 to 30 cm thick, underlain by a highly calcareous B or C

horizon. The B and C horizons often have dull colors and reddish spots and stains, indicative of imperfect soil drainage. Streaks of carbonate frequently occur throughout the upper horizons.

Saline Naicam - The saline Naicam soil usually occurs on lower slopes and in imperfectly to poorly drained depressions or along shallow drainage channels, where the upward movement of subsurface water results in the deposition of salts in upper horizons. Consequently, soluble salts are usually present within 50 cm of the surface. The salts occur as a light-colored surface crust or as small, white specks in the soil, although salts may not always be visible. Dull colors and reddish spots and stains, indicative of imperfect soil drainage, are often present in the subsoil. These soils frequently occur in association with carbonated Naicam soils.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Naicam Soils

Naicam soils are very good agricultural soils with an agricultural capability rating of class 1. These soils may be downrated based on soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. They are often downrated to agricultural capability class 2 due to the accumulative effect of two or more adverse characteristics of the soil and landscape which singly, are not serious enough to affect the class rating. Ratings for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Naicam soils are naturally fertile due to their thick surface horizon and high organic matter content. They are low in available phosphorus and high in available potassium. Crops grown on these soils will respond to additions of nitrogen and phosphorus fertilizers. Some soils may be highly alkaline, which may limit the yield of particular crops and the availability of some plant nutrients.

Stones are seldom a problem on most Naicam soils, however, due to the presence of glacial till near the surface, occasional clearing may be required. Similarly, water erosion is not a serious problem, but can become serious during periods of intense rainfall due to the relatively low infiltra-

tion rate of these soils. The potential for water erosion is greatest in areas with long slopes, and management practices such as cultivation across slopes and grassing of runways, should be followed as much as possible. Wind erosion may be a problem in Naicam soils unless conservation practices are followed. Such practices include continuous cropping and the maintenance of crop residues through reduced tillage or leaving stubble standing.

NORTHERN LIGHT (Nr) SOILS

Northern Light soils are gray soils that have formed in shallow (less than 1 m thick), silty lacustrine materials underlain by glacial till, in areas where wooded vegetation has had a strong influence on soil formation. Soils formed under these conditions are highly leached, resulting in lower organic matter levels and, hence, dark-gray to light grayish-colored surface horizons. Surface textures are predominantly loam and silt loam but can range to silty clay loam.

Northern Light soils are typically stone free to slightly stony, however, they can be moderately stony in areas where the lacustrine materials are very shallow or where they occur in complex with glacial till. Northern Light soils are usually associated with undulating to undulating dissected landscapes with very gentle to gentle slopes, but also occur on hummocky landscapes with gentle to strong slopes, particularly where they occur in complex with soils formed in glacial till.

Northern Light soils frequently occur in complex with soils of other associations. In most of these complexes, the Northern Light soils occur on the mid- and lower slope positions.

Kinds of Northern Light Soils

Gray Wooded Northern Light - The gray wooded Northern Light soil usually occupies upper slope positions, but may extend to all slope positions in some landscapes. It is a well-drained soil which, under forested conditions, is characterized by the presence of a thin, dark-colored surface horizon below the forest litter, underlain by a gray to grayish-brown, strongly leached, platy horizon of variable thickness. Upon cultivation, part or all of this leached horizon is incorporated into the plow layer, producing a light-gray surface. Below these horizons is a relatively thick, dark-brown to dark grayish-brown, lime-free B horizon that usually has a strong (hard) angular blocky to prismatic structure due to an enrichment of clay from the leached horizon. The B horizon is underlain, in turn, by a grayish-colored, moderately calcareous C horizon.

Gleyed Gray Wooded Northern Light - The gleyed gray wooded Northern Light soil commonly occurs on mid- to lower slope positions. It is a moderately well- to imperfectly drained soil characterized by a light-gray A horizon, 14 to 18 cm thick, underlain by a light-gray, leached horizon with platy structure. This horizon is underlain, in turn, by a brownish-colored B horizon that has a distinct, angular

blocky structure when dry. The B horizon is underlain, in turn, by a grayish-colored, moderately calcareous C horizon. The B and C horizons often have dull colors and reddish spots and stains, indicative of imperfect soil drainage.

Dark Gray Wooded Northern Light - The dark gray wooded Northern Light soil occurs on mid- to lower slope positions. It is a well- to moderately well-drained soil which, under forested conditions, is characterized by the presence of a dark-colored surface horizon below the forest litter and a gray to grayish-brown, strongly leached, platy horizon of variable thickness. Upon cultivation, part or all of this leached horizon is incorporated into the surface layer, producing a dark-gray surface, 12 to 19 cm thick. Below these horizons is a relatively thick, dark-brown to dark grayish-brown, lime-free B horizon that usually has a strong (hard), angular blocky to prismatic structure due to an enrichment of clay from the leached horizon. The B horizon is underlain, in turn, by a grayish-colored, moderately calcareous C horizon.

Leached Poorly Drained Soils - Leached poorly drained soils occur mainly in sloughs and low-lying depressional areas which collect runoff from heavy rains and snowmelt. They are usually wet for the early part of the growing season and are often subject to flooding throughout the year after significant rainfall; drainage is restricted by a dense, slowly permeable subsoil. These soils have a platy (leached), light-colored layer below the surface underlain by a dense, grayish-brown B horizon that becomes very hard when dry. When cultivated, the upper layers are mixed, resulting in a light gray-colored surface which is subject to crusting. The underlying B and C horizons have dull colors and reddish spots and streaks, indicative of formation under poorly drained conditions.

Agricultural Properties of Northern Light Soils

Northern Light soils are good (dark gray wooded) to fair (gray wooded) agricultural soils of capability classes 2 and 3, respectively. They have a moderate to slight moisture deficit, due to the regional subhumid climate and a moderate water-holding capacity. The main limitations of these soils are structural. The low organic matter content in some of these soils results in a structure that makes seedbed preparation difficult and also makes these soils susceptible to crusting especially for small-seeded crops, resulting in poor seedling emergence. They have a dense B horizon that may restrict water infiltration and root penetration. These limitations are most strongly expressed in the gray wooded soils. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. Ratings for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Northern Light soils are low in available phosphorus and high in available potassium. Crops usually respond well to applications of nitrogen and phosphorus fertilizers. Sulphur fertilizer may be beneficial to some crops, particularly oil

seeds. These soils are moderately acid to neutral in reaction. The acidity does not affect the productivity of these soils although, there may be local areas with moderate acidity that may inhibit the growth of some sensitive crops such as alfalfa.

Stones are generally not a problem on Northern Light soils, however, due to the presence of glacial till near the surface, occasional clearing may be required. Water erosion is not a serious problem, but can become serious during periods of intense rainfall due to the relatively low infiltration rate of these soils. The potential for water erosion is greatest in areas with long slopes and management practices, such as cultivation across slopes and grassing of runways, should be followed as much as possible. Wind erosion can be a problem unless conservation practices are followed. Such practices include maintenance of crop residues through reduced tillage or leaving stubble standing, continuous cropping or the establishment of forages in seriously affected areas. Management practices, such as the use of legumes in crop rotations, would help to increase the soil organic matter content and, thus, improve the surface structure of these soils.

NISBET (N) SOILS

Nisbet soils are Dark Gray soils that have formed in sandy fluvial materials, in areas where wooded vegetation has had an influence on soil formation. Surface textures range from loamy sand to sandy loam.

Nisbet soils are usually stone free, however, some stones may occur where the sandy deposits are shallow (less than 1 m thick) and underlain by glacial till or gravel. These soils are typically associated with hummocky landscapes having gentle and moderate slopes. Undulating landscapes with very gentle to gentle slopes occur less frequently.

The Nisbet soils often occur in complex with soils of other associations. In most of these complexes, the Nisbet soils occur on mid- and upper slope positions. However, in complexes with soils developed on glacial till, the Nisbet soils usually occur on the mid- and lower slope positions.

Kinds of Nisbet Soils

Orthic Nisbet - The orthic Nisbet soil usually occupies mid- and lower slope positions in the landscape, however, in can extend onto lower slopes in some landscapes. It is a well-drained soil that is characterized by a dark-gray A horizon, 11 to 22 cm thick, underlain by a brown, lime-free B horizon and a light-brown, weakly calcareous C horizon.

Gleyed Orthic Nisbet - The gleyed orthic Nisbet soil usually occurs on lower slopes and in depressional areas. It is a moderately well- to imperfectly drained soil characterized by a dark-gray A horizon, underlain by B and C horizons that often have drab colors and reddish spots and stains, indicative of formation under conditions of imperfect soil drainage.

Calcareous Nisbet - The calcareous Nisbet soil usually occurs on locally dry upper slopes and knolls where runoff

reduces the amount of water entering the soil. This results in a thinner soil with less organic matter than the Nisbet soils on mid- and lower slopes. It has a thin, usually calcareous A horizon, and a thin, calcareous B horizon. This soil has usually been affected by erosion to some degree, but the extent of erosion is less than on the eroded Nisbet soil.

Eroded Nisbet - The eroded Nisbet soil, as the name implies, is a Nisbet soil whose topsoil has been partially or almost totally removed by erosion. The A horizon is usually less than 10 cm thick. It occurs on upper slopes and knolls and can be easily recognized in cultivated fields by its light brownish- to grayish-colored surface. It occurs most often in hummocky landscapes with moderate to steep slopes.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas which collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Nisbet Soils

Nisbet soils with fine sandy loam and sandy loam surface textures are fair agricultural soils of capability class 3 due primarily to their sandy nature. The occurrence of these soils in an area of more favorable precipitation compensates slightly for the low water-holding capacity. The coarser-textured Nisbet soils, those with loamy sand and sand surface textures, are considered to be poor agricultural soils of capability class 4. Nisbet soils may be further downrated based on soil and landscape limitations (i.e. salinity, wetness, topography, stones, erosion, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

The Nisbet soils have a low organic matter content which, combined with the sandy nature of this soil, results in one of the poorest agricultural soils in the Dark Gray soil zone. Fertilizer applications, particularly nitrogen and phosphorus, can increase their productivity somewhat. In addition, applications of potassium and sulphur may be required for certain crops. The potential for erosion is high on these soils because of their sandy texture, weak structure and low organic matter content, making it difficult to keep a cloddy surface that is resistant to wind erosion. It is recommended that soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, frequent inclusion of forages in crop rotations, and shelter-

belts, be utilized wherever possible to control soil erosion and maintain or enhance soil organic matter content and aggregate stability. Stones are generally not a problem. Areas of poorly drained soils, especially those with peaty surfaces, often have little potential for arable agriculture and are best suited for forage production or pasture.

OXBOW (Ox) SOILS

Oxbow soils are Black soils that have formed in loamy glacial till. Surface textures are predominantly loam but can range from sandy loam to clay loam.

Oxbow soils are usually slightly to moderately stony, but some areas are very stony. Oxbow soils can occur on a variety of landscapes, but most commonly occur on undulating landscapes with very gentle to moderate slopes and on hummocky landscapes with slopes ranging from gentle to moderate in some areas to steep in others.

Oxbow soils frequently occur in complex with soils of other associations. In most of these complexes, the Oxbow soils occur on the mid- and upper slopes, however, in complex with Dark Brown or Dark Gray soils, they may occur on lower slopes.

Kinds of Oxbow Soils

Orthic Oxbow - The orthic Oxbow soil occurs on mid-slopes in most Oxbow landscapes, however, it may extend onto upper and lower slope positions in landscapes with gentle slopes. It is a well-drained soil characterized by a black A horizon, 11 to 18 cm thick, underlain by a brownish-colored B horizon and a grayish-colored, moderately calcareous C horizon.

Gleyed Orthic Oxbow - The gleyed orthic Oxbow soil commonly occurs on mid- and lower slope positions in most Oxbow landscapes. It is a moderately well- to imperfectly drained soil characterized by a black A horizon, 11 to 19 cm thick, underlain by B and C horizons that often have dull colors and reddish spots and stains, indicative of restricted drainage.

Calcareous Oxbow - The calcareous Oxbow soil occurs on locally dry upper slopes and knolls where runoff reduces the amount of water entering the soil. This results in a thinner soil with less organic matter than the surrounding orthic Oxbow soils. It is characterized by a thin, usually calcareous A horizon, 10 to 16 cm thick, that is usually underlain by a thin, calcareous B horizon and a grayish-colored, moderately calcareous C horizon.

Eroded Oxbow - The eroded Oxbow soil, as the name implies, is an Oxbow soil whose topsoil has been partially or almost totally removed by erosion. It occurs on knolls and upper slopes and can be easily recognized in cultivated fields by its light-brown to grayish-colored surface. It occurs most often on hummocky landscapes with moderate to steep slopes and, in severely eroded areas, may occupy 30% or more of the landscape.

Eluviated Oxbow - The eluviated Oxbow soil occurs on mid- to lower slopes and, occasionally, in depressional areas. It is a well- to moderately well-drained soil with a black A horizon, 14 to 19 cm thick, underlain by a lighter-colored, leached horizon with platy structure. This horizon, in turn, is underlain by a dense, dark-gray to dark grayish-brown B horizon and a moderately calcareous, grayish-colored C horizon.

Carbonated Oxbow - The carbonated Oxbow soil occurs on lower slopes, frequently surrounding sloughs or poorly drained depressions. It is characterized by a highly calcareous A horizon underlain by a highly calcareous B or C horizon. The B and C horizons often have drab colors and reddish spots and stains, indicative of imperfect soil drainage. Streaks of carbonate may occur throughout most or all of the profile.

Saline Oxbow - The saline Oxbow soil occurs on lower slopes, often surrounding sloughs or poorly drained depressional areas, and along drainage channels or gullies. It is characterized by the presence of soluble salts, usually within 50 cm of the surface. The salts occur as a white surface crust or as small, white specks within the soil, although salts may not always be visible. Dull colors and reddish spots and stains, indicative of imperfect soil drainage, are often present in the subsoil. This soil frequently occurs intermixed with carbonated Oxbow soils on lower slopes surrounding sloughs.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Oxbow Soils

The best Oxbow soils are those with a clay loam surface texture. These are very good agricultural soils of capability class 1. The more common Oxbow soils, those with a loam surface texture, are good agricultural soils of capability class 2; a slight moisture deficit, imparted by the subhumid regional climate and a moderate water-holding capacity, is their main limitation. Oxbow soils with a sandy loam surface texture are fair agricultural soils of capability class 3. Oxbow soils are often downrated further based on other soil and landscape limitations (i.e. salinity, wetness, topography, stones, erosion, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Most Oxbow soils have a moderate amount of organic matter in the A horizon resulting in reasonably fertile soils of good tilth. These soils, however, are usually low in available phosphorus but high in available potassium. Crops grown on most Oxbow soils respond to additions of nitrogen and phosphorus fertilizers. Oxbow soils range from slightly to very stony, hence, the number of clearing operations required each year will vary considerably.

Oxbow soils occur on a variety of landscapes. Where they occur on landscapes with very gentle to moderate slopes, they have a low susceptibility to wind and water erosion. However, it is not unusual for these soils to occur on landscapes with strong to steep slopes (sometimes dissected), in which case they will have a high to very high susceptibility to water erosion. The eroded Oxbow soils on steep knolls and ridges have low nutrient reserves and are locally dry because of the rapid runoff associated with these slopes. Where Oxbow soils have undergone severe erosion or where they are very susceptible to erosion, it is recommended that soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, grassing water runways, cultivation across slopes and establishment of forages, be utilized to control soil erosion.

PATHLOW (Pa) SOILS

Pathlow soils are a mixture of Gray and Dark Gray soils that have formed in a complex mixture of glacial till, eroded glacial till, and shallow, silty lacustrine materials underlain by eroded glacial till, in areas where wooded vegetation has influenced soil formation. Soils formed under these conditions are usually leached; the degree of leaching is dependent on surface drainage and slope position. These soils have low to moderate amounts of organic matter, resulting in gray to dark-gray surface colors. Surface textures range from loam to clay loam.

These soils are associated with broad ridged landscapes that are usually very gently to moderately sloping and frequently dissected. Areas of Pathlow soils are dominated by a shallow, silty lacustrine veneer overlying severely eroded glacial till. At the lacustrine-glacial till contact, there is a dense layer of stones and/or gravel indicative of the severe erosion that occurred prior to the deposition of the silty lacustrine material. On some ridge tops, the eroded glacial till extends to the surface. Where this occurs, the till is very stony to exceedingly stony. Consequently, many of the ridge tops have not been cultivated and remain treed.

Kinds of Pathlow Soils

Orthic Pathlow - The orthic Pathlow soil usually occurs on lower slope positions in the landscape. It is a well- to moderately well-drained soil with a dark-gray A horizon, 10 to 15 cm thick. The A horizon is underlain, in turn, by a brownish-colored, clay-enriched B horizon which usually has a strong, angular blocky structure. There may be a thin, grayish-colored, leached layer with platy structure between the A and B horizons which, if cultivated, is often incorpo-

rated into the plow layer. The glacial till usually occurs below the A and B horizons.

Gray Wooded Pathlow - The gray wooded Pathlow soil occurs mainly on upper slopes and, occasionally, on midslopes. It is a well-drained soil which, under forested conditions, is characterized by the presence of a thin, dark-colored surface horizon below the forest litter, underlain by a light-gray, leached horizon with platy structure and a brownish-colored B horizon that has been enriched by clay leached from upper horizons. Where these soils are cultivated, the leaf litter has been mixed with part or all of the leached horizon, resulting in a light-gray A horizon that is 10 to 16 cm thick.

Dark Gray Wooded Pathlow - The dark gray wooded Pathlow soil occurs mainly on mid- to lower slopes. It is a well- to moderately well-drained soil with a dark-gray A horizon, 12 to 16 cm thick, overlying a lighter-colored, leached horizon which, in turn, overlies a brownish-colored B horizon enriched by clay leached from the upper horizons. The transition from the silty lacustrine material to glacial till often occurs within the A or B horizon.

Agricultural Properties of Pathlow Soils

Pathlow soils are, at best, good to fair agricultural soils of capability classes 2 to 3, respectively. They have a slight moisture deficit due to the regional subhumid climate and a moderate water-holding capacity. Pathlow soils also have soil structural limitations due to a dense and compact B horizon, that is enriched by clay leached from the A horizon, which may restrict root penetration. Also, they have low to moderate amounts of organic matter and are very susceptible to crusting. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to an individual delineation. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Management practices that include forages in crop rotations should be used to enhance soil organic matter content and improve soil structure. Stones, particularly on upper slopes, often pose a serious hindrance to cultivation. If these soils are broken, extensive clearing of stones and boulders is usually required prior to cultivation of the land. Sulphur fertilizer may benefit some crops such as oil seeds, particularly on the ridge tops where gray wooded Pathlow soils occur. As well, these soils have silty textures and are associated with long slopes, making them very susceptible to water erosion. Stubble conservation and contour cultivation will help to slow water movement over the soil surface.

PERLEY (Pe) SOILS

Perley soils are Thick Black soils that have formed in sandy fluvial materials. They are distinguished from Meota soils by having a surface horizon that is greater than 20 cm thick. Surface textures range from very fine sandy loam to loamy sand.

Perley soils are usually stone free, however, some stones may occur where the sandy deposits are shallow (less than 1 m thick) and underlain by glacial till or gravel. Perley soils frequently occur on very gently to gently undulating landscapes, but in some areas, these soils commonly occur on hummocky landscapes with gentle to moderate slopes.

Perley soils frequently occur in complex with soils of other associations. In most of these complexes, the Perley soils occur on the mid- and lower slope positions.

Kinds of Perley Soils

Orthic Perley - The orthic Perley soil occupies mid- and upper slope positions in most Perley landscapes. It is a well- to rapidly drained soil with a black A horizon, 20 to 36 cm thick, underlain by a brownish-colored B horizon and a light-brown, weakly to moderately calcareous C horizon.

Gleyed Orthic Perley - The gleyed orthic Perley soil usually occurs on lower slopes and in depressional areas, but it can extend to upper slopes in some Perley landscapes. It is a moderately well- to imperfectly drained soil with a black A horizon, greater than 20 cm thick, underlain by B and C horizons which often have drab colors and reddish spots and stains, indicative of formation under imperfectly drained soil conditions.

Calcareous Perley - The calcareous Perley soil usually occurs on mid- and upper slopes and knolls, but it can extend onto lower slopes in some Perley landscapes. It is a well- to moderately well-drained soil with a black, usually calcareous A horizon, 20 to 37 cm thick, underlain by a brownish-colored, calcareous B horizon and a grayish-colored, weakly to moderately calcareous C horizon.

Carbonated Perley - The carbonated Perley soil occurs in low-lying depressional areas or on lower slope positions, often surrounding sloughs. It is a moderately well- to imperfectly drained soil with a black, usually calcareous A horizon, 20 to 40 cm thick, underlain by a highly calcareous B or C horizon. The B and C horizons often have drab colors and reddish spots and stains, indicative of imperfect soil drainage. Salts are frequently present in the C horizon of carbonated Perley soils.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that often include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Perley Soils

The best Perley soils, particularly the typical or orthic Perley soils with a very fine sandy loam surface texture, are good agricultural soils of capability class 2; a moderate moisture deficit, imparted by the subhumid regional climate and a moderate water-holding capacity, is their main limitation. Most Perley soils, however, have fine sandy loam to sandy loam surface textures, and are, at best, only fair agricultural soils of capability class 3. Their low water-holding capacity is their main limitation. Perley soils with coarser textures (loamy sands) are even more droughty and are considered to be poor agricultural soils of capability class 4. Perley soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to an individual delineation. For example, Perley soils are downrated to class 4 or 5 if the potential for crop production is severely affected by salinity. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

A moderate to high organic matter content makes most Perley soils reasonably fertile, although additions of phosphorus and nitrogen are usually required to obtain optimum yields. Due to their sandy nature and weak structure, these soils are often subject to wind erosion, particularly when the surface is not protected by a trash cover. Tillage practices that maintain a trash cover and management practices that maintain or increase the organic matter content will help reduce this hazard. Water erosion is generally not a problem because of high infiltration rates. Likewise, stones are seldom a problem on these soils.

PORCUPINE PLAIN (Pp) SOILS

Porcupine Plain soils are Gray soils that have formed in silty lacustrine materials, in areas where wooded vegetation has had a strong influence on soil formation. Soils that have formed under these conditions are usually strongly leached, resulting in low organic matter levels and, hence, a dark-gray to light gray-colored surface horizon upon cultivation. Surface textures are predominantly loam and silt loam but can range to silty clay loam.

Porcupine Plain soils are typically stone free. However, where these soils are shallow (less than 1 m thick) and underlain by gravel, a few pebbles may occur at depth or throughout the profile. In areas where these soils are underlain by eroded till, large boulders may be present causing a serious hindrance to cultivation. Porcupine Plain soils are usually associated with undulating or undulating dissected landscapes with very gentle to gentle slopes, but also occur on hummocky landscapes with gentle to strong slopes.

Porcupine Plain soils frequently occur in complex with soils of other associations. In complexes with soils of sandy fluvial or loamy lacustrine origin, they often occur on the mid- to lower slopes. In complexes with soils of finer textured lacustrine origin, they usually occur on the upper slopes.

Kinds of Porcupine Plain Soils

Gray Wooded Porcupine Plain - The gray wooded Porcupine Plain soil usually occupies upper slope positions, however, it may extend to all slope positions in some landscapes. It is a well-drained soil which, under forested conditions, is characterized by the presence of a thin, dark-colored surface horizon below the forest litter, underlain by a gray to grayish-brown, strongly leached, platy horizon of variable thickness. Upon cultivation, part or all of this leached horizon is incorporated into the plow layer, producing a light-gray surface. Below these horizons is a relatively thick, dark-brown to dark grayish-brown, lime-free B horizon that usually has a strong (hard), angular blocky to prismatic structure due to an enrichment of clay from the leached horizon. The B horizon, in turn, is underlain by a grayish-colored, moderately calcareous C horizon.

Gleyed Gray Wooded Porcupine Plain - The gleyed gray wooded Porcupine Plain soil commonly occurs on mid- to lower slope positions. It is a moderately well- to imperfectly drained soil characterized by a light-gray A horizon, 8 to 15 cm thick, underlain by a light-gray, leached horizon with platy structure. This horizon, in turn, is underlain by a brownish-colored B horizon that has a distinct, angular blocky structure when dry. The B horizon is underlain by a grayish-colored, moderately calcareous C horizon. The B and C horizons often have dull colors and reddish spots and stains, indicative of imperfect soil drainage.

Peaty Poorly Drained Soils - Peaty poorly drained soils occur mainly in sloughs and low-lying depressional areas. They are characterized by a shallow layer of peaty material, from 15 to 40 cm thick, that overlies a dark-colored A horizon. The underlying horizons have drab colors that often include reddish spots and stains, indicative of formation under poorly drained conditions. When the organic material is primarily fibric peat, the organic layer may be up to 60 cm thick. Unless artificially drained, most of these soils are frequently wet for all or a significant portion of the growing season and are often flooded.

Agricultural Properties of Porcupine Plain Soils

Porcupine Plain soils are good (dark gray wooded) to fair (gray wooded) agricultural soils of capability classes 2 and 3, respectively. They have a slight moisture deficit, due to the regional subhumid climate and a moderate water-holding capacity. The main limitations in these soils are structural. The low organic matter content in some of these soils results in a structure that makes seedbed preparation difficult and also makes the soil susceptible to crusting after heavy rains, resulting in poor seedling emergence, especially for small-seeded crops. They have a dense B horizon that may restrict water infiltration and root penetration. These limitations are most strongly expressed in the gray wooded soils. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations.

Ratings for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Porcupine Plain soils are low in available phosphorus and high in available potassium and crops usually respond well to applications of nitrogen and phosphorus fertilizers. Sulphur fertilizer may be beneficial to some crops, particularly oil seeds. These soils are moderately acid to neutral in reaction. The acidity does not affect the productivity of these soils, although there may be local areas with moderate acidity that may inhibit the growth of some sensitive crops such as alfalfa.

Stones are generally not a problem in Porcupine Plain soils, however, where the lacustrine materials are shallow and underlain by eroded till, stones often cause a serious hindrance to cultivation and usually require clearing on an annual basis. Water erosion is not a serious problem, but can become serious during periods of intense rainfall due to the relatively low infiltration rate of these soils. The potential for water erosion is greatest in areas with long slopes so management practices, such as cultivation across slopes and grassing of runways, should be followed as much as possible. Wind erosion can be a problem unless conservation practices are followed. Such practices include maintenance of crop residues through reduced tillage or leaving stubble standing and continuous cropping. Management practices, such as the use of legumes in crop rotations, would help to increase the soil organic matter content and, thus, improve the surface structure of these soils.

RUNWAY (Rw) SOILS

Runway soils are formed in various deposits associated with the sides and bottoms of shallow drainage channels. This group of soils range from weakly developed to poorly drained and are primarily associated with dissected landscapes. As a result, surface texture, degree of stoniness, slope class and salinity are extremely variable.

Agricultural Properties of Runway Soils

Runway soils are usually rated as class 4, 5 or 6 for agricultural capability. Most of these soils, however, are nonarable in that the bottom lands are poorly drained and the side slopes are often too steep to permit cultivation. A few areas, where slopes permit crossing with field implements, have some potential for cultivation. As well, many areas have little potential for grazing land as they occur as narrow strips cutting through cultivated areas. Where they are large enough to be fenced, they do have some value as pasture land depending upon steepness of slope, density of tree cover and availability of water. Ratings for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

SEDGE PEAT (SP) SOILS

Sedge Peat soils are Organic soils that have formed in accumulations of vegetative materials, primarily sedges, meadow grasses and shrubs. These partly decomposed plant residues have accumulated in wet lowland areas. The native vegetation in these areas consist mainly of sedges, cattails, rushes, reeds and grasses, along with scattered willows and other shrubs. Trees such as tamarack and black spruce may also be present. The peat is derived from the accumulation of this vegetative material and is often saturated at or near the surface. In these areas, it is not unusual for the center of the deposit to have a small water body. The deposits are usually less than 2 m thick, but greater than 0.6 m thick, with the thickest part occurring towards the center of the deposit. The surface is generally level with strong, hummocky micro-relief.

Sedge Peat soils may occur in complex with soils of other associations. In these complexes, the Sedge Peat soils typically occur on lower landscape positions.

Kinds of Sedge Peat Soils

Fibric Sedge Peat - The fibric Sedge Peat soil is characterized by an accumulation of relatively undecomposed plant residues that are derived mainly from the accumulation of sedges, meadow grasses and shrubs. Decomposition of this material has been limited and residues are readily identifiable as to their origin.

Mesic Sedge Peat - The mesic Sedge Peat soil is characterized by an accumulation of moderately decomposed plant residues that are derived mainly from the accumulation of sedges, meadow grasses and shrubs. Decomposition of this material has occurred to a moderate extent, such that plant residues are difficult to identify as to their origin.

Humic Sedge Peat - The humic Sedge Peat soil is characterized by an accumulation of strongly decomposed plant residues that are derived mainly from the accumulation of sedges, meadow grasses and shrubs. Decomposition of this material has occurred to such an extent that plant residues are impossible to identify as to their origin.

Agricultural Properties of Sedge Peat Soils

Sedge Peat soils are not classified for agricultural capability unless they are being used for agricultural purposes. The rating for each delineation is listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report. Areas that are not classified are given the symbol O.

Sedge Peat soils, in their natural state, are very wet and are suitable only for native grazing. Areas with open water are not suitable for agricultural use. If these lands are to be

developed for agricultural use, they will require drainage and, usually, some clearing before breaking. Disposal of drainage water may not be feasible in some locations. Due to their low-lying position, these areas are susceptible to cold air drainage and, thus, to late spring and early fall frosts. The organic mat is also an excellent insulator and, thus, these soils are slow to warm up in the spring. This may shorten the growing season to the extent that crops may not have sufficient time to mature. Also, once the vegetative mat is disturbed by cultivation and the surface dries, these soils become very susceptible to wind erosion. If drainage is provided, the growing of perennial forage crops may be more feasible than annual cropping. Considering the cost of drainage and clearing of these areas, the possible requirement for specialized agricultural equipment, and the potential need for soil amendments such as fertilizer and lime, it may not be economically feasible to grow annual crops on these soils.

SHELLBROOK (Sb) SOILS

Shellbrook soils are Dark Gray soils that have formed in loamy lacustrine materials, in areas of mixed grassland and forest, where wooded vegetation has had some influence on soil formation. Soils that have formed under these conditions are usually slightly leached. They have lower organic matter levels than similar soils occurring in the Black soil zone, as evidenced by a dark-gray surface horizon. Surface textures range from sandy loam to loam.

Shellbrook soils are usually stone free. Some stones may occur where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel, or where the lacustrine materials occur in complex with glacial till or gravel. Shellbrook soils usually occur on undulating landscapes with very gentle to gentle slopes, however, they can also occur on hummocky landscapes with gentle to moderate slopes.

Shellbrook soils occur in complex with soils of many other associations. In most of these complexes, the Shellbrook soils occur on mid- and lower slopes, however, in complex with certain soils, such as Black soils formed in loamy lacustrine materials, they tend to occur on upper slope positions.

Kinds of Shellbrook Soils

Orthic Shellbrook - The orthic Shellbrook soil usually occurs on mid- and lower slopes, however, it can extend onto upper slopes in some landscapes. It is a well-drained soil characterized by a dark-gray A horizon, 11 to 22 cm thick, underlain by a reddish-brown B horizon and a grayish-colored, moderately calcareous C horizon. There may be a thin, grayish-colored layer with platy structure between the A and B horizons which, if cultivated, is often incorporated into the plow layer.

Agricultural Properties of Shellbrook Soils

The best Shellbrook soils are those with loam to very fine sandy loam surface textures. These are good agricultural soils of capability class 2; a slight moisture deficit, imparted by the subhumid regional climate and a moderate water-holding capacity, is their main limitation. Shellbrook soils with fine sandy loam to sandy loam surface textures are fair agricultural soils of capability class 3 due primarily to their low water-holding capacity. Shellbrook soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, erosion, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Shellbrook soils have a moderate amount of organic matter in the A horizon. They are low in available phosphorus and high in available potassium. Shellbrook soils have a weak to moderate cloddy structure that breaks to fine granular and single grain, making them moderately susceptible to wind erosion. Relatively high infiltration rates, coupled with very gentle to gentle slopes, often result in a low susceptibility to water erosion. Shellbrook soils occurring on landscapes with moderate slopes, however, are more susceptible to water erosion. It is recommended that soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, inclusions of forages in crop rotations, strip cropping and shelterbelts, be utilized wherever possible to control soil erosion and maintain or enhance soil organic matter content and aggregate stability.

Stones are not generally a problem on Shellbrook soils, however, periodic clearing may be required where the lacustrine materials are shallow (less than 1 m thick) and underlain by glacial till or gravel.

SYLVANIA (Sy) SOILS

Sylvania soils are Gray soils that have formed in sandy fluvial materials, in areas where wooded vegetation has had a strong influence on soil formation. Soils that have formed under these conditions are usually strongly leached, resulting in low organic matter levels and, hence, a grayish-colored surface upon cultivation. Surface textures are predominantly loamy sand to very fine sandy loam.

Sylvania soils are usually stone free, however, some stones may occur where the fluvial materials are shallow (less than 1 m thick) and underlain by glacial till or gravel. Sylvania soils usually occur on hummocky landscapes with gentle to moderate slopes, but can also occur on undulating landscapes with very gentle to gentle slopes.

Sylvania soils frequently occur in complex with soils of other associations. In complexes with soils formed in finer-textured lacustrine materials, the Sylvania soils tend to occur on mid- and upper slope positions, whereas, in complexes

with soils formed in coarser-textured fluvial materials or glacial till, they usually occur on the mid- and lower slope positions of the landscape.

Kinds of Sylvania Soils

Gray Wooded Sylvania - The gray wooded Sylvania soil usually occurs on mid- and upper slopes and knolls but, in some landscapes, it may be restricted to lower slopes and depressions. It is a well- to excessively drained soil which, under forested conditions, is characterized by the presence of a thin, dark-colored surface horizon below the forest litter, underlain by a strongly leached, gray to grayish-brown, platy horizon of variable thickness. Upon cultivation, part or all of this leached horizon is incorporated into the plow layer, producing a light-gray surface horizon, 11 to 16 cm thick. In some areas, a portion of the strongly leached, grayish-colored horizon remains below the cultivated surface layer. Below these horizons is a relatively thick, dark-brown to dark grayish-brown B horizon that usually has a strong (hard), angular blocky to prismatic structure, and a dark grayish-brown, weakly calcareous C horizon with a weak, blocky to massive structure.

Dark Gray Wooded Sylvania - The dark gray wooded Sylvania soil usually occurs on mid- and lower slopes, but it can extend onto upper slopes in some landscapes. It is a well-drained soil which, under forested conditions, is characterized by the presence of a dark-colored A horizon below the forest litter, underlain by a strongly leached, gray to grayish-brown horizon with platy structure. Upon cultivation, the forest litter, A horizon and part of the underlying leached horizon are incorporated together, producing a dark-gray surface horizon, 10 to 18 cm thick.

These horizons are underlain, in turn, by a brownish or reddish-brown B horizon having a strong, angular blocky structure, and a dark grayish-brown, weakly calcareous C horizon with a weak, blocky to massive structure.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Sylvania Soils

The best Sylvania soils are those with very fine sandy loam textures. They are considered to be fair agricultural soils of capability class 3. Areas with fine sandy loam to

sandy loam surface textures are, at best, only poor agricultural soils of capability class 4. A low water-holding capacity is the main agricultural limitation of these soils. Areas with very light surface textures (loamy sand) are rated class 5 (nonarable and suitable only for perennial forages or improved pasture) due to their very low water-holding capacity and low natural fertility. Sylvania soils may be further downrated based on other soil and landscape limitations (i.e. topography, wetness, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

The sandy nature and low organic matter content of the Sylvania soils result in soils of low fertility and poor tilth. They are coarse textured and lack drought resistance, while their loose surface structure makes them susceptible to wind erosion when cultivated. They are not usually susceptible to water erosion due to their high infiltration rate. To ensure productivity, management practices should be followed that will increase the organic matter content, maintain or increase fertility levels and prevent wind erosion. Sylvania soils may be better utilized for the production of forage crops than cereal crops. This would add organic matter to the soil and, along with proper tillage methods, protect against erosion. As well, the use of fertilizers would help ensure their continual use as arable agricultural land.

TIGER HILLS (Tg) SOILS

Tiger Hills soils are Dark Gray soils that have formed in shallow (less than 1 m thick), silty lacustrine materials underlain by glacial till, in areas of mixed grassland and forest, where wooded vegetation has had some influence on soil formation. Surface textures are predominantly loam and silt loam, but can range to silty clay loam and clay loam.

Tiger Hills soils are typically slightly stony to stone free, but may be moderately stony in areas where the lacustrine materials are very shallow or where they occur in complex with glacial till or gravel. They are usually associated with undulating or undulating dissected landscapes with very gentle to gentle slopes, but can also occur on more strongly sloping hummocky landscapes, particularly where they occur in complex with soils formed in glacial till.

Tiger Hills soils frequently occur in complex with soils of other associations. In most of these complexes, the Tiger Hills soils occur on the mid- and lower slope positions.

Kinds of Tiger Hills Soils

Orthic Tiger Hills - The orthic Tiger Hills soil usually occurs on lower slopes, however, it can extend onto mid- and upper slopes in some landscapes. It is a well-drained soil characterized by a dark-gray A horizon, 12 to 22 cm thick, underlain by a dark brownish-colored B horizon and a yellowish-brown, moderately calcareous C horizon.

Gleyed Orthic Tiger Hills - The gleyed orthic Tiger Hills soil usually occurs on mid- to lower slopes. It is a moderately well- to imperfectly drained soil characterized by

a dark-gray A horizon, 13 to 20 cm thick, underlain by a lime-free B horizon and a moderately calcareous C horizon. The B and C horizons often have dull colors and reddish spots and stains, indicative of imperfect soil drainage.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Tiger Hills Soils

Tiger Hills soils are generally good agricultural soils of capability class 2. A slight moisture deficit, imparted by a moderate water-holding capacity and the subhumid regional climate, is their main limitation. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. Ratings for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Tiger Hills soils have a moderate amount of organic matter in the A horizon. They are low in available phosphorus and high in available potassium. Crops grown on these soils usually respond well to applications of nitrogen and phosphorus fertilizers.

Stones are seldom a problem on Tiger Hills soils, however, due to the presence of glacial till near the surface, a few stones may be present and occasional clearing may be required. Similarly, water erosion is not a serious problem, but can become serious during periods of intense rainfall due to the relatively low infiltration rate of these soils. The potential for water erosion is greatest in areas with long slopes and management practices, such as cultivation across slopes and grassing of runways, should be followed as much as possible. Wind erosion may be a problem, unless conservation practices are followed. Such practices include maintenance of crop residues through reduced tillage or leaving stubble standing, continuous cropping, or the establishment of forages in seriously affected areas.

WHITEWOOD (Wh) SOILS

Whitewood soils are Dark Gray soils that have formed in loamy glacial till, in areas of mixed grassland and forest, where wooded vegetation has had an influence on soil formation. Soils formed under these conditions are usually

slightly leached, resulting in lower organic matter levels than similar soils occurring in the Black soil zone, and resulting in a dark gray-colored surface horizon. Surface textures are predominantly loam but may range from sandy loam to clay loam.

These soils are moderately to very stony. They usually occur on hummocky landscapes having gentle to moderate slopes, although steeper slopes are common in some areas.

Where Whitewood soils occur in complex with Black soils such as Oxbow, their occurrence in the landscape is a function of drainage and precipitation. In southern parts of the Black soil zone, Whitewood soils tend to occur on the lower slopes, whereas, in the northern more humid parts of the zone, the Whitewood soils occur on the well-drained upper slopes. Whitewood soils also tend to occupy upper slope positions when they occur in complex with soils of the Gray soil zone. When Whitewood soils occur in complex with soils formed in lacustrine or fluvial materials, they usually occupy mid- to upper slope positions.

Kinds of Whitewood Soils

Orthic Whitewood - The orthic Whitewood soil occupies lower slopes in most Whitewood landscapes, but may extend to the midslope in some areas, and upper slopes and knolls in other areas. It is a well-drained soil with a dark-gray A horizon, 10 to 18 cm thick, underlain by a brownish or reddish-brown B horizon that usually has a moderate, angular blocky structure when dry. There may be a thin, grayish-colored layer with platy structure between the A and B horizons which, if cultivated, is often incorporated into the plow layer. This mixing imparts a dark-gray color to the soil surface.

Gleyed Orthic Whitewood - The gleyed orthic Whitewood soil usually occurs on lower slope positions, although it may extend into depressions in some areas. It is a moderately well- to imperfectly drained soil with a dark-gray A horizon that is usually 10 to 20 cm thick, although it may be thicker in some lower slope positions. The underlying lime-free B horizon and calcareous C horizon have dull colors and reddish spots and stains, indicative of imperfect soil drainage.

Calcareous Whitewood - The calcareous Whitewood soil occurs on locally dry upper slopes and knolls where runoff reduces the amount of water entering the soil. This results in a thinner soil with less organic matter than the surrounding Whitewood soils. It has a thin, usually calcareous A horizon and a thin, calcareous B horizon. This soil is often affected by erosion but the extent of erosion is less than on the eroded Whitewood soil.

Eroded Whitewood - The eroded Whitewood soil, as the name implies, is a Whitewood soil whose topsoil has been partially or almost totally removed by erosion. It occurs on upper slopes and knolls and is easily recognized by its lighter, grayish-colored A horizon. This A horizon is low in organic matter, is often calcareous and directly overlies the C horizon.

WHITESAND (Ws) SOILS

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Leached Poorly Drained Soils - Leached poorly drained soils occur mainly in sloughs and low-lying depressional areas which collect runoff from heavy rains and snowmelt. They are usually wet for the early part of the growing season and are often subject to flooding throughout the year after significant rainfall; drainage is restricted by a dense, slowly permeable subsoil. These soils have a platy (leached), light-colored layer below the surface underlain by a dense, grayish-brown B horizon that becomes very hard when dry. When cultivated, the upper layers are mixed, resulting in a gray-colored surface which is subject to crusting. The underlying B and C horizons have dull colors and reddish spots and streaks, indicative of formation under poorly drained conditions.

Agricultural Properties of Whitewood Soils

Whitewood soils are good agricultural soils of capability class 2. Loam and sandy loam-textured Whitewood soils have a slight moisture deficit as their main limitation, which is a result of the subhumid regional climate and a moderate water-holding capacity. These soils may be further down-rated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to an individual delineation. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

The degree of leaching that has occurred in these soils is important when evaluating their productivity. Those that are only slightly leached are nearly equal to similar soils developed in the Black soil zone, whereas the more strongly leached Whitewood soils are lower in organic matter and natural fertility and have poorer structure. Some of the more strongly leached soils may be susceptible to crusting after rains and the dense B horizon may restrict rooting.

Stones, in many cases, pose a hindrance to cultivation and regular clearing is often required. Wind erosion has not been a serious problem but can be expected to worsen when all the land is cleared and cultivated. Water erosion is generally not a serious problem, except on moderately or strongly sloping landscapes. Management practices that include forages in the crop rotations should be used to enhance soil organic matter content and improve soil structure.

Whitesand soils are Black soils that have formed in gravelly fluvial materials. Surface textures usually range from loamy sand to sandy loam or loam, however, in some areas, gravelly phases of these textures may be more common.

Stoniness is variable, ranging from nonstony in some areas to moderately stony in others. Whitesand soils are usually associated with undulating landscapes having very gentle to gentle slopes and hummocky landscapes with gentle to moderate slopes.

Whitesand soils occur in complex with soils of many other associations. In most of these complexes, the Whitesand soils can occur in any slope position.

Kinds of Whitesand Soils

Orthic Whitesand - The orthic Whitesand soil usually occurs on mid- and lower slope positions but may extend onto upper slopes and knolls in some landscapes. It is a rapidly drained soil with a black A horizon, 10 to 20 cm thick, underlain by a brownish-colored B horizon and a lighter-colored, weakly calcareous C horizon. The C horizon is often much more gravelly than either the A or B horizon.

Calcareous Whitesand - The calcareous Whitesand soil usually occurs on upper slopes and knolls. It is a rapidly drained soil characterized by a thin, usually calcareous A horizon and a thin, calcareous B horizon. These soils are often subject to erosion and are frequently lighter in color than the surrounding soils on mid- and lower slopes.

Eroded Whitesand - The eroded Whitesand soil usually occurs on upper slopes and knolls. It is a rapidly drained soil that has had part or almost all of its topsoil removed by erosion. A lower organic matter content and the presence of carbonates in the A horizon often result in these soils having a much lighter surface color in cultivated fields than the surrounding Whitesand soils.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Whitesand Soils

Whitesand soils are fair to poor agricultural soils of capability classes 3 and 4, respectively. Their sandy surface textures, coupled with gravelly subsurface textures, result in

very low to low water-holding capacities. The finer-textured Whitesand soils, the loams to fine sandy loams, are rated as class 3, while the coarser-textured Whitesand soils, the sandy loams to loamy sands, are rated as capability class 4 due primarily to their lower water-holding capacity. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. For example, exceedingly stony Whitesand soils, which cannot be cultivated unless considerable clearing is done, are rated as capability class 5. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

The low to moderate organic matter content and low inherent fertility of Whitesand soils contribute to the low potential that these soils have for the production of annual crops. Whitesand soils are also highly susceptible to wind erosion. Their sandy textures and weak structure make it difficult to maintain a cloddy surface that is resistant to wind erosion. Where these soils are cultivated, it is recommended that soil conservation practices, such as maintenance of crop residues, through reduced tillage or leaving stubble standing, strip cropping and frequent inclusion of forages, be utilized to control soil erosion and maintain or enhance the organic matter content of the soil. Water erosion is not a problem on most of these soils because of their high infiltration rate. Stoniness varies from one area to another and even within the same area of Whitesand soils. Some areas are stone free while others have sufficient stones to require periodic removal if the soils are to be cultivated. At best, these soils are marginal for arable crops, with many areas best utilized for forage production or native grazing.

WAITVILLE (Wv) SOILS

Waitville soils are Gray soils that have formed in loamy glacial till, in areas where wooded vegetation has had a strong influence on soil formation. Soils formed under these conditions are highly leached, resulting in lower organic matter levels and, hence, dark-gray to gray surface colors. Surface textures range from loam to sandy loam.

Waitville soils are typically slightly to moderately stony but range to exceedingly stony in some areas. They usually occur on hummocky landscapes with slopes ranging from gentle to steep.

In most complexes in the Gray soil zone, the Waitville soils tend to occur on the mid- and upper slope positions, however, when these soils occur in complexes in the Black soil zone, the Waitville soils usually occur on the more moist and strongly leached lower slopes.

Kinds of Waitville Soils

Gray Wooded Waitville Soil - The gray wooded Waitville soil usually occupies upper slope positions, however, it may extend to all slope positions in some landscapes. It is a well-drained soil which, under forested conditions, is charac-

terized by the presence of a very thin, dark-colored surface horizon below the forest litter, underlain by a gray to grayish-brown, strongly leached horizon with platy structure. Upon cultivation, part or all of this leached horizon is incorporated into the plow layer, producing a light-gray surface. Below these horizons is a relatively thick, dark-brown to dark grayish-brown B horizon that usually has a strong (hard), angular blocky to prismatic structure due to an accumulation of clay leached from upper horizons. The B horizon, in turn, is underlain by a grayish-colored, moderately calcareous C horizon.

Gleyed Gray Wooded Waitville - The gleyed gray wooded Waitville soil commonly occurs on mid- to lower slope positions. It is a moderately well- to imperfectly drained soil characterized by a light-gray A horizon, 8 to 15 cm thick, underlain by a light-gray, leached horizon with platy structure. This horizon is underlain by a brownish-colored B horizon that has a distinct, angular blocky structure when dry, and a grayish-colored, moderately calcareous C horizon. The B and C horizons often have dull colors and reddish spots and stains, indicative of imperfect soil drainage.

Dark Gray Wooded Waitville - The dark gray wooded Waitville soil usually occurs on mid- to lower slope positions. It is a well- to moderately well-drained soil which, under forested conditions, is characterized by the presence of a thin, dark-colored surface horizon below the forest litter, underlain by a gray to grayish-brown, leached horizon with platy structure. Upon cultivation, part or all of this leached horizon is incorporated into the plow layer, producing a dark-gray surface. Below these horizons is a relatively thick, dark-brown to dark grayish-brown B horizon that usually has a strong (hard), angular blocky to prismatic structure due to an enrichment of clay leached from upper horizons. The B horizon is underlain, in turn, by a grayish-colored, moderately calcareous C horizon.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Leached Poorly Drained Soils - Leached poorly drained soils occur mainly in sloughs and low-lying depressional areas which collect runoff from heavy rains and snowmelt. They are usually wet for the early part of the growing season and are often subject to flooding throughout the year after significant rainfall; drainage is restricted by a dense, slowly permeable subsoil. These soils have a platy (leached), light-colored layer below the surface underlain by a dense, gray-

ish-brown B horizon that becomes very hard when dry. When cultivated, the upper layers are mixed, resulting in a light gray-colored surface which is subject to crusting. The underlying B and C horizons have dull colors and reddish spots and streaks, indicative of formation under poorly drained conditions.

Agricultural Capability of Waitville Soils

Waitville soils are fair agricultural soils of capability class 3. They have a slight moisture deficit due to the regional subhumid climate and a moderate water-holding capacity. The main limitations of these soils are related to soil structure. The low organic matter content of some of these soils results in a structure that makes seedbed preparation difficult and also makes the soil susceptible to crusting after heavy rains, resulting in poor seedling emergence especially for small-seeded crops. They have a dense B horizon that may restrict water infiltration and root penetration. These limitations are most strongly expressed in the gray wooded soils. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. Ratings for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Waitville soils are low in available phosphorus and high in available potassium. They are moderately acid to neutral in reaction, however, the acidity does not usually affect the yield of cereal crops. However, there may be local areas with moderate acidity where the growth of some sensitive crops, such as alfalfa, may be inhibited.

Stones can pose a serious hindrance to cultivation and annual clearing is often required. Water erosion is not a serious problem, but can be during periods of intense rainfall due to the relatively low infiltration rate of these soils. Management practices, such as cultivation across slopes and grassing of runways in affected areas, should be followed as much as possible. Wind erosion can be a problem unless conservation practices are followed. Such practices include maintenance of crop residues through reduced tillage or leaving stubble standing, strip cropping or the establishment of forages in seriously-affected areas. The use of legumes in crop rotations will help to increase the soil organic matter content and, thus, improve the surface structure of these soils.

WETLAND (Wz) SOILS

Wetland soils are poorly drained soils formed in a mixture of materials associated with depressional areas. Most occur as sloughs, too small to show separately on the soil map, and are included in the map units of the surrounding upland soils. Only the larger areas have been delineated on the map. The Wetland soils are made up of a variety of soils which are referred to collectively as poorly drained soils. All are wet for at least a portion of the growing season and many remain flooded for much or all of the growing season.

Kinds of Wetland Soils

Poorly Drained Wetland - Poorly drained Wetland soils occur in sloughs and, occasionally, in the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. The A horizon may be thin, thick and dark-colored, or leached and gray or grayish-brown in color. Peaty poorly drained soils have a layer of peaty material, from 15 to 40 cm thick, overlying mineral materials. Wetland soils have drab subsurface colors that are often dotted with reddish spots and streaks. Some of these poorly drained soils are also saline and carbonated. Most of the poorly drained Wetland soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought.

Agricultural Properties of Wetland Soils

All areas of Wetland soils have little or no potential for arable agriculture. Most have an agricultural capability rating of class 5 or lower. Some areas are suitable for tame or native forage production or as native grazing land. Those areas indicated on the soil map with the symbol Wz1 have the highest potential. These are areas that usually become dry at some point in the growing season and have potential for forage production. Some of the shallower ones may be arable during extended dry periods and will have some potential for the production of tame hay, or early maturing, moisture-tolerant crops. Those areas indicated on the soil map with the symbol Wz2 usually have at least central portions remaining flooded for all of the growing season. In these areas, only the outer margins have any potential for the production of native hay. Those areas indicated on the soil map with the symbol Wz3 usually remain flooded for all of the growing season and have little or no potential for agricultural use, even as grazing land.

YORKTON (Yk) SOILS

Yorkton soils are Thick Black soils that have formed in highly calcareous, loamy glacial till, in more humid areas of the Black soil zone, where soils with thicker-than-normal surface horizons have formed. They are distinguished from Oxbow soils by having a surface horizon that is greater than 20 cm in thickness. Surface textures are predominantly loam and clay loam but can range from sandy loam to silty clay loam.

Yorkton soils are usually slightly to very stony and are commonly associated with hummocky landscapes with slopes ranging from gentle to moderate although undulating landscapes with very gentle slopes are common in some areas.

The Yorkton soils occur in complex with many other soil associations. In complexes with other soils formed in glacial till, they usually occur on the mid- to lower slopes, however, when in complex with soils formed in lacustrine or fluvial

materials, they often occur on the mid- to upper slope positions.

Kinds of Yorkton Soils

Orthic Yorkton - The orthic Yorkton soil occurs mainly on lower slopes but may extend onto mid- and upper slopes in some landscapes. It is a well- to moderately well-drained soil, with a black A horizon 20 to 33 cm thick, underlain by a brownish-colored B horizon and a grayish-colored, strongly calcareous C horizon.

Gleyed Orthic Yorkton - The gleyed orthic Yorkton soil usually occurs on lower slope positions although it may extend into depressions in some areas. It is a moderately well- to imperfectly drained soil with a black A horizon, 21 to 32 cm thick, underlain by a lime-free B horizon and a strongly calcareous C horizon that often have dull colors and reddish spots and stains, indicative of imperfect soil drainage.

Calcareous Yorkton - The calcareous Yorkton soil occurs mainly on well-drained upper slopes but can also occur on some moderately well-drained lower slopes. It has a black, often calcareous A horizon, 21 to 33 cm thick, that is usually underlain by a brownish to grayish-brown, calcareous B horizon and a grayish-colored, strongly calcareous C horizon.

Weakly Developed Yorkton - The weakly developed Yorkton soil usually occurs on upper slopes and knolls. It is a well-drained soil, with a black, usually calcareous A horizon, 21 to 34 cm thick, underlain by a grayish-colored, strongly calcareous C horizon.

Carbonated Yorkton - The carbonated Yorkton soil occurs mainly on lower slopes, frequently surrounding sloughs or poorly drained depressions, but may also occur in imperfectly drained, nearly level or very gently sloping areas. The black-colored A horizon is highly calcareous and is usually thicker than the A horizon of the Yorkton soils on mid- and upper slopes. The underlying B, if present, and C horizons are highly calcareous and often have drab colors and reddish spots and stains, indicative of imperfect soil drainage. In some areas, these soils may be affected to some degree by salinity.

Saline Yorkton - Saline Yorkton soils usually occur on lower slopes and in imperfectly drained depressions where the upward movement of subsurface water results in the deposition of salts in upper horizons. Consequently, soluble salts are usually present within 50 cm of the surface. The salts occur as a white surface crust, or as small, white specks within the soil, although salts may not always be visible. Dull colors and reddish spots and stains, indicative of imperfect

soil drainage, are often present in the subsoil. It frequently occurs intermixed with the carbonated Yorkton soil.

Poorly Drained Soils - Poorly drained soils represent a variety of wet soils. They occur mainly in sloughs and, occasionally, on the bottom of small drainage channels and low-lying depressional areas. They occur in areas that collect runoff from heavy rains and snowmelt, and usually remain wet for much of the growing season. They often have thick, dark-colored A horizons and drab subsurface colors that include reddish spots and streaks. In some cases, they may have a peaty layer 15 to 40 cm thick on the surface. In other areas, there may be a very light-gray, strongly leached horizon below the darker A horizon. Most of these soils are not cultivated unless drained, although some may become dry enough to cultivate during periods of prolonged drought. Due to their location in the landscape, some of these soils have become saline and/or carbonated.

Agricultural Properties of Yorkton Soils

The best Yorkton soils, those with loam and clay loam surface textures, are very good agricultural soils of capability class 1. The less common sandy loam-textured Yorkton soil has a capability of class 2 and is considered to be a good agricultural soil. A slight moisture deficit, imparted by a moderate water-holding capacity and the subhumid regional climate, is the main limitation to the production of common field crops. These soils may be further downrated based on other soil and landscape limitations (i.e. salinity, topography, stones, etc.) that are peculiar to individual delineations. Limitations for each delineation are listed under the heading "Agricultural Capability" in the Interpretive Data Tables section of this report.

Yorkton soils are naturally fertile. They have a moderately high organic matter content, are neutral to moderately alkaline in reaction, and are reasonably easy to keep in good tilth. Most fields require periodic clearing of stones but the stones are generally not a serious problem. Wind erosion has not been serious but may have occurred to some extent on some knolls. Water erosion is generally not a problem except on sloping lands, particularly in areas with long slopes. Soil conservation practices, such as maintenance of crop residues through reduced tillage or leaving stubble standing, grassing runways, cultivation across slopes and establishment of forages, may be used to control soil erosion. Saline soils occur infrequently, however, where they do occur, they are usually associated with depressional areas. Consequently, crop yields in areas adjacent to poorly drained depressions may be reduced in some years.

4. SOIL INTERPRETATIONS

4.1 SALINITY

Saline soils occur sporadically throughout the agricultural area of Saskatchewan. These soils contain sufficient water soluble salts to inhibit the uptake of moisture by plants, resulting in moisture stress and reduced plant growth. The presence of saline soils can often be recognized by bare spots in the crop or by uneven stands of grain or forage. Very strongly saline soils usually develop a white surface crust during dry weather. Where less salt is present, the soil is grayish in color when dry and the subsoil often has streaks or specks of salt at a depth of 5 to 25 cm or deeper. In weakly saline or moderately saline soils that are very wet, it may not be possible to see the salt.

Development of Saline Soils

Saline soils result almost invariably from the movement of salts carried by groundwater and subsequent concentration in the soil upon evaporation of this water at or near the soil surface.

Soluble salts are present in the parent materials of all soils as the result of on-going natural, chemical and physical weathering processes. When the amount of water evaporating from the soil is greater than the amount infiltrating, salts may accumulate in the soil and may result in saline soils. Areas are subject to soil salinization where water tables are high and the amount of infiltration of precipitation is limited. In most cases, this is a natural process which has been going on since the time of deglaciation. Agriculture has, however, aggravated the problem in some areas by the use of cropping systems that are not as water efficient as the natural prairie.

Management of Saline Soils

Management of saline soils requires the effective management of soil water in both the saline and nonsaline parts of the landscape. In terms of water management for soil salinity control this means making the most efficient use of soil moisture possible. Extending the cropping rotation or continuously cropping in nonsaline areas will cycle more precipitation through crops rather than allowing it to reach the water table where it may contribute to salinity in some other location. Leaving stubble standing promotes a more even distribution of snow cover reducing the amount that blows off the land into large snow drifts or depressions where, upon melting, it has a greater chance of infiltrating to the watertable. Saline soils should be seeded to long-term forage or continuously cropped with crops having the appro-

prate degree of salt tolerance. The objective in the saline areas is to reduce the amount of evaporation from the soil surface, lower the watertable level and move salts downward with infiltrating precipitation.

Table 1. The relative tolerance of common field crops to soil salinity. (Differences of one or two places in the ranking may not be significant.)

Degree of Salinity Tolerated			
	Nonsaline	Moderately Saline	Strongly to Very Strongly Saline
Annual Field Crops			
Increasing ↓ Tolerance	Soybeans	Canola	Barley may produce some crop but this land best suited to tolerant forages.
	Field Beans	Mustard	
	Faba Beans	Wheat	
	Peas	Flax	
	Corn	Fall Rye*	
	Sunflowers	Oats	
		Barley* Sugar Beets	
Forage Crops			
Increasing ↓ Tolerance	Red Clover	Reed Canary	Altai Wild Rye
	Alsike	Meadow Fescue	Russian Wild Rye
	Timothy	Intermediate Wheat	Slender Wheatgrass ^b
		Crested Wheat	Tall Wheatgrass ^b
		Brome	
		Alfalfa	
		Sweetclover ^a	

* These crops not tolerant of flooding, which is common in some saline areas.

^b Under dry conditions slender wheatgrass is more tolerant than tall wheatgrass.

For more information on saline soils and their management, see the publication **The Nature and Management of Salt Affected Land in Saskatchewan** by Saskatchewan Agriculture, Soils and Crops Branch.

Explanation of the Salinity Symbol

The soil salinity symbol is made up of three components indicating the extent of saline soils, the degree of the salts in the saline soils, and the position in the landscape occupied by the saline soils within the delineation and is based on field observation alone.

Example: 1WPA	1 - Extent Class
	W - Degree Class
	PA - Landscape Position

Soil Salinity Extent Class Limits

Table 2. Soil salinity extent class limits.

Extent Class	% Of Area Affected
0	0
1	0 - 3
2	3 - 10
3	10 - 20
4	20 - 40
5	40 - 70
6	> 70

Soil Salinity Degree

Table 3. Description of soil salinity degree classes.

Salinity Degree	Electrical Conductivity of 0-60 cm depth (mS/cm)	Effect on Crop Growth and Estimate of Potential Yield Loss
Nonsaline	0 - 2	There are no visible effects of salts on the growth of crops. No yield loss.
(W) Weak	2 - 4	Yields of very sensitive crops may be restricted. Cereals are generally unaffected.
(M) Moderate	4 - 8	Yields of many crops are restricted. Wheat yields may be reduced by 30%.
(S) Strong	8 - 16	Only tolerant crops yield satisfactorily. Wheat yields may be reduced by 60%.
(V) Very Strong	16+	Only a few very tolerant crops yield satisfactorily. Wheat yields may be reduced by 80-100%.

Note: Electrical conductivity values based on a saturated paste extract.

Yield loss estimates are based on recent research and only apply to the saline soils, not to the entire delineated area.

Landscape Position

The landscape position describes where in the landscape saline soils occur. In some areas, saline soils occur in more than one landscape position. These situations are indicated by the use of two letters.

Table 4. Description of landscape position symbol.

Symbol	Description
P	Saline soils occur on the edges of depressions, sloughs or runways. All soils in the bottoms of the depressions are leached and nonsaline.
A	Saline soils occur throughout the bottoms of depressions and sloughs.
D	Saline soils extend throughout the bottoms of dissections and small runways.
S	Saline soils occur on the sides of hills and slopes well above any slough or depression.
I	Saline soil parent materials within 60 cm of the soil surface generally occur on knolls and upper slopes.

4.2 IRRIGATION SUITABILITY

The irrigation suitability rating is based on soil and landscape characteristics. The suitability rating uses limiting factors to predict the potential landscape-water-crop interaction. It also considers the potential long-term consequences of irrigation such that the soil will remain permanently productive while being irrigated. It does not consider water availability and quality, climate, or economics. Within any one map delineation there may be smaller soil areas that have a higher or lower irrigation suitability than that indicated by the map symbol. **Any decision regarding irrigation should be made only after a field-specific examination is made.**

Symbol Interpretation

The combination of soil and landscape categories (Table 5), based upon the most limiting features present (Tables 6 and 7), determine the irrigation class and suitability rating (Table 8).

A maximum of three limitations are shown in the symbol. An ideal soil area to be used for irrigation will have the following characteristics:

- medium texture
- uniform texture vertically and horizontally
- uniformly well drained
- nonsaline
- permeable
- nearly level
- nonstony

Table 5. Soil and landscape categories.

Soil Category	Landscape Category	Description
1	A	nonlimiting
2	B	slightly limiting
3	C	moderately limiting
4	D	severely limiting

Irrigation Symbol

example: 2Cmvt₁

2C - Irrigation class

m - Soil limitations

v, t₁ - Landscape limitations

The example above indicates that the area in question has slight limitations (2) due to soil factors (m) and moderate limitations (C) due to landscape factors (v, t₁). This area, therefore, has a fair suitability rating (Table 8).

Symbol Evaluation

Excellent to good areas (Table 8) can usually be considered irrigable. Fair areas are marginally suitable for irrigation providing adequate management exists such that the soil and adjacent areas are not affected adversely by water application. Poor soils can usually be considered nonirrigable. The rating is given for the area based on soil characteristics in the upper 1.2 m and the main landscape features in the area. Depending on the type and severity of the limitation, it may be advisable to investigate an area further. Portions of the total area may also be significantly better or poorer than the general rating would indicate. For example, within a poor area with steep slopes, there may be areas of gentler topography that may be suitable for small scale irrigation if the detailed examination indicates that this smaller area is otherwise suitable.

Decision to Irrigate

The cost of irrigation development can be expected to increase with less suitable soils. The suitability rating does not take into account important factors such as climate, agronomy, availability of water, or economics in determining the feasibility of an irrigation project. If a field is indicated to be suitable for irrigation based on the information presented in this report, then an onsite inspection should be made. Other factors not used in this rating should also be considered during a site specific examination. These include geological uniformity to 3 m, local relief, depth to bedrock, drainability, sodicity, organic matter content and surface crusting potential. These factors may affect the suitability to some degree in terms of the type of irrigation system that can be used, the type and amount of surface preparation needed, the response of the soil and crop to applied water, and the type of management needed. A decision can then be made whether to irrigate if economic conditions are suitable and an adequate source of water is available.

Irrigation can lead to improved stability and flexibility in farm production through improved reliability of water application. Although maximum yields may be attainable only through irrigation, assuming adequate management, other climatic considerations may affect the feasibility. Climatic factors may limit the range of crops that can be grown due to heat or growing season limitations. In higher rainfall areas of the province, irrigation water may only be, in many years, a minor supplemental source of water that may not be needed every year. In these cases, the increased returns through higher yields, in some years, may not justify the expense of development. In dry regions where the risk of crop failure due to drought is relatively high and the range of crops that can be grown is lower, irrigation water may be a potentially important source of moisture needed for crop growth.

Table 6. Landscape limitations.

Symbol	Description
a	Impact on Target Areas - refers to the hazard resulting from the impact of applied irrigation water to the irrigated area. Impacts may include such effects as higher water tables, wetter soils, and increases in soil salinity.
c	Impact on Nontarget Areas - refers to the hazard resulting from the impact of applied irrigation water on an adjacent nonirrigated area. The hazards may include such effects as higher water tables, wetter soils, development or build-up of saline areas, or flooding and sedimentation caused by runoff.
i	Inundation - refers to the frequency of flooding. The inundation hazard is used mainly on areas adjacent to rivers.
p	Stones - refers to the amount of stone present on the surface and in the soil. Stones may reduce the available water-holding capacity of the soil, increase development costs and restrict the types of crops that may be grown.
t	Slope - refers to the presence of simple slopes (t_1) in undulating landscapes, or complex slopes (t_2) in hummocky or inclined landscapes. Complex slopes are often more limiting than simple slopes. Topography may affect the type of irrigation system, design and management required.
v	Horizontal Variability - refers to the horizontal variations caused by texture, soil structure, and landscape pattern that may result in the surface ponding of irrigated soils.

Table 7. Soil limitations.

Symbol	Description
d	Structure - soil structural properties that restrict root and water penetration. Commonly used with soils that have a dense B horizon and an A horizon that is subject to crusting.
g	Geological Uniformity - the uniformity of the soil texture with depth. The greater the textural difference between the surface and subsoil, the greater the potential for the development of perched water tables and lateral water movement.
k	Hydraulic Conductivity - the rate at which water moves through a saturated soil. Used mainly on soil areas that swell upon wetting, restricting water movement through the soil.
m	Available Water-Holding Capacity - the amount of water held by a soil that can be absorbed by plants. Coarse-textured soils with a low water-holding capacity are considered to be relatively inefficient for irrigation, as compared to medium-textured soils. Soils with this limitation also have relatively high hydraulic conductivities and intake rates.
q	Intake Rate - the rate of movement of water into the soil. It is closely associated with hydraulic conductivity which controls the rate at which water moves through the soil, and thus affects the rate at which water is able to enter the soil. Usually used on fine-textured soils that have relatively low intake rates requiring relatively light water application rates.
r	Depth to Bedrock - the presence of near-surface bedrock. Perched water tables may form, resulting in poor drainage and lateral movement of water and salts.
s	Salinity - the presence of soluble salts that may affect the growth of crops. The potential exists for lower yields, or for lateral salt movement into adjacent areas.
w	Drainage - the rate of removal of water from a soil in relation to supply. Indicates areas of mainly poorly drained soils.

Table 8. Irrigation suitability classes.

Class	Rating	Degree of Limitation	Description
1A	Excellent	No soil or landscape limitations	These soils are medium textured, well drained and hold adequate available moisture. Topography is level to nearly level. Gravity irrigation methods may be feasible.
2A 2B 1B	Good	Slight soil and/or landscape limitations	The range of crops that can be grown may be limited. As well, higher development inputs and management skills are required. Sprinkler irrigation is usually the only feasible method of water application.
3A 3B 3C 1C 2C	Fair	Moderate soil and/or landscape limitations	Limitations reduce the range of crops that may be grown and increase development and improvement costs. Management may include special conservation techniques to minimize soil erosion, limit salt movement, limit water table build-up or flooding of depressional areas. Sprinkler irrigation is usually the only feasible method of water application.
4A 4B 4C 4D 1D 2D 3D	Poor	Severe soil and/or landscape limitations	Limitations generally result in a soil that is unsuitable for sustained irrigation. Some lands may have limited potential when special crops, irrigation systems, and soil and water conservation techniques are used.

4.3 STONES

The stones rating is an estimation of the average severity of stoniness in a delineation. The estimation is based on the amount of stone clearing activity required and is related to the number and size of stones on the soil surface, number and size

of stone piles, and the soil parent material observed. The amount of stone clearing activity required is categorized into one of six stone severity classes listed in the table below.


Table 9. Stone classes.

Symbol	Description
S0	Nonstony.
S1	Slightly stony - stones seldom hinder cultivation. Light clearing is occasionally required.
S2	Moderately stony - stones are a moderate hindrance to cultivation. Annual clearing is usually required.
S3	Very stony - stones cause a serious hindrance to cultivation. Sufficient stones to require clearing on an annual basis.
S4	Excessively stony - stones prohibit cultivation or make clearing a major task. Cultivation is usually severely hindered, even after regular, heavy clearing.
U	Unclassified.

4.4 SOIL CAPABILITY FOR AGRICULTURE

The soil capability classification for agricultural use is an interpretive classification of soils based on limitations affecting their use for production of annual crops. These limitations are categorized according to **degree** or severity and **kind** of limitation. Degree of limitation is represented by

the capability class (numbers in the example below) and kind of limitation is represented by the capability subclass (letters in the example below). (The bracketed numbers in the example below indicate the percentage of each capability class present.) Capability classes and subclasses are briefly outlined below. A complete explanation of the system of soil capability classification for agriculture is contained in the publication, **A Guide to Soil Capability and Land Inventory Maps in Saskatchewan**.



Capability Class (Degree of Limitation)

The mineral soils of Saskatchewan are grouped into seven capability classes. Soils rated Classes 1 to 3 are considered suitable for sustained production of common cultivated field crops, those rated Class 4 are considered marginal for sustained production of common cultivated

field crops, those rated Class 5 are considered capable only of permanent pasture and hay production, those rated Class 6 are considered suitable only for use as native pasture, and those rated Class 7 are considered unsuitable for either the production of field crops or for use as native pasture.

Table 10. Description of capability classes.

CLASS 1	Soils in this class have no significant limitations in use for crops.
CLASS 2	Soils in this class have moderate limitations that restrict the range of crops or require moderate conservation practices.
CLASS 3	Soils in this class have moderately severe limitations that restrict the range of crops or require special conservation practices.
CLASS 4	Soils in this class have severe limitations that restrict the range of crops or require special conservation practices, or both.
CLASS 5	Soils in this class have very severe limitations that restrict their use to the production of native or tame species of perennial forage crops. Improvement practices are feasible.
CLASS 6	Soils in this class are capable of producing native forage crops only . Improvement practices are not feasible.
CLASS 7	Soils in this class have no capability for arable agriculture or permanent pasture .
CLASS O	Unimproved or virgin organic soils are not included in classes 1 to 7, and are designated by the letter 'O'.

Capability Subclass (Kind of Limitation)

The capability subclass represents a grouping of soils that have the same kind of limitations for crop production. If more than one limiting condition is recognized in a particular area, the subclasses are listed in order of their importance.

Table 11. Description of capability subclasses.

<u>Climatic Limitations</u> - Limitations due to climatic deficiencies.	
C	Depicts a moisture deficiency due to insufficient precipitation.
<u>Soil Limitations</u> - Limitations due to soil deficiencies are caused by adverse physical, chemical and morphological properties of the soil.	
D	Depicts adverse soil structure in the upper layers (A and B horizons) that affects the condition of the seedbed, prevents or restricts root growth and penetration, or adversely affects moisture permeability and percolation.
F	Depicts adverse fertility characteristics of soils having naturally low inherent fertility due to lack of available nutrients, high acidity or alkalinity, high calcium carbonate content or inadequate cation exchange capacity.
M	Depicts an insufficient soil water-holding capacity, due to the combined effects of the textural characteristics of the top 1 m and by the organic matter content of the surface horizon.
N	Depicts excessive soil salinity and applies to soils with either high alkalinity or a sufficient content of soluble salts to adversely affect crop growth or the range of crops which can be grown.
S	Depicts a variety of adverse soil characteristics. It is used in a collective sense in place of subclasses M, D, F and N, where more than two of them are present, or where two of these occur in addition to some other limitation.
<u>Landscape Limitations</u> - Limitations due to adverse characteristics of the soil landscape.	
T	Depicts a limitation in agricultural use of the soil as the result of unfavorable topography. It includes hazards to cultivation and cropping imposed by increasing degree of slope as well as by the irregularity of field pattern and lack of soil uniformity.
W	Depicts a limitation due to excess water caused by either poor soil drainage, a high groundwater table or to seepage and local runoff. It does not include limitations that are the result of flooding.
P	Depicts a limitation caused by excess stones and it applies to soils that are sufficiently stony that the difficulty of tillage, seeding and harvesting are significantly increased.
E	Depicts a limitation caused by actual damage from wind and/or water erosion.
I	Depicts a limitation due to inundation and applies to soils subjected to flooding by lakes or streams, but does not include local ponding in undrained depressions.
R	Depicts a limitation due to shallowness to bedrock and applies to soils where the rooting zone is restricted.
X	Soils having a moderate limitation due to the accumulative effect of two or more adverse characteristics of the soil and the landscape which singly are not serious enough to affect the class rating.

4.5 SURFACE pH

The pH scale, which ranges from 0 to 14 is used to indicate the relative acidity or alkalinity of a solution. A soil with a pH value of 7.0 is neutral, while one with a value less than 7.0 is acidic, and one with a value greater than 7.0 is alkaline.

The pH values indicated in the table in Section 6.0 are for the surface layer of soil which ranges in thickness from 10 to 20 cm. The soil pH was determined using a mixture of one part soil with one part distilled water. A soil with a pH between 6.5 and 7.5 provides the best environment for crop growth. Yields of sweet clover and alfalfa are reduced below

a pH of 6.0. A pH of 5.5 or less may reduce the yields of wheat, barley and canola.

Table 12. Surface pH classes.

pH Class	pH Range	Description
X	less than 5.5	Moderately acid
A	5.5 to 6.0	Slightly acid
B	6.1 to 6.7	Slightly acid to neutral
C	6.8 to 7.5	Neutral to slightly alkaline
D	greater than 7.5	Alkaline

Table 13. Definition of surface pH symbol.

Symbol	Percent Surface pH Class	Symbol	Percent Surface pH Class
X1	X ⁵ A ³ B ²	C1	C ⁵ B ⁴ D ¹
A0	A ⁵ B ⁵	C2	C ⁷ B ³
A1	A ⁷ B ² C ¹	C3	C ⁷ B ² D ¹
A2	A ⁵ B ² C ³	C4	C ⁹ D ¹
A3	A ³ B ⁴ C ³	C5	C ⁵ B ² D ³
A4	A ³ B ³ C ³ D ¹	C6	C ⁷ D ³
B0	B ⁷ A ² C ¹	C7	C ⁶ D ⁴
B1	B ⁴ C ⁴ A ²	D1	D ⁵ C ⁵
B2	B ⁷ C ³	D2	D ⁷ C ³
B3	B ⁵ C ⁵	D3	D ⁹ C ¹
B4	B ⁶ C ³ D ¹	D4	D ⁵ C ³ B ²
B5	B ⁷ A ³		

EXAMPLE

Symbol → B1 = B⁴C⁴A²

B⁴ - 40% of surface area has a pH in the "B" range (6.1 - 6.7)

C⁴ - 40% of surface area has a pH in the "C" range (6.8 - 7.5)

A² - 20% of surface area has a pH in the "A" range (5.5 - 6.0)

4.6 WETLANDS CLASSIFICATION


Wetlands are intermittent, semipermanent, or permanent waterbodies and include areas that have been drained as well as areas in which organic soils occur. Undisturbed wetlands are used extensively by wildlife for shelter, breeding habitat, and as feeding areas. Agricultural use of wetlands is often limited. However, some wetlands may provide native forage or hay for livestock, and trees may be used from some organic wetlands. If drained, these wetlands may support annual crop or hay production.

The size and type of wetland is often reflective of the permanence of a wetland. Generally the larger a wetland and the more open water there is, the more permanent it is. Nevertheless, wetlands will be influenced by recent climatic conditions. Due to the drought of the 1980's, for example, many wet meadows and shallow marshes have become dry enough to cultivate. In wetter years, it may be that many of these areas will become flooded and they may not be feasible to farm without installing drainage.

Wetlands occur in almost all areas, but are often not large enough to be mapped as individual areas. The wetlands classification attempts to indicate the types of wetlands that occur in each soil area mapped, as well as their approximate size and extent. It may, however, also be used to describe an individual wetland that is large enough to be delineated on the map. The wetland types can be identified using characteristic vegetation types, and the size and extent of wetlands distributed in the landscape is estimated using air photo interpretation.

Wetland Symbol

The wetlands symbol is composed of: an extent class, followed by a wetland type, a size class, and a water quality rating.



Wetland Types

W- Wet Meadow. Contains vegetation composed mainly of fine-textured grasses and sedges of low stature intermixed with various forbs. Occasionally willows will dominate the wet meadow. Normally wet meadows are flooded for only three or four weeks in the springtime.

- S - Shallow Marsh.** Contains vegetation composed of various intermediate-height grasses, sedges, and forbs. Flooding normally lasts until July or early August.
- E - Emergent Deep Marsh.** Contains vegetation composed of coarse grass-like plants such as bulrushes and cattails. Water normally persists into late summer and fall and occasionally throughout the winter.
- O - Open Water Marsh.** Contains an open water zone that occupies less than 75% of the wetland's diameter. Plants that occur in the open water area are either submerged rooted or floating. Water is normally persistent throughout the year, except in times of extreme drought.
- P - Shallow Open Water Wetland.** The shallow open water wetland has an open water zone that occupies greater than 75% of the wetland's diameter.
- A - Open Alkali Wetland.** Contains a high concentration of salts in the mineral material throughout the extent of the wetland. The length of time the wetland is flooded varies from a few weeks to months. When the wetland is dry, a salt crust usually forms on the soil surface.
- C - Cultivated Wetland.** Any wetland that has been altered by cultivation.
- B - Bog.** Wetlands having an accumulation of mosses and forest peat materials.
- F - Fen.** Wetlands associated with nutrient-rich groundwater, and having an accumulation of sedge residues.
- L - Lake.** Named lakes, and other large waterbodies that are predominantly open water and greater than 2 m in depth.
- N - No Wetlands.** Wetlands occupy less than 1% of the mapped area.

Wetland Extent and Size Classes

The proportion of the polygon that has wetlands is estimated and given an extent class. The main size classes of wetlands in the polygon is also determined.

Table 14. Wetland extent and size class limits.

Extent		Size	
Class	% Area	Class	Area (acres)
1	1 - 5	1	< 1
2	5 - 10	2	1 - 5
3	10 - 20	3	5 - 10
4	20 - 40	4	10 - 20
5	40 - 70	5	20 - 40
6	> 70	6	> 40

Water Quality

Quality of the water is estimated based upon the wetland vegetation.

- N - Fresh to Moderately Saline.** Water conductivity is less than 15,000 $\mu\text{s}/\text{cm}^3$.
- H - Saline to Highly Saline.** Water conductivity is greater than 15,000 $\mu\text{s}/\text{cm}^3$. Plants that are known to occur where it is saline are red samphire, alkali grass, and prairie bulrush.

4.7 WIND EROSION

Wind Erosion Potential

The calculation of wind erosion potential is based on the following formula:

$$E(p) = C \times T \times I \times K$$

E(p) Potential annual soil loss.

C **Climatic factor** (based on average wind velocity and temperature). Values are compiled from weather stations and are presented on a rural municipality basis.

T **Landscape factor** (based on slope class and surface form). Topography, including the differences in relief between one location and another, the direction, steepness and frequency of slopes, and the comparative roughness of the land's surface, has a pronounced effect on the potential erodibility of soils.

I **Soil erodibility factor** (based on texture). The relative proportions of sand, silt and clay present influence a soil's ability to absorb and retain moisture and, consequently, to form aggregates resistant to wind erosion. Coarse-textured soils have a "single grain" structure lacking sufficient amounts of silt and clay to bind individual sand particles together. Consequently, these soils are readily broken down and eroded by wind. Fine and medium-textured soils have a higher water-holding capacity and stronger surface attraction. This results in a good soil structure with a high degree of resistance to wind erosion.

K **Soil ridge roughness factor** (based on texture).

The E(p) values from the formula are used to predict a soil's susceptibility to wind erosion if the soil surface is bare (i.e. it is in summerfallow with no growth and no organic residue on the surface). Management practices and the actual amount of past wind erosion that has occurred are not considered.

Table 15. Wind erosion susceptibility classes.

Class	Susceptibility	Description
1	Very Low	Good soil management and average growing conditions will produce a crop with sufficient residue to protect these soils against wind erosion.
2	Low	Good soil management and average growing conditions may produce a crop with sufficient residue to protect these soils against wind erosion.
3	Moderate	Average growing conditions may not supply adequate residue to protect these soils against wind erosion. Enhanced soil management practices are necessary to control wind erosion.
4	High	Average growing conditions will not provide sufficient residue to protect these soils against wind erosion. Coarse-textured soils may be seeded to pasture or to forage crops to prevent severe degradation of the soil.
5	Very High	These soils should not be used for annual cropping, but rather for pasture and forage crops which will protect the surface from severe degradation.
6	Extremely High	These soils must be left in permanent pasture and are not capable of sustaining arable agriculture.
U	Unclassified	Unclassified areas (e.g. Wetlands).

4.8 WATER EROSION

Water Erosion Potential

The potential water erosion classes are obtained by using the Universal Soil Loss Equation. This equation is an erosion model developed in the United States to predict long term average soil losses from runoff. The equation is:

$$A = R \times K \times L \times S \times C \times P$$

- A** Computed loss per unit area (tons per acre per year).
- R** Rainfall erosivity factor (the amount and intensity of rainfall an area receives).
- K** Soil erodibility factor (calculated using several physical soil properties; texture, organic matter, infiltration rate and structure).

- L** Slope length factor.
- S** Slope steepness factor.
- C** Cover and management factor.
- P** Support practice factor.

Assessment of potential water erosion for a particular area is independent of current management practices and therefore the C and P factors in the equation are held constant.

When using this information, it should be remembered that the class assigned to an area is an estimation of potential erosion for the entire area and that individual soils may occur within the area that vary significantly from the assigned class.

Table 16. Water erosion susceptibility classes.

Class	Susceptibility	Description
1	Very Low	Conventional soil management will produce sufficient residue to protect the soil from water erosion.
2	Low	Conventional soil management and average growing conditions should produce sufficient residue to protect the soil from water erosion.
3	Moderate	Conventional farming practices will result in a steady loss of soil due to water erosion. Conservation practices should be instituted to prevent degradation of these soils.
4	High	Rapid loss of soil will occur unless conservation practices are instituted. All gullies in these areas should be grassed.
5	Very High	These soils should not be broken due to their water erosion hazard. If broken, perennial crops or permanent forage should replace annual crops.
U	Unclassified	Unclassified areas (e.g. Wetlands).
D or G Modifiers		If an area was observed to be gullied (G) or dissected (D) (dissections being shallow gullies that can be crossed with farm implements), these symbols were added to the erosion class symbol to indicate that higher rates of erosion may occur on the steeper slopes along the edges of the dissection or gully if they are left unprotected.

4.9 PAST WIND AND WATER EROSION

An erosion rating has been assigned to each soil area. This rating reflects the surveyor's best estimate of the extent and degree of erosion that has occurred in an area since cultivation. Areas that have never been cultivated usually have enough vegetative cover to protect the soil surface from erosion and, therefore, remain relatively unaffected. Some

uncultivated areas, however, do have clear evidence of recent erosion.

The rating system contains six classes with the degree of past wind and water erosion ranging from unaffected (W0) to very severe (W5). These classes, with the exception of W0 (unaffected), are assigned modifiers (G, K, B) which identify the type of erosion that has occurred. Wetlands, nonsoil areas and some uncultivated areas were not classified and are designated with the symbol 'U'.

Table 17. Past wind and water erosion classes.

Class	Description
U	Unclassified
W0	Unaffected. No evidence of past wind or water erosion.
W1	Weak. Soils are slightly eroded.
W1K	The knolls have slightly thinner A horizons and are lighter in color than midslopes. There is no noticeable thickening of the surface horizon on mid- to lower slopes.
W1B	Wind has removed part of the soil surface resulting in thinner A horizons. There is very little mixing of the A and B horizons and little sign of soil accumulation on mid- and lower slopes.
W1G	A few very shallow dissections are present indicating very slight evidence of water erosion.
W2	Moderate. Soils are moderately eroded.
W2K	Eroded knolls make up 5-15% of the area. The knolls are much lighter in color than midslopes. There is a noticeable thickening of the surface horizon on lower slopes due to accumulation of upper slope material.
W2B	Wind has removed part of the A horizon resulting in moderately thin A horizons. There is slight mixing of A and B horizons during tillage and some evidence of soil accumulation near fencelines and windbreaks.
W2G	Shallow dissections are present. The dissections may easily be crossed by farm implements and have little effect on cultivation. There is evidence of rill erosion (small channels a few centimeters deep, occurring after substantial rains or snowmelt).
W3	Strong. Soils are strongly eroded.
W3K	Eroded knolls make up 15-40% of the area. The knolls are much lighter in color than midslopes. A large portion of the A horizon has been removed and redistributed to lower slopes. On knolls, subsoil has been incorporated into the cultivated horizon.
W3B	Wind has removed a significant amount of the A horizon. Regular tillage results in a thorough mixing of the B horizon with the remaining A horizon. Accumulation of wind-blown material occurs along fencelines and windbreaks.
W3G	Distinct dissections are present. The dissections may be crossed by farm implements with some difficulty, and have a moderate effect on cultivation. These dissections should be seeded to grass to prevent further damage from erosion.
W4	Severe. Soils are severely eroded.
W4K	Eroded knolls make up 40-70% of the area. The eroded knolls are white in color, with light colors extending well onto the midslope position. Erosion has destroyed the soil profile on upper slopes.
W4B	Wind has removed most of the A horizon and frequently part of the B horizon. Occasional blowout areas are present, creating a very unstable surface.
W4G	Occasional shallow gullies are present. The gullies cannot be crossed by farm implements, and therefore, should not be cultivated for annual cropping. Reclamation for improved pasture is difficult unless erosion can be controlled.
W5	Very Severe. Soils are very severely eroded.
W5K	Eroded knolls make up greater than 70% of the area. The knolls and midslopes are white in color. Erosion has destroyed the soil profile on upper and midslope positions.
W5B	Wind has removed most of the soil profile. Blowout holes are numerous and easily carved into the subsoil or parent material. Areas between blowouts are deeply buried by eroded soil material. At best, this land should be utilized for native or improved pasture.
W5G	Deep gullies occur frequently. Soil profiles have been destroyed except in small areas between gullies. These areas should be permanently grassed. Reclamation of eroded areas is a difficult process.

4.10 SAND AND GRAVEL

The sand and gravel symbol shows the location of near surface sources of sandy and gravelly materials. The materials can range from mixtures of sand and silt to coarse gravelly sand. These materials may be used for concrete, sub-base for roads, traffic gravel or pervious borrow for fill

purposes. This symbol does not suggest whether any of these areas contain sands and gravels of sufficient volume or quality to enable commercial development.

The term sand refers to materials with greater than 50% sand and with less than 15% clay. Gravel refers to materials having a significant component of particles greater than 2 mm in diameter.

Table 18. Description of sand and gravel symbol.

Symbol	Description	Symbol	Description
SG0	No sandy or gravelly materials recognized	G3	Gravelly materials occupy 40-70% of landscape
S1	Sandy materials occupy 1-15% of landscape	SG3	Sandy and gravelly materials occupy 40-70% of landscape
G1	Gravelly materials occupy 1-15% of landscape	S4	Sandy materials occupy greater than 70% of landscape
SG1	Sandy and gravelly materials occupy 1-15% of landscape	G4	Gravelly materials occupy greater than 70% of landscape
S2	Sandy materials occupy 15-40% of landscape	SG4	Sandy and gravelly materials occupy greater than 70% of landscape
G2	Gravelly materials occupy 15-40% of landscape	U	Unclassified
SG2	Sandy and gravelly materials occupy 15-40% of landscape		
S3	Sandy materials occupy 40-70% of landscape		

Characteristics of Sand and Gravel Pits

The following table lists the legal location, and provides information on various characteristics, of sand and gravel pits in this municipality. The data was compiled by the Sas-

katchewan Research Council based on detailed field investigations by the Saskatchewan Department of Highways and Transportation.

Table 19. Gravel pit characteristics.

1/4	Legal Location				Tested Thickness (m)	Texture (%)			Comments
	Sec	Twp	Rg	Mer		Gravel*	Sand	Fines	
SE	15	40	18	2	2.7	21.9	72.6	5.4	Gravelly sand with few boulders.
S	32	42	18	2	2.2	36.2	62.1	1.6	Clean gravelly sand with some boulders.
<p>* "Gravel" refers to material greater than 5 mm in diameter (Industrial Classification). "Sand" refers to material greater than 0.071 mm and less than 5 mm in diameter. "Fines" refers to material less than 0.071 mm in diameter.</p>									

4.11 SOIL MOISTURE AND YIELD

An estimate of the amount of available soil moisture may be of value in decision-making regarding the seeding of stubble land and the level of fertilizer inputs required. In most cases, such decisions rely, in part, on an estimate of expected yield. Since yield is largely a function of soil moisture conditions, an estimate of the amount of soil moisture available to the crop over the growing season can be used to assess the probability of obtaining a given yield.

Calculation of Available Soil Moisture

To calculate the amount of available water in the soil, multiply the soil's available water-holding capacity (see Table 20) by the depth of moist soil.

Table 20. Available soil water-holding capacity in relation to soil texture.

Texture Class	Available water-holding capacity per unit depth of moist soil
Loamy sand	0.05
Sandy loam	0.08
Fine sandy loam	0.09
Very fine sandy loam	0.10
Sandy clay loam	0.12
Loam	0.14
Silt loam	0.16
Clay loam	0.17
Silty clay loam	0.20
Clay	0.21
Silty clay	0.22
Heavy clay	0.23

Example: If a loam-textured soil were moist to a depth of 500 mm, it would contain about 70 mm (500 x 0.14) of available water.

Estimation of Potential Yield

The amount of soil moisture available to the crop equals the amount stored in the soil prior to seeding, plus that received as precipitation during the growing season. Available soil moisture prior to seeding can be estimated using Table 20. The probability of receiving various amounts of precipitation over the growing season is given in Table 21.

Table 21. Probability (%) of receiving at least the indicated amounts of growing-season precipitation (May 15 to August 13).

Precipitation (mm) ^a	65	90	105	125	145	170	190	210	235	250
Probability (%)	>95	90	85	75	65	50	35	25	15	10

^a Precipitation data from Muenster weather station.

Example: If the probability of receiving 170 mm of precipitation were 50%, then at least 170 mm of precipitation could be expected in 5 out of 10 years.

To assess the probability of obtaining a given yield:

1. Estimate the available soil moisture prior to seeding.
2. Determine the total moisture requirements (Table 22).
3. Assess the probability of receiving enough precipitation during the growing season to make up the difference.

Example: If a loam-textured soil were moist to a depth of 500 mm, what would be the probability of obtaining a wheat yield of at least 1500 kg/ha?

1. Stored soil moisture = (500 x 0.14) = 70 mm.
2. Total moisture requirements = 200 mm.
3. Moisture required = (200 - 70) = 130 mm.
4. Probability of receiving at least 130 mm is about 72%.

Table 22. Estimated yields^b (kg/ha) in relation to available moisture requirements.

Crop	Moisture requirements (mm)										To convert kg/ha to bu/ac multiply by:
	125	150	175	200	225	250	275	300	325	350	
Wheat	600	900	1200	1500	1800	2100	2400	2700	3000	3300	.015
Oats	640	950	1270	1590	1910	2230	2540	2860	3180	3500	.026
Barley	740	1110	1480	1850	2210	2580	2950	3320	3690	4060	.019
Flax	360	500	660	830	990	1160	1320	1490	1650	1820	.016
Canola	360	540	720	900	1080	1270	1440	1620	1800	1980	.018

^b Yields are based on good management and reasonably normal seasonal climatic conditions, particularly with respect to the timeliness of rainfall events. Actual yields may exceed estimates under abnormally favorable conditions, or be below estimates due to weeds, disease, low fertility, or adverse climatic extremes.

5. ACREAGE FACTS

Rural Municipality of Pleasantdale, Number 398

Hectares Acres		Hectares Acres					
TOTAL AREA	85552	211400	SURFACE pH (Soil Acidity)			
SOIL CAPABILITY FOR AGRICULTURE				X (< 5.5)	0	0
Class 1	14070	34768	A (5.5 - 6.0)	437	1081
Class 2	19691	48656	B (6.1 - 6.7)	11032	27260
Class 3	22910	56610	C (6.8 - 7.5)	33649	83146
Class 4	9015	22277	D (> 7.5)	36306	89711
Class 5	6588	16278	SURFACE TEXTURE			
Class 6	4553	11251	Sands	3317	8196
Class 7	1728	4271	Sandy Loams	7219	17837
Class O	2868	7086	Loams	66779	165010
IRRIGATION SUITABILITY				Clay Loams	881	2176
Excellent	0	0	Clays	0	0
Good	33499	82776	Organics	3229	7980
Fair	23690	58537	WIND EROSION POTENTIAL			
Poor	24235	59885	Very Low	64572	159557
SALINITY				Low	13686	33817
Very Strong	87	215	Moderate	864	2136
Strong	157	387	High	79	195
Moderate	765	1890	Very High	0	0
Weak	195	482	Extremely High	0	0
None	84349	208426	WATER EROSION POTENTIAL			
SAND AND GRAVEL				Very Low	13770	34026
Sandy	2956	7305	Low	49581	122515
Sandy and Gravelly	2039	5039	Moderate	10616	26231
Gravelly	1955	4832	High	980	2422
STONES				Very High	1024	2531
Non- to Slightly Stony	39332	97189	WETLANDS AND POORLY DRAINED SOILS			
Moderately Stony	34419	85050	Open water and lakes	5841	14432
Very Stony	3953	9767	Wet, poorly drained soils	8805	21758
Excessively Stony	3720	9192				

6. INTERPRETIVE DATA TABLES

Area No.	Map Unit	Slope Class	Land Form	Surface Texture	Salinity	Irrigation	Stone Class	Agricultural Capability	pH	Wetlands Classification	Wind Eros.	Water Eros.	Past Eros.	Sand/Gravel	Acres (ac)
1	Mw6	2	ud	l-cl	1MPA	4Cwi	S1	3(7)W5(3)W	D3	5S3-N,4C3-N	1	1D	W1G	SG0	97.4
2	WhWv3	3	ud	l-cl	1MP	2Bdst1	S1	2(7)M3(3)D	C3	1C1-N	1	2D	W1K	SG0	89.6
3	YkOx15	2-3	u	l	1MPA	2Csv	S2	1(4)2(4)M3(2)W	C1	3C2-N,1S2-N	1	1	W1K	SG0	483.2
4	WvSy1	3	ud	l-sl	0	2Bdt1v	S2	4(6)MP3(4)D	C1	N	2	2D	W1G	S2	571.8
5	OxMe1	2-3	ud	l-sl	1WP	1Dv	S2	3(6)MW2(4)M	D1	2C2-N	2	1D	W1G	S2	966.9
6	Kitako Lake									6L6-N					1155.2
7	Mh1	2-3	u	scl	4SA	4Dwsv	S2	5(7)WN6(3)W	D3	6W6-N,2E3-N	1	1	U	SG0	112.3
8	WvWh1	3-4	urd	l-sl	0	2Bdt1v	S2	3(7)D2(2)DM4(1)M	C3	N	1	2D	W1G	SG0	789.1
9	MwSP1	2	h	l-o	0	4Cwi	S2	6(6)WO(4)	D3	5W6-N,5F6-N	1	1	U	SG0	631.5
10	OxYk7	3-2	ud	l	1WP	1Bt1v	S2	2(6)M1(3)4(1)W	D2	1S2-N,1C2-N	1	2D	W1G	SG1	126.3
11	WvWh1	3	hr	l	0	2Bdtv2	S2	2(7)DM3(3)D	C1	N	1	2	W1K	SG0	163.7
12	HmMe10T	3-2	ud	fl-sl	2MPA	3Dsv	S1	3(6)M4(4)MW	D3	2C2-N	2	2D	W1B	S2	82.2
13	Wz2	1	l	l	1MP	4Cwi	S2	6(4)W7(4)W5(2)W	C3	5W6-N,5E6-N	U	U	U	SG0	228.7
14	WhOx1	3	ud	l	1WP	1Bt1	S2	2(10)MD	D1	N	1	2D	W1G	SG0	169.7
15	WhWv3	3	urd	l	0	2Bdt1v	S2	2(6)DM3(4)D	C3	N	1	2D	W1G	SG0	208.0
16	WhOx3	3	ud	l	1WP	1Bt1v	S2	2(9)DM5(1)W	D1	2C2-N	1	2D	W1G	SG0	143.4
17	Mw6	2	u	cl	1WP	4Cwi	S1	6(6)W5(4)W	D3	5W5-N,5C4-N	1	1	W0	SG0	48.3
18	WvWh1	3	hd	l-vl	0	2Ddv	S1	2(5)DM3(5)DW	C4	2C1-N	2	2D	W1K	SG0	225.0
19	SP1F	1	H	o	1MP	4Cwi	S0	0(5)6(5)W	D3	5C6-N,5F6-N	1	U	W1B	SG0	48.6
20	WvWh1	3	ur	l	0	2Ddv	S1	2(5)DM3(5)DW	C4	3C1-N	1	2	W2K	SG0	273.8
21	McKnight Lake									6L6-N					92.3
22	WvWh5	4-5	h	l	1WP	2Ddv	S2	3(3)DT4(3)T5(4)WT	C3	3P4-N,2E2-N	1	3	W2K	SG0	781.4
23	Wz2	1	l	l	1WP	4Cwi	S2	6(10)W	C3	5E6-N,5W6-N	U	U	U	SG0	84.3
24	WvWh1	3-4	urd	l	0	2Bdt1v	S2	3(8)D2(2)DM	C3	N	1	2D	W1G	SG0	1350.5
25	Wz3	1	l	l	1WP	4Cwi	S2	7(10)W	C3	6L6-N	U	U	U	SG0	75.5
26	WhOx3	3	ud	l	1WP	1Bt1v	S2	2(9)MD5(1)W	D1	2W2-N	1	2D	W2G	SG0	191.2
27	WhOx3	3-2	ud	l-cl	1WP	1Cv	S2	2(6)MD1(2)4(2)WP	D1	2C2-N	1	2D	W1G	SG0	1072.1
28	Wv9	4-5	h	l	1WP	2Cdt2v	S2	3(4)D4(4)T5(2)W	C3	2S2-N,1W1-N	1	2	W2K	SG0	101.2
29	SyGb1	3	h	sl-gsl	1WP	2Bdmt2	S1	4(10)M	D1	1S1-N	2	2	W2B	SG4	288.8
30	Mh1	3	u	scl	1MP	4Dwv	S2	6(10)W	D3	6E6-N	1	2	U	SG0	107.6
31	WvWh1	3	u	l	1WP	2Bdt1v	S2	3(7)D2(3)D	C3	N	1	2	U	SG0	284.3
32	Av11	2	u	l-sl	6VA	4Dwsv	S2	6(10)NW	D3	6W6-H	1	2	U	SG0	268.8
33	OxMe1	2-3	ud	l-sl	1WP	1Dv	S2	3(6)MW2(4)M	D1	2C2-N	2	1D	W1G	S2	208.5
34	WhOx4	3	h	l-sl	1WP	1Cv	S2	2(8)MD3(2)MW	D2	2C1-N	1	2	W1K	SG0	161.6
35	WvSy1	3	ud	l-sl	0	2Bdt1v	S2	4(6)MP3(4)D	C1	N	2	2D	W1G	S2	304.0
36	OxWh1	2	u	l-sl	1WA	1Bv	S2	2(9)M3(1)M	D1	1C1-N	1	1	W1K	SG0	502.0
37	MwSP1	2	u	l-o	0	4Cwi	S2	0(5)5(5)W	D3	5W5-N,5F5-N	1	1	U	SG0	29.6
38	Mw6	2-3	u	l-scl	1WP	4Cwi	S2	6(7)W5(3)W	D3	6W6-N,3C5-N	1	1	W0	SG0	760.8
39	Ox3	3	ud	l	1WP	1Bt1	S2	2(10)M	D1	N	1	2D	W1G	SG0	155.5
40	WvWh1	3-4	urd	l	0	2Bdt1v	S3	3(8)DP5(2)P	C3	N	1	2D	W1G	SG0	459.5
41	Mh1	3	u	scl	2MPA	4Dwv	S2	5(7)W6(3)W	D3	6W6-N,3E4-N	1	2	U	SG0	589.2
42	Kinistino Lake									6L6-N					384.1
43	WvWh1	3	u	l-cl	2MP	3Bst1v	S2	3(7)D2(3)D	C3	1E2-N	1	2	W1K	SG0	259.9
44	OxWh5	3-2	ud	l	1MP	2Csv	S2	2(8)MD3(1)W4(1)W	D2	2C2-N,1W2-N	1	2D	W2G	SG0	198.9
45	OxHm6	3	u	sl	3SA	4Bst1v	S2	3(8)M4(2)NP	D3	1C2-N	2	1	W2B	SG1	41.2
46	SP1M	2	H	o	1WP	4Cwi	S0	6(10)W	D3	6C6-N	1	U	W2B	SG0	87.4
47	OxHm6	3	u	sl	3SA	4Bst1v	S2	3(8)M4(2)NP	D3	1C2-N	2	1	W2B	SG1	113.9
48	OxWh5	3-2	ud	l	1MP	2Csv	S2	2(8)MD3(1)W4(1)W	D2	2C2-N,1W2-N	1	2D	W2G	SG0	263.8
49	WvWh1	3-4	urd	l-sl	0	2Bdt1v	S2	3(6)D2(2)D4(2)M	C3	N	1	3D	W2G	SG0	1043.5
50	Rw	3-5	hd	l	1MD	4Dct2	S2	5(5)WE6(5)WE	D3	4W4-N,3E3-N	1	4D	U	SG0	191.7
51	WhWv3	3-2	ud	l	0	2Bdt1v	S2	2(8)MD3(2)D	C3	N	1	2D	W1G	SG0	134.1
52	WhOx3	3-2	u	l	1WP	1Bt1	S1	2(10)MD	D1	1C2-N	1	2	W1K	SG0	592.3
53	Hubbel Lake									6L6-N					412.9
54	WvWh1	3	urd	l	0	2Bdt1v	S2	3(7)D2(3)DM	C3	1P3-N	1	2D	W1G	SG0	1060.1
55	Lemoine Lake									6L6-N					306.6
56	WhWv3	3-2	ud	l	0	2Bdt1v	S1	2(6)MD3(4)D	C3	N	1	2D	W1G	SG0	653.8
57	WhOx4	3	h	l	1WP	1Bt2v	S2	2(9)MD5(1)W	D2	1W1-N,1C1-N	1	2	W1K	SG0	216.1
58	Mw6	3	h	l-cl	1WP	4Cwi	S2	6(6)W5(4)W	D3	5W6-N,5S6-N	1	2	W1K	SG0	339.2
59	WhOx4	3	h	l	1WP	1Bt2v	S2	2(9)MD5(1)W	D2	1W1-N,1C1-N	1	2	W1K	SG0	1009.7
60	Mw6	3	h	l-cl	1WP	4Cwi	S2	6(6)W5(4)W	D3	5W6-N,5S6-N	1	2	W1K	SG0	75.2
61	WhWv3	3	h	l	1WP	2Bdtv2	S2	2(7)MD3(2)D5(1)W	C3	2W2-N,1C2-N	1	2	W1K	SG0	1524.9
62	WhOx3	3	u	l	1WP	1Bt1v	S2	2(9)MD5(1)W	D1	2C2-N,1W2-N	1	2	W1K	SG0	351.6
63	Mw6	3	h	l-cl	1WP	4Cwi	S2	6(6)W5(4)W	D3	5W6-N,5S6-N	1	2	W1K	SG0	231.2
64	WhOx3	3	u	l	1WP	1Bt1v	S2	2(9)MD5(1)W	D1	2C2-N,1W2-N	1	2	W1K	SG0	60.1
65	WvWh1	3	hd	l	0	2Bdtv2	S2	3(6)D2(4)MD	C3	N	1	2D	W1G	SG0	479.8

Area No.	Map Unit	Slope Class	Land Form	Surface Texture	Salinity	Irrigation	Stone Class	Agricultural Capability	pH	Wetlands Classification	Wind Eros.	Water Eros.	Past Eros.	Sand/Gravel	Acreeage (ac)
66	Mw6	3	h	l-cl	1WP	4Cwi	S2	6(6)W5(4)W	D3	5W6-N,5S6-N	1	2	W1K	SG0	124.3
67	Mw6	3-2	u	l-cl	1WP	4Cwi	S2	6(7)W5(3)W	D3	6E6-N,3O6-N	1	2	U	SG0	49.9
68	WvWh5	3	h	l	0	2Ddv	S1	2(5)MD3(5)DW	C4	3C1-N,1S1-N	1	2	W2K	SG0	1266.3
69	Mw6	3-2	u	l-cl	1WP	4Cwi	S2	6(7)W5(3)W	D3	6E6-N,3O6-N	1	2	U	SG0	141.6
70	Charron Lake														
71	WvWh5	3	h	l	0	2Ddv	S1	2(5)MD3(5)DW	C4	3C1-N,1S1-N	1	2	W2K	SG0	89.6
72	WhWv2	3-4	h	l-cl	0	2Cdt2v	S1	2(5)D3(3)DT4(2)W	C3	3C1-N	1	2	W2K	SG0	286.1
73	Wz2	1	l	l	1WP	4Cwi	S2	6(6)W7(4)W	C3	6O6-N	U	U	U	SG0	85.6
74	WhWv4	3	h	l	0	2Ddv	S2	2(6)D3(4)DW	C3	3C1-N	1	2	W1K	SG0	303.0
75	Mw6	2-3	u	sil	3SA	4Cwi	S0	6(10)W	D3	6S6-N,3C3-N	1	1	W0	SG0	168.3
76	WhWv3	3	urd	l-cl	0	2Bdt1v	S1	2(9)MD3(1)D	C3	N	1	2D	W1G	SG0	1585.1
77	OxYk7	3-2	u	l	1MA	2Bst1v	S1	2(7)M1(2)3(1)W	D3	2C1-N	1	2	W1K	SG0	15.3
78	Wz2	1	l	l	1MP	4Cwi	S1	5(5)W6(5)W	D3	6S6-N	U	U	U	SG0	30.3
79	OxYk7	3	hd	l	2WA	2Bst2	S2	2(10)MT	D2	1W3-N	1	2D	W1K	SG0	225.9
80	YkNc7	3-2	u	l-sil	2MA	3Bst1v	S1	1(6)2(3)X3(1)NW	D3	2C2-N	1	2	W1K	SG0	1854.1
81	Rw	3	id	sil	2MPD	4Dct2	S2	5(7)WE3(3)WE	D3	4S6-N,3W6-N	1	3D	W3G	SG0	217.3
82	NcYk7	2-3	ud	l-sil	2MA	3Bst1v	S1	1(9)3(1)NW	D2	2C2-N	1	1D	W1G	SG0	232.1
83	Wz3	1	l	l-sil	1MP	4Cwi	S1	7(10)W	D2	6P6-N	U	U	U	SG0	116.6
84	NcYk7	2-3	ud	l-sil	2MA	3Bst1v	S1	1(9)3(1)NW	D2	2C2-N	1	1D	W1G	SG0	265.9
85	Rw	3	id	sil	2WPD	4Dct2	S2	5(7)WE3(3)TE	D3	4S6-N,3W6-N	1	3D	W4G	SG0	93.2
86	WhWv3	3	urd	l-cl	0	2Bdt1v	S1	2(9)MD3(1)D	C3	N	1	2D	W1G	SG0	1102.6
87	Wz2	1	l	l	1MP	4Cwi	S2	6(6)W7(4)W	C3	6S6-N	U	U	U	SG0	113.0
88	WvWh1	3	hrd	l	0	2Bdvt2	S2	3(7)D2(3)DM	C3	N	1	2D	W2G	SG0	295.3
89	Rw	3	id	sil	2WPD	4Dct2	S2	5(7)WE3(3)TE	D3	4S6-N,3W6-N	1	3D	W4G	SG0	36.7
90	Wz3	1	l	l	1MP	4Cwi	S2	7(8)W6(2)W	D2	6P6-N	U	U	U	SG0	168.5
91	Yk5	3-2	u	cl-l	3MA	4Dsv	S1	1(6)3(2)NW5(2)W	D3	3S2-N,2W2-N	1	2	W0	SG0	736.7
92	Wz3	1	l	l	1WP	4Cwi	S1	7(10)W	C3	6O6-N	U	U	U	SG0	276.0
93	WhWv3	3	ud	l	0	2Bdt1v	S1	2(9)MD3(1)D	C3	N	1	2D	W1G	SG0	929.6
94	Wz3	1	l	l-fl	1WP	4Cwi	S1	7(10)W	D1	6P6-N	U	U	U	SG0	48.7
95	Mw6	2-3	u	sil	3MPA	4Cwi	S0	3(10)WN	D3	6S6-N	1	1	W0	SG0	81.4
96	WhOx4	3-4	h	l	2WA	2Dsv	S1	2(7)MD3(3)TW	D2	2W2-N	1	2	W2K	SG0	390.1
97	OxWh1	2-3	ud	l	0	1Bt1	S1	2(10)MD	D1	N	1	1D	W1G	SG0	1031.4
98	Wz2	1	l	l-sil	2MP	4Cwi	S1	6(10)W	D2	6S6-N	U	U	U	SG0	76.8
99	Rw	4-5	id	sil	2WPD	4Dct2	S2	4(7)TE3(3)TE	D3	1C2-N	1	4D	W2G	SG0	77.6
100	NcYk7	2-3	ud	l-sil	2MA	3Bst1v	S1	1(9)3(1)NW	D2	2C2-N	1	1D	W1G	SG0	5790.1
101	WhWv3	3	hrd	l	0	2Bdvt2	S1	2(9)MD3(1)D	C3	N	1	2D	W1G	SG0	132.7
102	Yk11	3-2	u	l	2MA	3Bst1v	S1	1(9)3(1)NW	D3	2C2-N	1	2	W0	SG0	312.7
103	Wz2	1	l	l	2MP	4Cwi	S1	6(10)W	D3	6S6-N	U	U	U	SG0	78.8
104	Yk11	3-2	u	l	2MA	3Bst1v	S1	1(9)3(1)NW	D3	2C2-N	1	2	W0	SG0	249.2
105	Wz2	1	l	l-sil	2MP	4Cwi	S1	6(6)W7(4)W	D3	6O6-N	U	U	U	SG0	93.3
106	Rw	3-4	id	sil	2WPD	4Dct2	S2	5(7)WE3(3)WP	D3	3S3-N,2W2-N	1	3D	W3G	SG0	165.8
107	Yk2	3-2	ud	l	1WD	1Bt1	S1	1(10)	D2	1C1-N	1	2D	W1G	SG0	383.7
108	YkOx14	3	urd	l	1WD	1Bt1	S1	1(7)2(3)M	D2	N	1	2D	W1G	SG0	254.4
109	Wz2	1	l	l	1MP	4Cwi	S1	6(6)W7(4)W	D2	6O6-N	U	U	U	SG0	78.8
110	NcYk7	2-3	ud	l-sil	2MA	3Bst1v	S1	1(9)3(1)NW	D2	2C2-N	1	1D	W1G	SG0	519.1
111	YkNc7	3-2	u	l-sil	2MA	3Bst1v	S1	1(6)2(3)X3(1)NW	D3	2C2-N	1	2	W1K	SG0	471.5
112	Yk2	3-2	ud	l	1WD	1Bt1	S1	1(10)	D2	1C1-N	1	2D	W1G	SG0	1378.7
113	YkOx4	3-2	h	l	2MA	3Cst2	S2	2(9)X3(1)W	D2	3C1-N	1	1	W2K	SG0	291.5
114	Wz2	1	l	l	2MP	4Cwi	S2	7(6)W5(4)W	D2	6E6-N,3W3-N	U	U	U	SG0	52.8
115	YkOx4	3-2	h	l	2MA	3Cst2	S2	2(9)X3(1)W	D2	3C1-N	1	1	W2K	SG0	67.0
116	Wz3	1	l	l	1MP	4Cwi	S2	7(10)W	D2	6P6-N	U	U	U	SG0	85.3
117	YkOx14	3	urd	l	1WD	1Bt1	S1	1(7)2(3)M	D2	N	1	2D	W1G	SG0	352.4
118	Wz3	1	l	l	1WP	4Cwi	S1	7(10)W	D2	6P6-N	U	U	U	SG0	155.7
119	WhWv3	3-4	hrd	l	0	2Cdt2	S2	2(7)MD3(3)TD	C3	N	1	2D	W2G	SG0	209.1
120	Wz3	1	l	l	1WP	4Cwi	S1	7(10)W	D2	6P6-N	U	U	U	SG0	12.8
121	Rw	3-5	id	sil	2MP	4Dct2	S2	5(10)WP	D3	6S6-N	1	4D	U	SG0	37.8
122	OxWh4	3	ud	l-cl	1WA	1Bvt1	S1	2(10)MD	D2	N	1	2D	W2G	SG0	269.3
123	NcYk7	2-3	u	sil-l	2MA	3Bst1v	S1	1(9)3(1)WN	D2	2P3-N	1	1	W0	SG0	483.4
124	YkOx14	3	urd	l	1WD	1Bt1	S1	1(7)2(3)M	D2	N	1	2D	W1G	SG0	3753.5
125	WhOx1	3	ud	l	1WD	1Bt1	S1	2(10)MD	D1	N	1	2D	W1G	SG0	202.4
126	Wz3	1	l	l	2MP	4Cwi	S1	7(8)W6(2)W	D2	6P6-N	U	U	U	SG0	233.4
127	Wz2	1	l	l	2MP	4Cwi	S1	7(6)W6(4)W	D2	6P6-N	U	U	U	SG0	72.1
128	WhTg1	3	ud	l-sil	1WD	1Bvt1	S1	2(10)MD	D1	N	1	2D	W1G	SG0	303.1
129	Yk2	3-2	u	l	1WA	1Bt1	S1	1(10)	D2	N	1	2	W1K	SG0	586.2
130	OxWh4	3	h	l	0	1Bt2	S2	2(10)MD	D2	N	1	2	W2K	SG0	140.6
131	Wz3	1	l	l	1MP	4Cwi	S2	7(10)W	D3	6P6-N	U	U	U	SG0	69.5
132	YkNc13	3-2	u	fl-l	2MA	3Bst1v	S2	1(6)2(3)M3(1)W	D3	2C1-N	2	2	W1K	SG0	23.7
133	YkOx2	3-4	h	l-cl	3MA	4Dsv	S2	2(5)TM3(4)TW4(1)N	D2	3C1-N	1	2	W2K	SG0	31.3
134	YkOx4	3	h	l	1MA	2Bst2v	S1	2(9)TM4(1)W	D3	2C1-N	1	1	W1K	SG0	414.0
135	YkNc6	2-3	u	l-sil	3MA	4Dsv	S1	1(6)3(3)WN5(1)W	D3	2C2-N,1S2-N	1	1	W0	SG0	43.0

Area No.	Map Unit	Slope Class	Land Form	Surface Texture	Salinity	Irrigation	Stone Class	Agricultural Capability	pH	Wetlands Classification	Wind Eros.	Water Eros.	Past Eros.	Sand/Gravel	Acreage (ac)
136	Mw7	2-3	u	l-cl	5MA	4Cwi	S1	4(8)NW5(2)W	D3	6S6-N,4C3-N	1	1	W0	SG0	90.4
137	Mw7	2-3	u	l-cl	5MA	4Cwi	S1	4(8)NW5(2)W	D3	6S6-N,4C3-N	1	1	W0	SG0	63.9
138	YkOx7	3	h	l	3MA	4Csv	S1	2(8)TM3(2)WN	D3	2C1-N	1	2	W2K	SG0	32.3
139	YkNc6	2-3	u	l-sil	3MA	4Dsv	S1	1(6)3(3)WN5(1)W	D3	2C2-N,1S2-N	1	1	W0	SG0	1439.7
140	SP1M	2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	308.4
141	SP1M	2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	133.2
142	SP1M	2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	229.6
143	YkNc7	2-3	u	l-sil	2MA	3Bst1v	S1	1(9)3(1)NW	D3	2C2-N	1	1	W1K	SG0	1076.4
144	OxYk10	3-2	u	l-cl	1MA	2Bst1v	S2	2(7)M1(3)	D3	2C1-N	1	2	W1K	SG0	182.2
145	WhTg4	3-2	u	l-sil	1WP	1Bvt1	S1	2(7)D1(2)3(1)W	D3	3C1-N,1W3-N	1	2	W0	SG0	584.0
146	WhNr2	3-2	u	l-cl	1WP	2Cdv	S1	2(8)MD3(1)WD5(1)W	D3	3C1-N,2S2-N	1	2	W1K	SG0	3881.0
147	Lewis Lake									6P6-N					66.2
148	Wz1	1	l	l	2MPA	4Cwi	S1	5(10)W	D3	6S6-N	U	U	U	SG0	72.4
149	Yk11	3-2	u	l	1WA	1Bt1	S1	1(10)	D3	1C1-N	1	2	W0	SG0	1279.0
150	Wz2	1	l	l-cl	1WP	4Cwi	S1	6(6)W7(4)W	D3	6P6-N	U	U	U	SG0	476.9
151	WhOx1	3	ud	l	1WD	1Bt1	S1	2(10)MD	D1	N	1	2D	W1G	SG0	537.4
152	Errol Lake									6L6-N					193.1
153	YkNc4	2-3	u	l-sil	2MA	3Bst1v	S1	1(9)3(1)NW	D2	2C2-N	1	1	W0	SG0	345.3
154	SP1M	2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	275.0
155	WhOx1	3	ud	l	1WD	1Bt1	S1	2(10)MD	D1	N	1	2D	W1G	SG0	197.0
156	YkNc4	2-3	u	l-sil	2MA	3Bst1v	S1	1(9)3(1)NW	D2	2C2-N	1	1	W0	SG0	515.5
157	Rw	3-4	id	sil	2MP	4Dct2	S2	5(10)WE	D3	6S6-N	1	3D	U	SG0	39.9
158	Mw6	2-3	u	sil	5SA	4Cwi	S0	6(10)W	D3	5S6-N,4W6-N	1	1	U	SG0	548.2
159	YkNc7	2-3	u	l-sil	2MP	3Bst1v	S1	1(9)3(1)WN	D3	2C1-N	1	1	W0	SG0	833.1
160	YkOx10	3-2	u	l	2MA	3Bst1v	S1	1(7)2(2)M3(1)NW	D3	2C1-N	1	2	W0	SG0	468.1
161	Rw	3	id	sil	2MP	4Dct2	S2	4(7)WE5(3)WE	D3	3C3-N,2S3-N	1	3D	W2G	SG0	57.8
162	Mw6	2-3	u	sil	5SA	4Cwi	S0	6(10)W	D3	5S6-N,4W6-N	1	1	U	SG0	145.8
163	YkNc10	3	ud	l-sil	1WA	1Bvt1	S1	1(10)	D3	N	1	2D	W1G	SG0	285.4
164	YkNc10	3	ud	l-sil	1WA	1Bvt1	S1	1(10)	D3	N	1	2D	W1G	SG0	1137.4
165	OxYk1	3	urd	l	0	1Bt1	S1	2(6)M1(4)	D2	N	1	2D	W1G	SG0	96.7
166	Lac Vert									6L6-N					274.7
167	YkNc10	3	ud	l-sil	1WA	1Bvt1	S1	1(10)	D3	N	1	2D	W1G	SG0	328.3
168	Rw	3	id	sil	2MP	4Dct2	S2	5(10)WE	D3	5S3-N,4W3-N	1	3D	W2G	SG0	71.2
169	Yk4	2-3	u	l-cl	3MA	4Dsv	S1	1(6)3(3)WN5(1)W	D3	3C2-N,2W1-N	1	1	W0	SG0	609.0
170	SP1M	2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	164.9
171	TgWh1	3	ud	sil-l	1WA	1Bvt1	S1	2(10)M	D3	1C1-N	1	2D	W1K	SG0	1096.5
172	WvWh1	3-2	ud	l-sl	0	2Bdt1v	S1	3(5)D2(3)DM4(2)M	C3	N	1	3D	W1G	SG0	237.2
173	Rw	3	id	sil	1WP	4Dct2	S2	5(10)WE	D3	6S5-N	1	3D	U	SG0	59.6
174	Wz3	1	l	sil-l	1WP	4Cwi	S1	7(10)W	D3	6L6-N	U	U	U	SG0	335.2
175	SP1M	2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	593.8
176	Mw6	2-3	u	sil	2SA	4Cwi	S0	5(10)WN	D3	6S6-N	1	1	U	SG0	72.8
177	WvWh1	3-4	hrd	l-sl	0	2Cdt2	S1	3(6)DT2(2)DM4(2)M	C3	N	1	3D	W1G	SG0	687.8
178	OxYk1	3	urd	l	0	1Bt1	S1	2(6)M1(4)	D2	N	1	2D	W1G	SG0	86.4
179	Lac Vert Nord									6L6-N					652.9
180	WhWv3	3	ud	l	0	2Bdt1v	S1	2(7)MD3(3)D	C3	N	1	2D	W1G	SG0	1365.7
181	Wz3	1	l	l-cl	1WP	4Cwi	S2	7(10)W	C3	6P6-N	U	U	U	SG0	167.8
182	WvWh1	3-4	hrd	l-cl	0	2Cdt2	S2	3(7)DT2(3)DM	C3	N	1	3D	W1G	SG0	977.4
183	YkNc9	2-3	ud	l-cl	2MA	3Bst1v	S1	1(9)3(1)WN	D3	1C1-N	1	1D	W0	SG0	176.0
184	SP1M	2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	862.2
185	WvWh1	3	ud	l-sl	0	2Bdt1v	S2	3(5)D2(3)DM4(2)M	C3	N	1	3D	W1G	SG0	635.4
186	WhWv3	3	ud	l-sl	0	2Bdt1v	S1	2(6)MD3(4)D	C3	N	1	2D	W1G	SG0	327.6
187	Rw	3	ud	sil-l	1WA	4Dct2	S2	5(7)WP4(3)WP	D3	6S6-N	1	3D	U	SG0	111.9
188	Wz3	1	l	l-sl	1WP	4Cwi	S2	7(10)W	C3	6P6-N	U	U	U	SG0	98.1
189	WvWh1	3-4	h	l	1WA	2Cdt2	S2	2(5)DM3(5)DT	C3	1C1-N	1	2	W2K	SG0	546.3
190	Wh5	3-4	hd	l	1MP	2Dsv	S1	2(7)M4(2)W5(1)W	C3	3C2-N,2S3-N	1	2D	W1G	SG0	26.0
191	Wv4	3	ud	l-sl	0	2Bdt1p	S2	3(7)DP2(2)D5(1)P	C3	N	1	2D	W1G	SG0	696.7
192	Mw1	3-2	u	l-cl	1MA	4Cwi	S2	3(6)W4(4)W	D3	5C3-N,5S3-N	1	2	W1K	SG0	99.8
193	Mw1	3-2	u	l-cl	1MA	4Cwi	S2	3(6)W4(4)W	D3	5C3-N,5S3-N	1	2	W1K	SG0	65.4
194	WvWh1	3-4	hrd	l-sl	0	2Cdt2	S2	3(7)D2(3)DM	C3	N	2	3D	W2G	SG0	1987.6
195	Mw6	2	u	sil-sl	2MA	4Cwi	S2	5(10)W	D3	6W6-N	1	1	W0	SG0	113.3
196	WhTg1	3-2	u	l	1WA	1Bvt1	S2	2(10)M	C3	1C1-N	1	2	W1K	SG0	142.2
197	Mw1	2-3	u	ls-l	2MA	4Cwi	S0	5(10)W	D3	6W6-N	2	1	W0	SG0	72.2
198	Rw	3-4	ud	l	1MD	4Dct2	S2	5(6)WE6(4)WE	D3	5C6-N,5C5-N	1	3D	W4G	SG0	67.9
199	Ox1	3-2	ud	l	1WD	1Bt1	S2	2(10)M	D1	N	1	2D	W2G	SG1	118.3
200	WvWh1	3	hrd	sl	0	2Bdt2p	S2	4(6)MP3(4)MD	C3	N	2	2D	W2K	SG0	729.2
201	Wz3	1	l	ls-sl	1WP	4Cwi	S2	7(6)W6(4)W	C3	6P6-N	U	U	U	SG0	329.9
202	WhWv3	3-2	ud	l-sl	0	2Bdt1v	S1	2(7)MD3(3)DM	C3	N	1	2D	W1G	SG0	583.9
203	Mw6	2-3	u	sil-sl	2MA	4Cwi	S0	5(10)W	D3	6S6-N	1	1	U	SG0	168.1
204	Me4	3-2	u	ls-s	0	3Bmt1v	S0	4(6)M5(4)MF	D2	N	4	2	W4B	S4	83.7
205	Me12	3-2	u	ls	0	3Bmt1	S0	4(10)M	D3	N	3	2	W2B	S4	169.2

Area No.	Map Unit	Slope Class	Land Form	Surface Texture	Salinity	Irrigation	Stone Class	Agricultural Capability	pH	Wetlands Classification	Wind Eros.	Water Eros.	Past Eros.	Sand/Gravel	Acreeage (ac)
206	NcYk7	3-2	ud	sil-l	2MP	3Bst1v	S1	1(9)3(1)NW	D2	2C1-N	1	2D	W1G	SG0	3025.0
207	Sandy Beach Lake									6L6-N					270.8
208	WvWh1	3-4	hd	l	0	2Cdt2	S1	3(8)DT2(2)DM	C3	N	1	3D	W1K	SG0	345.1
209	OxYk1	3-2	ud	l	0	1Bt1	S1	2(8)M1(2)	D2	N	1	2D	W1G	SG0	525.7
210	Linstrum Lake									6L6-N					506.1
211	WhWv3	3	ud	l	0	2Bdt1v	S1	2(8)MD3(2)D	C3	N	1	2D	W1G	SG0	152.8
212	NcYk7	3-2	ud	sil-l	2MP	3Bst1v	S1	1(9)3(1)NW	D2	2C1-N	1	2D	W1G	SG0	984.4
213	Rw	3-4	id	sil-l	1MP	4Dct2	S2	5(6)WE4(4)WP	D3	5S5-N	1	3D	U	SG0	284.0
214	Mw1	2-3	u	sil-sic	2MA	4Cwi	S2	4(8)W5(2)P	D3	6C5-N	1	1	W0	SG0	40.0
215	Yk2	3-2	ud	l-sl	1WA	1Bvt1	S1	1(8)2(2)M	D2	N	1	2D	W1G	SG0	578.6
216	Wz3	1	l	l-sl	1WP	4Cwi	S1	7(7)W6(3)W	D2	6P6-N	U	U	U	SG0	433.9
217	Wh1	3	ud	l	0	1Bt1	S1	2(10)MD	D1	N	1	2D	W1G	SG0	209.4
218	Mw6	2-3	u	sil-l	2MA	4Cwi	S2	5(10)W	D3	6W6-N	1	1	U	SG0	87.8
219	Yk11	2-3	u	l	1WP	1Bt1	S1	1(9)2(1)M	D3	1C1-N	1	1	W0	SG0	833.4
220	Rw	3-4	id	sil-l	1MP	4Dct2	S2	5(6)WE4(4)WE	D3	4S4-N,3C4-N	1	3D	W2G	SG0	205.0
221	Yk11	3-2	ud	l	2MA	3Bst1	S1	1(9)2(1)M	D3	N	1	2D	W1G	SG0	3767.0
222	OxYk1	3	urd	l	0	1Bt1	S2	2(7)M1(3)	D2	N	1	2D	W1G	SG0	210.5
223	NcYk7	2-3	ud	l-sil	2MA	3Bst1v	S1	1(9)3(1)NW	D2	2C2-N	1	1D	W1G	SG0	121.2
224	Rw	3-5	hd	sil-l	1WPD	4Dct2	S2	6(7)WE4(3)WE	D3	5C6-N,5E6-N	1	4D	W4G	SG0	579.5
225	Yk2	3-2	ud	l	0	1Bt1	S2	1(10)	D2	N	1	2D	W1G	SG0	200.0
226	WhWv3	3	urd	l	0	2Bdt1v	S2	2(8)MD3(2)D	C3	N	1	2D	W1G	SG0	84.5
227	WhWv3	3	ud	l	0	2Bdt1v	S1	2(7)MD3(3)D	C3	N	1	2D	W1G	SG0	64.7
228	Lawley Lake									6O6-N					258.1
229	NcYk3	3-2	ud	l-sil	3MA	4Bst1v	S1	1(9)3(1)N	D3	1C2-N	1	2D	W1G	SG0	761.3
230	Nc6	3-2	u	sil-l	1SP	2Bst1v	S1	1(9)3(1)WN	D3	1C3-N	1	2	W0	SG0	513.3
231	Nc6	3-2	u	sil-l	1SP	2Bst1v	S1	1(9)3(1)WN	D3	1C3-N	1	2	W0	SG0	30.2
232	Nc6	3-2	u	sil-l	1SP	2Bst1v	S1	1(9)3(1)WN	D3	1C3-N	1	2	W0	SG0	48.5
233	Rw	3-5	hd	l-cl	1WP	4Dct2	S2	5(10)WE	D3	6W6-N	1	4D	W4G	SG0	109.9
234	OxYk7	3-2	u	l	1WP	1Bt1v	S2	2(6)M1(3)5(1)W	D2	1C2-N,1W2-N	1	2	W1B	SG0	1335.5
235	YkNc9	3-2	ud	l	3MP	4Bst1v	S2	1(9)3(1)NW	D3	2C3-N	1	2D	W1G	SG0	465.6
236	WhWv3	3-2	urd	l	0	2Bdt1v	S1	2(6)MD3(4)D	C3	N	1	2D	W1G	SG0	1085.3
237	NcYk3	3-2	ud	l-sil	3MA	4Bst1v	S1	1(9)3(1)N	D3	1C2-N	1	2D	W1G	SG0	418.5
238	Wv4	3	urd	l	0	2Bdt1v	S1	3(10)D	C3	N	1	3D	W1G	SG0	183.3
239	Stewarts Lake									6L6-N					266.3
240	WvWh1	3	urd	l	0	2Bdt1v	S2	3(7)D2(3)DM	C3	N	1	2D	W1G	SG0	226.8
241	WvWh1	4-3	hrd	l	0	2Cdt2	S2	3(8)DT2(2)DM	C3	N	1	3D	W2G	SG0	278.6
242	WhOx4	3-4	hd	l	1MP	2Cst2	S2	2(7)MD3(2)T4(1)W	D2	2C2-N	1	2D	W2K	SG0	163.5
243	WhOx3	3-2	ud	l	1MP	2Bst1	S2	2(10)MD	D1	N	1	2D	W2G	SG0	208.9
244	Wv4	3	urd	l	0	2Bdt1v	S2	3(10)D	C3	1P2-N	1	3D	W1G	SG0	145.9
245	WvWh1	3	hd	l	1WP	2Bdvt2	S2	3(7)D2(3)DM	C3	1W2-N	1	2D	W2G	SG0	1298.4
246	McArthur Lake									6L6-N					766.7
247	WvWh1	3	urd	l	0	2Bdt1v	S2	3(7)D2(3)DM	C3	N	1	2D	W1G	SG0	698.9
248	WhOx3	3-2	ud	l	1WP	1Bt1v	S2	2(9)MD5(1)W	D1	1C2-N,1W2-N	1	2D	W1G	SG0	385.9
249	Sempel Lake									6L6-N					398.4
250	YkNc6	2-3	ud	l	3MA	4Dsv	S1	1(6)3(3)WN5(1)W	D3	3W3-N,2C3-N	1	1D	W1G	SG0	295.3
251	Mw6	2-3	u	l	3MA	4Cwi	S1	5(7)W6(3)W	D3	5C5-N,5W5-N	1	1	W1B	SG0	49.4
252	WvWh1	3-4	urd	l	0	2Bdt1v	S2	3(7)D2(3)DM	C3	N	1	2D	W2G	SG0	259.5
253	WhOx1	3	hd	l	0	1Bt2	S2	2(10)MD	D1	N	1	2D	W1G	SG0	360.6
254	Knutson Lake									6L6-N					335.7
255	WvWh1	3-4	urd	l	0	2Bdt1v	S2	3(7)D2(3)DM	C3	N	1	2D	W2G	SG0	548.5
256	WvWh1	3-4	urd	l-sl	0	2Bdt1v	S2	3(6)D2(2)D4(2)M	C3	N	1	3D	W2G	SG0	5665.3
257	Mw6	3	h	l	1WP	4Cwi	S2	6(7)W5(3)W	D3	5E5-N,5W5-N	1	2	U	SG0	67.7
258	Clearwater Lake									6L6-N					92.9
259	WhOx3	3	hd	l	1WP	1Bt2pv	S2	2(7)MD4(2)P5(1)W	D1	1C2-N,1W2-N	1	2D	W1G	SG0	714.5
260	Gilkey Lake									6L6-N					102.2
261	WvWh1	3-4	hrd	l	0	2Cdt2	S2	3(8)DT2(2)D	C3	N	1	2D	W2G	SG0	349.1
262	McPhail Lake									6L6-N					153.4
263	Wz2	1	l	l	2WP	4Cwi	S2	6(10)W	D2	6E6-N	U	U	U	SG0	148.5
264	WvNr7	3-4	hrd	l-sil	0	2Cdt2	S2	3(9)D4(1)P	C3	N	1	3D	W2G	SG0	666.2
265	WvNr7	3	urd	l-sil	0	2Bdt1p	S2	3(8)D4(2)P	C3	N	1	2D	W1G	SG0	685.0
266	Wv4	3	urd	l	0	2Bdt1v	S2	3(10)D	C3	1P2-N	1	3D	W1G	SG0	627.5
267	Yk5	2-3	ud	l-sl	2MA	3Dsv	S2	1(4)2(3)M4(3)W	D3	3C3-N,2W2-N	1	1D	W1G	SG0	320.1
268	Burnett Lake									6L6-N					140.7
269	WhOx3	3	hd	l	1WP	1Bt2v	S2	2(9)MD4(1)W	D1	2C2-N	1	2D	W1G	SG0	430.5
270	WvNr7	3	hrd	l-sil	0	2Bdvt2	S2	3(10)D	C3	N	1	2D	W2G	SG0	523.9
271	WvNr3	3	urd	l-sil	0	2Bdt1v	S2	3(10)D	C3	N	1	2D	W2G	SG0	539.0
272	NcYk6	2-3	ud	sil-l	3MA	4Dsv	S2	1(7)3(2)NW4(1)W	D3	2W3-N,2C3-N	1	1D	W1G	SG0	210.3
273	Rw	3-4	uc	l	1WP	4Dct2	S0	5(6)WE7(4)W	D3	5P4-N,3E3-N	1	3	W3G	SG0	441.5
274	WhWv5	3	h	l	0	2Cdv	S2	2(4)MD3(4)DP5(2)W	C3	2W2-N,2S2-N	1	2	W1K	SG0	1053.2
275	Wv1	3-4	urd	l	0	2Bdt1v	S2	3(10)D	C3	N	1	3D	W2G	SG0	326.6

Area No.	Map Unit	Slope Class	Land Form	Surface Texture	Salinity	Irrigation	Stone Class	Agricultural Capability	pH	Wetlands Classification	Wind Eros.	Water Eros.	Past Eros.	Sand/Gravel	Acreeage (ac)
276	SP2M	2	H	o	0	4Cwi	S0	O(10)	D3	6F6-N	1	U	U	SG0	659.5
277	WvSy1	3-2	u	l-sl	0	2Cdv	S2	3(7)D5(3)MF	C1	1S2-N	2	2	W2B	S2	709.5
278	WhOx3	3	hd	l	0	1Bt2v	S2	2(9)MD5(1)W	D1	2W2-N	1	2D	W2G	SG0	151.8
279	Woods Lake									6L6-N					82.5
280	WhOx3	3	hd	l	0	1Bt2v	S2	2(9)MD5(1)W	D1	2W2-N	1	2D	W2G	SG0	443.2
281	OxCf7	3-2	u	l	1WP	1Bvt1	S2	2(9)M4(1)W	D2	1W2-N,1C2-N	1	2	W1B	SG0	401.0
282	WvWh1	3-2	ud	l	0	2Bdt1v	S2	3(6)D2(4)D	C3	N	1	2D	W1G	SG0	749.3
283	Laytons Lake									6L6-N					209.2
284	WvWh1	3-4	urd	l-sl	0	2Bdt1v	S2	3(6)D2(2)D4(2)M	C3	N	1	3D	W2G	SG0	865.7
285	HmWs6	3-2	u	fl-sl	1WP	2Dmv	S1	3(5)M4(4)MW5(1)W	C3	3C4-N	2	2	W0	G2	159.2
286	HmOx13	2-3	u	l-fl	1WP	1Cv	S2	2(4)M3(4)M5(2)W	D3	2C3-N,2W3-N	1	2	W1B	SG0	534.1
287	WvWh1	3-4	urd	l	0	2Bdt1v	S2	3(7)D2(3)D	C3	N	1	3D	W1G	SG0	608.8
288	WsMe4	3	h	gls-ls	1WP	4Bmvt2	S2	4(9)M5(1)W	D3	2S1-N	3	1	W1K	SG4	167.8
289	WsMe4	3	h	gls-ls	1WP	4Bmvt2	S2	4(9)M5(1)W	D3	2S1-N	3	1	W1K	SG4	48.1
290	Mw6	3	u	l	1MPA	4Cwi	S2	6(7)W5(3)W	D3	5W6-N,5S6-N	1	2	U	SG0	103.8
291	HmOx13	2-3	u	l-fl	1WP	1Cv	S2	2(4)M3(4)M5(2)W	D3	2C3-N,2W3-N	1	2	W1B	SG0	820.4
292	Mw6	3	u	l	1WP	4Cwi	S2	6(8)W5(2)W	D3	6W6-N,2C3-N	1	2	W0	SG0	195.9
293	HmOx13	2-3	u	l-fl	1WP	1Cv	S2	2(4)M3(4)M5(2)W	D3	2C3-N,2W3-N	1	2	W1B	SG0	99.0
294	SyNt5	3	h	s-ls	1WP	4Dmv	S0	5(6)MW4(4)MW	C3	2W1-N,2C1-N	4	2	W2B	S4	111.0
295	HmOx13	2-3	u	l-fl	1WP	1Cv	S2	2(4)M3(4)M5(2)W	D3	2C3-N,2W3-N	1	2	W1B	SG0	522.9
296	Prost Lake									6L6-N					144.6
297	WhOx3	3-2	u	l	1WP	1Bt1	S1	2(10)MD	D1	1C2-N	1	2	W1K	SG0	116.0
298	WhOx4	4-3	h	l	1WP	1Ct2	S2	3(5)T2(4)MD5(1)W	D2	2W2-N	1	2	W2K	SG0	94.0
299	Mw6	2-3	u	l	2MP	4Cwi	S1	6(7)W5(3)W	D3	5E6-N,5O6-N	1	1	W1B	SG0	189.6
300	Wz3	1	l	l	1WP	4Cwi	S2	7(10)W	C3	6L6-N	U	U	U	SG0	374.9
301	WvWh1	3-4	urd	l	0	2Bdt1v	S2	3(6)D2(4)D	C3	N	1	3D	W1G	SG0	478.0
302	OxMe2	2-3	u	l-sl	1MP	2Cdv	S2	2(5)M3(3)M4(2)W	D2	3C2-N	2	1	W0	S2	135.6
303	WvWh6	3	h	l	1WP	2Ddv	S4	5(6)PW2(2)DM3(2)D	C3	3W2-N	1	2	W0	SG0	162.0
304	Mw1	2	u	l	1WP	4Cwi	S3	4(5)WP5(5)WP	D3	5W6-N,5C6-N	1	1	W0	SG0	164.4
305	Ox20	2-3	ud	l-sl	1WP	1Cv	S2	2(5)M3(3)M4(2)W	D3	2C3-N,2W2-N	1	1D	W1G	SG0	353.8
306	Wv8	3	h	l	0	2Bdt2	S3	3(5)D4(3)P5(2)P	D1	N	1	2	W0	SG0	130.4
307	MwSP1	2	u	l-o	0	4Cwi	S2	0(5)5(5)W	D3	5W5-N,5F5-N	1	1	U	SG0	6.1
308	Mizhask Lake									6L6-N					431.3
309	Wh5	3-2	u	l	0	1Dv	S3	2(4)M4(3)WP5(3)PW	D1	3W2-N,2S2-N	1	2	W0	SG0	8.7
310	WvBd1	3	h	l-sl	0	2Bdt2	S3	3(4)D4(4)MP5(2)P	C3	1S1-N	2	2	W0	G2	180.4
311	Wv4	3-4	urd	l	0	2Bdt1v	S2	3(10)D	C3	N	1	3D	W1G	SG0	703.3
312	Wh1	3	ud	l	0	1Bt1	S2	2(10)MD	D1	N	1	2D	W1G	SG0	204.6
313	HmOx4	3-2	h	l	1WP	1Ct2	S2	2(9)M4(1)W	D1	2C2-N	1	2	W0	SG1	262.5
314	Mw6	2	u	l	2WP	4Cwi	S1	6(6)W4(4)W	D3	5S6-N,5C6-N	1	1	W1B	SG0	146.1
315	Wz3	1	l	sl-l	1WP	4Cwi	S4	7(10)W	C3	6E6-N	U	U	U	SG1	18.4
316	Wv4	3	urd	l	0	2Bdt1v	S2	3(10)D	C3	N	1	3D	W1G	SG0	686.0
317	Wz3	1	l	l	1WP	4Cwi	S3	7(10)W	D1	6L6-N	U	U	U	SG0	105.1
318	WhOx3	3	h	l	1WP	1Bt2v	S3	2(7)MD4(2)P5(1)W	D1	2W2-N,2S2-N	1	2	W1K	SG0	615.0
319	Wv4	3	urd	l	0	2Bdt1v	S2	3(10)D	C3	N	1	3D	W1G	SG0	313.1
320	WhOx3	3-2	u	l	0	1Bt1pv	S2	2(7)MD3(2)P4(1)W	D1	2W2-N	1	2	U	SG0	322.7
321	Wv4	3-4	h	sl-l	0	2Ddmv	S4	5(4)PW6(2)P4(4)MP	C3	1P2-N,1W2-N	2	2	W0	SG1	2699.8
322	Hw	6-7	ihg	sl-fl	0	4Dct2	S3	5(5)TP6(5)TE	D1	N	2	5G	U	SG0	268.1
323	Wz3	1	l	sl-l	0	4Cwi	S4	7(10)W	C3	6E6-N	U	U	U	SG1	94.8
324	Wz3	1	l	sl-l	0	4Cwi	S4	7(10)W	C3	6L6-N	U	U	U	SG1	165.1
325	Kipabiskau Lake									6L6-N					529.6
326	Hw	6-7	ihg	sl-fl	0	4Dct2	S3	5(5)TP6(5)TE	D1	N	2	5G	U	SG0	406.7
327	WvBd3	3-4	hd	sl-l	0	2Cdm2	S3	4(5)MP3(3)D5(2)P	C3	1W2-N	2	2D	W1G	G2	562.1
328	Wz3	1	l	sl-l	0	4Cwi	S3	7(10)W	C3	6P6-N	U	U	U	SG0	29.4
329	Mw6	2	u	ls-sl	0	4Cwi	S2	5(7)W6(3)W	D3	6W6-N	2	1	U	SG0	146.7
330	WvNr7	3-2	urd	l-sil	0	2Bdt1v	S2	3(10)D	C3	N	1	3D	W1G	SG0	288.1
331	MwSP1	2	u	l-o	0	4Cwi	S2	0(5)6(5)W	D3	5W5-N,5F5-N	1	1	U	SG0	381.8
332	WhTg1	2-3	id	l	0	1Bvt2	S2	2(9)MD5(1)W	D1	2W3-N	1	1D	W0	SG0	99.3
333	WvLc1	2-3	u	l-fl	0	2Bdt1v	S2	3(10)DM	C1	N	1	2	W0	SG0	773.8
334	WvSy1	3	hd	l-sl	0	2Bdt2	S3	4(6)MP3(3)D5(1)P	C1	N	2	2D	W0	S2	365.8
335	Me1	3	u	sl	0	2Bmt1	S0	3(10)M	C3	1W2-N	2	2	W0	S4	137.9
336	MP2M	1-2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	91.5
337	Wv4	3-4	hd	l-sl	0	2Cdt2	S4	3(5)D4(3)MP6(2)P	C3	N	1	2D	W0	SG1	449.4
338	HwAv1	3-6	hic	sl-cl	1WP	4Dct2	S4	5(5)TE6(5)WP	D3	4W6-N,3E5-N	2	4	W5G	SG1	1012.6
339	Ws1	3	hd	sl	0	4Bmt2	S4	4(6)M5(3)P6(1)P	C1	N	2	2D	W1G	G4	64.0
340	Wv1	3	ud	l	0	2Bdt1v	S3	3(4)D4(3)P5(3)P	C3	N	1	2D	W1G	SG1	541.0
341	GbWv1	3	ud	sl-l	0	3Dmv	S2	4(10)MP	C1	N	2	2D	W1G	G3	162.0
342	WsHm1	3	ud	gsl-l	0	3Dmv	S2	4(7)M2(3)M	C1	1W1-N	2	1D	W0	G3	87.9
343	Ws1	4-5	hd	gsl-sl	0	4Cmt2	S3	4(7)M5(3)P	C1	N	2	2D	W2G	G4	45.5
344	HmCf1	3	ud	fl-l	0	1Bt1	S1	2(7)M3(3)M	C3	N	2	2D	W1G	SG1	194.0
345	HmWs1	3	h	l-gsl	0	2Cmv	S2	2(5)M4(5)MP	C3	1C1-N	2	2	W1K	G2	84.4

Area No.	Map Unit	Slope Class	Land Form	Surface Texture	Salinity	Irrigation	Stone Class	Agricultural Capability	pH	Wetlands Classification	Wind Eros.	Water Eros.	Past Eros.	Sand/Gravel	Acreage (ac)
346	Ws1	4-5	hd	gs-l	0	4Cmt2	S3	4(7)M5(3)P	C1	N	2	2D	W2G	G4	358.8
347	Hw	6	ig	gs-l	0	4Dct2	S3	5(9)TE6(1)P	D3	N	2	5G	W5G	SG1	215.3
348	MwSP1	2	u	cl-o	0	4Cwi	S1	0(5)6(5)W	D3	5W6-N,5F6-N	1	1	U	SG0	231.3
349	SyGb1	3	h	sl-l	0	2Bmvt2	S2	4(5)MP3(4)M5(1)W	C1	2W3-N	2	2	W1B	SG4	710.1
350	WvWh1	3	hd	l-sl	0	2Bdvt2	S3	3(5)D4(2)P5(3)P	C3	N	1	2D	W1G	SG0	426.8
351	Spence Lake									6L6-N					35.9
352	WvSy1	3	h	l-sl	1WP	2Ddv	S3	4(5)MP3(2)D5(3)PW	C3	2W2-N	2	2	W1K	S2	1117.9
353	Mw6	2-3	u	cl-l	0	4Cwi	S2	5(6)W4(4)W	D3	5W6-N,5C6-N	1	1	W0	SG0	155.5
354	SP2M	1-2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	122.4
355	MeNi2	2	u	fl-sl	0	2Bmv	S0	3(10)M	B2	N	2	1	W1B	S4	240.1
356	WvGb1	3	ud	l-sl	0	2Bdt1v	S2	3(5)D4(5)MP	C3	1W2-N	2	2D	W1G	G2	543.0
357	Bu9	2	u	l	0	1Cv	S0	2(8)M3(1)W5(1)W	D3	2W2-N,2C2-N	1	2	W0	SG0	119.3
358	Bu9	2	u	l	0	1Cv	S0	2(8)M3(1)W5(1)W	D3	2W2-N,2C2-N	1	2	W0	SG0	82.5
359	WvLc1	2-3	urd	l-vl	0	2Bdt1p	S2	3(8)DM4(2)P	C1	N	1	2D	W2G	SG0	506.0
360	Sy2	3	h	sl-fl	0	2Bmvt2	S0	4(10)M	B2	1W2-N	2	2	W1B	S4	452.9
361	Rw	3-5	ihd	l-cl	0	4Dct2	S1	6(8)W3(2)TE	D1	5P5-N,3E4-N	1	4D	U	SG0	27.5
362	SbGo1	3-2	u	sl-vl	0	2Bmt1v	S0	3(7)M2(3)M	B0	1C2-N	2	2	W2B	SG0	46.4
363	Hm1GT	2	u	l-sl	0	2Bgv	S1	2(5)M3(4)M4(1)W	C1	2C2-N	1	2	W0	G2	115.7
364	Pa1	2-3	urd	sil-l	0	2Bdt1p	S2	3(8)DP2(2)DM	B0	N	1	3D	W1K	G1	333.0
365	SbGo1	3-2	u	sl-vl	0	2Bmt1v	S0	3(7)M2(3)M	B0	1C2-N	2	2	W2B	SG0	454.3
366	NiCr5	3	h	ls-sl	1WP	3Dmv	S0	3(5)M4(3)MW5(2)W	B2	4W3-N	2	2	W1B	S4	353.3
367	SP2M	2	H	o	0	4Cwi	S0	6(10)W	D1	6F6-N	1	U	U	SG0	404.7
368	Ni2	3-4	h	sl-ls	0	2Cmt2	S0	3(7)ME4(3)M	B2	N	2	2	W3B	S4	284.4
369	BuHm4	2-3	u	l-fl	1WP	1Dv	S0	2(6)M3(2)MW5(2)W	B2	4W4-N,3C4-N	1	2	W1B	SG0	1017.2
370	Mw6	2	u	sil-sl	2MA	4Cwi	S0	5(7)W6(3)W	D3	6W6-N,4E3-N	1	1	W0	SG0	148.4
371	Pingwi Lake									6L6-N					59.8
372	Pa1	3-2	rud	sil-l	0	2Cdt2	S2	3(6)DP2(3)DM4(1)P	B0	N	1	3D	W2G	G1	903.0
373	Nikk Lake									6L6-N					221.4
374	HmBb13	3-2	ud	l-fl	3MPA	4Dsv	S1	2(4)M3(4)MW5(2)W	D3	3E3-N,2W2-N	1	2D	W2G	SG0	346.7
375	LcGb1	3-4	h	vl-gls	0	2Ddmv	S1	3(5)M4(5)MW	B2	2W2-N	2	2	W1B	G2	142.7
376	SP2M	1-2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N	1	U	U	SG0	277.2
377	Pa1	3-2	rud	sil-l	0	2Cdt2	S2	3(6)DP2(3)DM4(1)P	B0	N	1	3D	W2G	G1	1703.8
378	SP2M	1-2	H	o	0	4Cwi	S1	0(8)7(2)W	D1	6F6-N,3P6-N	1	U	U	SG0	544.7
379	Gb1	3-2	ur	gs-l	0	4Bmt1p	S2	4(10)M	C1	N	2	1	W1K	G4	166.4
380	Pp7S	2	u	sil	1WP	2Cgv	S0	3(8)D4(1)W5(1)W	D1	3W2-N	1	2	W0	S3	217.6
381	GbSb1	3	hd	gs-l	0	3Dmv	S2	4(7)M3(3)M	C1	N	2	1D	W1G	G3	310.0
382	Hw	4-6	hid	l	0	4Dct2	S4	5(5)PT6(1)P4(4)PT	D3	N	1	5D	U	SG0	150.1
383	Ws1	3	hd	gs-l	0	4Bmt2	S3	4(7)M5(3)P	C1	N	2	1D	W1G	G4	42.6
384	WvBd1	3	h	l-gsl	0	2Cdv	S3	3(4)D4(4)MP5(2)P	C3	1W1-N	2	2	W0	G2	103.0
385	WvBd1	3	h	l-gsl	0	2Cdv	S3	3(4)D4(4)MP5(2)P	C3	1W1-N	2	2	W0	G2	653.0
386	Kwatapiu Lake									6L6-N					256.6
387	Hw	4-6	hid	l	0	4Dct2	S4	5(5)PT6(1)P4(4)PT	D3	N	1	5D	U	SG0	292.5
388	WvNr3	3-4	hd	l-sil	0	2Cdt2	S2	3(9)D4(1)T	C3	N	1	2D	W2G	SG0	258.1
389	Gb5	3	h	gs-l	0	4Dmv	S1	4(5)MW3(2)M5(3)W	C1	4W5-N	2	1	W0	G4	777.1
390	WvNr7	2-3	u	l-sil	1WP	2Cdv	S3	3(7)D4(2)PW5(1)P	C3	2W3-N	1	2	W0	SG1	951.5
391	Rw	3-5	hd	l	1WP	4Dct2	S3	5(6)WE4(4)TE	D3	4W3-N	1	4D	U	SG0	201.0
392	Pa3	3-2	urd	l-sil	0	2Bdt1p	S2	3(8)DP4(2)P	C1	N	1	2D	W2G	G1	220.1
393	Wz3	1	l	l-sil	1MP	4Cwi	S3	7(10)W	C3	6L6-N	U	U	U	SG1	183.9
394	Wv4	3	ud	l	0	2Bdt1v	S3	3(7)D4(2)P5(1)P	C3	N	1	2D	W1G	SG1	1453.2
395	Rw	3-5	hd	l	0	4Dct2	S2	4(5)TE5(5)WE	D3	4W3-N,2E2-N	1	4D	W4G	SG0	80.4
396	Mw6	2	u	cl	1WP	4Cwi	S2	5(4)W6(4)W7(2)W	D3	5C4-N,5P4-N	1	1	W0	SG0	195.3
397	Olsons Lake									6L6-N					157.6
398	Mw6	2-3	h	sl-l	0	4Cwi	S1	6(6)W5(3)W7(1)W	D3	6W6-N,4C4-N	2	1	W0	SG0	121.0
399	YkNc9	3-2	hd	l	2MA	3Cst2	S1	2(5)T1(4)3(1)WN	D3	2C2-N	1	2D	W1K	SG0	85.8
400	WvNr7	3	urd	l-sil	0	2Bdt1p	S2	3(8)D4(2)P	C3	N	1	2D	W2G	SG0	287.9
401	Tg5	2-3	u	l-sil	0	1Dv	S2	2(7)M3(2)W5(1)W	D1	3C3-N,2W3-N	1	2	W1B	SG0	963.7
402	WvNr3	3-4	hrd	l-sil	0	2Cdt2	S2	3(10)D	C3	N	1	3D	W3G	SG0	560.2
403	WvNr7	3-2	u	l-sil	0	2Bdt1v	S2	3(10)D	C3	N	1	2	W0	SG1	130.9
404	Mw6	3	h	l-sl	0	4Cwi	S2	5(5)W6(4)W3(1)D	D3	5W3-N,5S2-N	1	2	U	SG0	94.8
405	WvNr7	3-2	u	l-sil	0	2Bdt1v	S2	3(10)D	C3	N	1	2	W0	SG1	379.8
406	Wv4	3-4	h	l	0	2Cdt2	S2	3(8)D4(2)P	C3	N	1	2	U	SG0	152.7
407	Ka1GT	3-2	u	sil-l	0	2Bgt1v	S1	1(5)2(4)M5(1)W	C3	1W3-N,1C2-N	1	2	W0	G2	523.9
408	Wz3	1	l	sil-l	1WP	4Cwi	S1	7(10)W	C3	6L6-N	U	U	U	SG0	149.2
409	WhWv3	3-4	h	l	0	2Cdt2	S2	2(5)DM3(4)DT4(1)W	C3	2W1-N	1	2	W1K	SG0	540.9
410	Wv1	3-4	urd	l	0	2Bdt1v	S1	3(10)D	C3	N	1	2D	W3G	SG0	349.7
411	HwAv1	4-6	hid	l-cl	0	4Dct2	S4	5(4)PT6(4)PW4(2)PT	D3	3E6-N,1W5-N	1	5D	U	SG0	1198.4
412	Wading Eagle Lake									6L6-N					134.9
413	WvWh1	3-4	h	l	0	2Cdt2p	S4	5(5)P6(5)P	C3	1W1-N	1	2	W1K	SG1	1680.1
414	Wv1	2-3	ud	l	0	2Bdt1v	S2	3(10)D	C3	N	1	2D	W1G	SG0	594.7
415	SP1H	2	H	o	0	4Cwi	S0	0(10)	D3	6F6-N,2C6-N	1	U	W0	SG0	287.2

Area No.	Map Unit	Slope Class	Land Form	Surface Texture	Salinity	Irrigation	Stone Class	Agricultural Capability	pH	Wetlands Classification	Wind Eros.	Water Eros.	Past Eros.	Sand/Gravel	Acreeage (ac)
416	Wv1	3-4	urd	l	0	2Bdt1v	S2	3(10)D	C3	N	1	2D	W3G	SG0	1521.3
417	WhTg5	3	ud	l	1WP	1Cv	S1	2(5)MD1(3)A(2)W	D1	3C3-N,1W2-N	1	2D	W1G	SG0	502.8
418	Bd1	3-4	hd	gl-gsl	0	4Cmt2	S2	4(10)MF	C1	N	2	1D	W1G	G4	166.2
419	SP1H	2	H	o	0	4Cwi	S0	O(10)	D3	6F6-N	1	U	U	SG0	91.4
420	Wascana Lake									6L6-N					228.2
421	WvWh3	3	h	l	0	2Bdt2	S2	3(7)D2(3)DP	B0	N	1	2	W1K	SG0	210.4
422	Furgstrom Lake									6L6-N					45.1
423	WvWh1	3-4	h	l	0	2Cdt2	S1	3(7)DT2(3)D	C3	N	1	2	W2K	SG0	157.7
424	Perch Lake									6L6-N					207.4
425	Wz3	1	l	l	1WP	4Cwi	S1	7(6)W6(4)W	C3	6O6-N	U	U	U	SG0	73.1
426	Wv1	3-4	urd	l	0	2Bdt1v	S1	3(10)D	C3	N	1	2D	W3G	SG0	951.3
427	Wh1	2-3	ud	l	0	1Bt1	S2	2(10)MD	D1	N	1	1D	W1G	SG0	637.5
428	Lc4T	2-3	ud	sl	0	2Bdmt1	S1	3(10)M	B0	N	2	2D	W2K	SG0	353.0
429	Moss Dale Lake									6L6-N					224.4
430	Pa1	3-4	rud	sil-l	0	2Cdt2	S2	3(6)DP2(4)DM	B0	N	1	3D	W2G	G1	6.3
431	Pa1	3-4	rud	sil-l	0	2Cdt2	S2	3(6)DP2(4)DM	B0	N	1	3D	W2G	G1	1.8
432	Tg5	2-3	ud	l	1MA	2Csv	S1	2(8)M4(2)W	B2	3C3-N,2W2-N	1	2D	W1K	SG0	192.1
433	NtSy1	3	ru	sl-ls	0	2Bmt2	S0	3(6)M4(4)M	B2	N	2	2	W1B	S4	34.3
434	BuPe11	2	u	sl-ls	1MA	2Dmsv	S0	3(6)M4(3)MW5(1)W	D3	4C4-N	2	1	W1B	S2	127.2
435	Wv4	3-4	ud	l	0	2Bdt1v	S2	3(10)D	C3	N	1	2D	W2G	SG0	314.6
436	BuPe1	3	hd	sl-ls	1WD	2Bmt2	S0	3(7)M4(3)M	D1	1W2-N	2	2D	W1B	S2	566.0
437	BbHh1	2-3	u	sil-l	0	1Bt1	S0	2(6)M1(4)	D1	N	1	1	W0	SG0	123.4
438	BuPe11	3-2	u	sl-ls	1MA	2Dmsv	S0	3(6)M4(3)MW5(1)W	D3	4C4-N	2	2	W1B	S2	403.5
439	BuPe11	3-2	u	sl-ls	1MA	2Dmsv	S0	3(6)M4(3)MW5(1)W	D3	4C4-N	2	2	W1B	S2	383.5
440	WvLc1	2-3	ud	l-vl	0	2Bdt1v	S2	3(10)D	C1	N	1	2D	W1G	SG0	266.8
441	WvLc1	2-3	ud	l-vl	0	2Bdt1v	S2	3(10)D	C1	N	1	2D	W1G	SG0	225.6
442	NcCf10	2-3	ud	l-sil	1WP	1Bvt1	S1	1(5)2(4)M4(1)W	D3	2W2-N	1	1D	W2G	SG0	712.5
443	BuPe11	3-2	u	sl-ls	1MA	2Dmsv	S0	3(6)M4(3)MW5(1)W	D3	4C4-N	2	2	W1B	S2	573.0
444	NtCr1	3-2	u	ls-sl	1WP	3Dmv	S0	4(7)MW3(3)M	B4	2C2-N	2	2	W2B	S4	366.4
445	Mw6	2	u	sil-sl	1WP	4Cwi	S1	5(4)W6(3)W4(3)W	D3	5W6-N,4C5-N	1	1	W0	SG0	1146.7
446	Nt1	3	h	sl	0	2Bmt2	S0	3(10)M	B0	N	2	2	W2K	S4	124.1
447	PeWs10	3-2	u	sl-l	1MP	2Bmt1v	S1	2(6)M3(2)M4(2)M	D3	N	2	1	W0	SG4	516.5
448	NtSy6	3-2	u	ls-sl	0	3Bmt1v	S0	4(6)M3(2)M5(2)MF	B4	N	2	2	W2B	S4	976.6
449	SP1M	1	H	o	0	4Cwi	S0	O(10)	D3	6F6-N	1	U	U	SG0	234.9
450	NtCr1	3-2	u	ls-sl	1WP	3Dmv	S0	4(7)MW3(3)M	B2	2C2-N	2	2	W2B	S4	778.8
451	PeMe8	2-3	u	ls-sl	1MP	3Bmt1v	S0	3(6)M4(4)M	D3	3S2-N,1C2-N	2	1	W0	S4	153.0
452	Mw6	2-3	u	sil-sl	2MP	4Cwi	S2	5(7)W4(3)W	D3	6W6-N	1	1	W0	SG0	575.7
453	MeWs7	3-2	u	ls-gsl	1WP	3Bmt1	S1	4(10)M	B4	1W2-N	2	2	W2B	SG4	1365.9
454	Mw1	2	u	sl-ls	2MP	4Cwi	S2	3(7)W5(3)W	D3	5W6-N,4C5-N	2	1	W0	SG0	962.4
455	Rw	3-4	id	sl-l	2MP	4Dct2	S2	5(10)WE	D3	6W6-N	2	3D	U	SG0	403.5
456	PeWs10	3-2	u	sl-l	1MP	2Bmt1v	S1	2(6)M3(2)M4(2)M	D3	N	2	1	W0	SG4	619.3
457	SP1M	1	H	o	0	4Cwi	S0	O(7)6(3)W	D3	6F6-N,3C6-N	1	U	U	SG0	508.8
458	Nt1	3-2	u	ls	0	3Bmt1	S0	4(10)M	B0	N	3	2	W3B	S4	1080.7
459	SP2M	1	H	o	0	4Cwi	S0	6(7)W5(3)W	D1	6F6-N	1	U	U	SG0	54.6
460	SP2M	1	H	o	0	4Cwi	S0	6(7)W5(3)W	D1	6F6-N	1	U	U	SG0	21.3
461	SP2M	1	H	o	0	4Cwi	S0	6(7)W5(3)W	D1	6F6-N	1	U	U	SG0	22.4
462	SyNt1T	3-4	r	sl-ls	0	2Cgmt2	S1	3(5)M4(5)M	B0	N	2	2	W2B	S4	186.7
463	Mw6	1	l	sil	0	4Cwi	S0	4(7)W6(3)W	D1	5W6-N,4P4-N	1	1	W0	SG0	172.7
464	PeWs4	3-2	u	ls-sl	1MP	3Bmt1v	S1	4(8)M3(2)M	B4	N	2	2	W0	SG4	847.5
465	Mw6	2-3	h	sil	1MP	4Cwi	S0	5(6)W4(4)W	D3	6W6-N	1	1	U	SG0	145.5
466	SP2M	1	H	o	0	4Cwi	S0	6(8)W0(2)	D3	6C6-N,3F6-N	1	U	W1B	SG0	264.6
467	WhGb1	3-2	u	sl-l	1WA	2Dmv	S2	2(7)M3(3)M	B2	N	2	2	W1B	G2	669.8
468	WsHh1	3-2	u	l-sil	1WA	3Dmv	S1	3(7)M1(3)	B2	N	1	2	W0	G3	26.1
469	Rw	3-5	hc	l-sic	1MA	4Dct2	S2	6(8)WT2(2)TI	D1	6W6-N	1	4	U	SG0	104.2
470	WsHh1	3-2	u	l-sil	1WA	3Dmv	S1	3(7)M1(3)	B2	N	1	2	W0	G3	247.8
471	NrTg1	3-2	ud	sil-l	1WP	2Bdt1v	S1	2(10)MD	B2	1C2-N	1	3D	W2G	SG0	427.1
472	Eagle Lake									6L6-N					126.9
473	TgNr3	3	h	l	1WP	2Bdt2	S2	2(8)TD3(2)D	C4	1C1-N	1	2	W2K	SG0	88.8
474	Ws4	3-2	u	l-sl	0	4Bmt1v	S1	3(7)M4(3)M	C3	N	1	2	W0	G4	87.4
475	SP2M	1	H	o	0	4Cwi	S0	6(8)W0(2)	D3	6C6-N,3F6-N	1	U	W1B	SG0	76.3
476	Ws4	3-2	u	l-sl	0	4Bmt1v	S1	3(7)M4(3)M	C3	N	1	2	W0	G4	1272.9
477	NtGb1	3-2	h	ls-sl	0	3Cmt2	S1	4(7)M3(3)M	B4	N	2	2	W2B	SG4	228.8
478	SP1M	1	H	o	0	4Cwi	S2	O(10)	D3	6F6-N	1	U	U	SG0	713.0
479	WhNt1	3-2	h	l-ls	0	1Cvt2	S2	2(7)M4(3)M	C3	N	2	2	W2K	S2	414.7
480	Wv4	3	h	sil	1WP	2Bdt2p	S2	3(6)DP2(4)D	C3	N	1	2	W2K	SG0	103.7
481	TgNr3	3	h	l	1WP	2Bdt2	S2	2(8)TD3(2)D	C4	1C1-N	1	2	W2K	SG0	77.6
482	SP2M	1	H	o	0	4Cwi	S0	6(10)W	D1	6F6-N	1	U	U	SG0	260.3
483	Ws7	3-2	u	l-sl	0	4Bmt1v	S2	3(6)MP4(3)M5(1)W	D1	2W2-N	1	2	W0	G4	501.7
484	OxWs4	3-2	u	l-gls	1WP	1Dv	S2	2(6)M3(4)M	D1	N	2	2	W0	G2	818.7
485	Mw6	2-3	u	sil-sl	2MP	4Cwi	S2	5(7)W4(3)W	D3	6W6-N	1	1	U	SG0	226.5

Area No.	Map Unit	Slope Class	Land Form	Surface Texture	Salinity	Irrigation	Stone Class	Agricultural Capability	pH	Wetlands Classification	Wind Eros.	Water Eros.	Past Eros.	Sand/Gravel	Acreege (ac)
486	WhNt1	3-2	h	l-ls	0	1Cvt2	S2	2(7)M4(3)M	C3	N	2	2	W2K	S2	1129.8
487	WvWh1	2-3	u	l	0	2Bdt1p	S2	3(7)D2(3)DM	C3	N	1	2	W1K	SG0	174.9
488	OxWs4	3-2	u	l-gls	1WP	1Dv	S2	2(6)M3(4)M	D1	N	2	2	W0	G2	156.2
489	SyNt1	3-2	u	ls	0	3Bmt1	S0	5(6)MF4(4)M	B4	N	3	1	W0	S4	670.5
490	SP2M	1	H	o	0	4Cwi	S0	O(10)	D3	6F6-N	1	U	U	SG0	580.8
491	WvWh1	3-4	hrd	l-sl	0	2Cdt2	S2	3(7)D2(3)DM	C3	N	1	3D	W2G	SG0	715.6
492	WhGb1	2-3	ud	l-sl	0	1Dv	S2	2(7)M3(3)MP	C3	1C1-N	2	1D	W2G	G2	77.2
493	SP2H	1	H	o	0	4Cwi	S0	4(7)W5(3)W	D1	6F6-N	1	U	U	SG0	61.1
494	Gb1	2	u	gl-sl	0	4Am	S2	3(10)M	C4	1C1-N	2	1	W1K	G4	38.9
495	Wv4	2-3	urd	l-sl	0	2Bdt1v	S3	3(5)DP4(3)P2(2)D	C3	N	1	2D	W2G	SG0	178.1
496	WvNr4	2-3	u	l-sil	0	2Ddv	S2	3(6)D4(4)PW	C3	2W4-N	1	2	W0	SG1	1220.2
497	Wv4	3-2	hd	l	0	2Cdt2p	S4	5(4)P6(3)P3(3)DP	C3	N	1	2D	W0	SG0	411.5
498	Wv4	4-3	hd	l	1WP	2Ddpv	S4	6(6)P5(4)PW	C3	2W2-N	1	3D	W1G	SG0	793.2
499	WvWh1	3-4	hd	l	0	2Cdt2	S2	3(6)DT2(3)DM4(1)TE	C3	N	1	2D	W3G	SG0	229.0
500	PeMe8	2-3	u	ls-sl	1MP	3Bmt1v	S0	3(6)M4(4)M	D3	3S2-N,1C2-N	2	1	W0	S4	245.7
501	OxHm1	2-3	ud	l-fl	0	1Bvt1	S1	2(7)M3(3)M	D1	1W1-N	1	1D	W1G	SG0	783.6

7. GLOSSARY

- Acid soil** - A soil having a pH of less than 7.0.
- Aggregate** - A group of soil particles sticking together in such a way that they behave mechanically as a unit.
- Alkali soil** - (i) A soil having a high degree of alkalinity (pH of 8.5 or higher), or having a high exchangeable sodium content (15% or more of the exchange capacity), or both. (ii) A soil that contains enough alkali (sodium) to interfere with the growth of most crops.
- Note:** The term "alkali soil" is often incorrectly used to describe "saline soil".
- Alkaline soil** - A soil having a pH greater than 7.0.
- Alluvial deposit** - Refer to section "2.3 Surface Deposits" on page 2-1 of this report.
- Apron** - Refer to section "2.4 Surface Forms" on page 2-2 of this report.
- Aspect** - The particular direction in which a slope faces.
- Available water** - The portion of water in a soil that can be readily absorbed by plant roots. See also "field capacity".
- Bedrock** - The preglacial sediments, exclusive of stratified deposits in preglacial valleys, that underlie the surficial glacial sediments. These bedrock materials may or may not be consolidated into solid rock and may be exposed at the surface.
- Blanket** - A mantle of unconsolidated materials thick enough to mask minor irregularities in the underlying unit but still conforming to the general underlying topography.
- Blowout** - A small area from which soil material has been removed by wind.
- Channel** - The bed where a natural stream of water runs or has run.
- Clay** - (i) A soil particle that is less than 0.002 mm in diameter. (ii) A soil textural class. See also "texture, soil".
- Clod** - A compact, coherent mass of soil varying in size, usually produced by plowing or digging.
- Cobble** - Rounded or partially rounded rock or mineral fragment between 8 and 25 cm in diameter.
- Cobbly** - Containing appreciable quantities of cobbles. The term is used to describe both soil and land.
- Colluvium** - A heterogeneous mixture of material that has moved down a slope and settled at its base, as a result of gravitational action.
- Degradation** - (i) The decline in a soil's fertility status as a result of loss of organic matter, erosion by wind or water, compaction, salinization, or acidification. (ii) The changing of a soil to a more highly leached and more highly weathered condition, usually accompanied by morphological changes such as the development of an eluviated (leached), light-colored Ae horizon.
- Dissected** - Where the original surface has been cut by running water, leaving a network of channels, shallow gullies, or valleys.
- Dunes** - Wind-built ridges and hills of sand formed in the same manner as snowdrifts. They are started where some obstruction, such as a bush, boulder, or fence, causes an eddy or otherwise thwarts the sand-laden wind. Once begun, the dunes themselves offer resistance and they grow to form various shapes.
- Eluviation** - The transportation of soil material in suspension or solution within the soil by the downward or lateral movement of water.
- Eolian deposit** - Refer to section "2.3 Surface Deposits" on page 2-1 of this report.
- Erosion** - The wearing away of the land surface by running water, wind, ice or gravity.
- Erosivity** - The tendency for a soil to erode or permit erosion.
- Esker** - A winding ridge of irregularly stratified sand, gravel, and cobbles deposited under the ice by a rapidly flowing glacial stream.
- Fan** - Refer to section "2.4 Surface Forms" on page 2-2 of this report.
- Fibric layer** - A layer of organic material containing large amounts of weakly decomposed fiber whose botanical origin is readily identifiable.
- Field capacity** - The percentage of water remaining in the soil two or three days after the soil has been saturated and free drainage has practically ceased.
- Fluvial deposit** - Refer to section "2.3 Surface Deposits" on page 2-1 of this report.
- Genesis** - The mode of origin of the soil, especially the processes or soil-forming factors responsible for the development of the solum, the true soil, from unconsolidated parent material.
- Glacial till** - See "till".
- Glaciofluvial** - Refer to section "2.3 Surface Deposits" on page 2-1 of this report.
- Glaciolacustrine** - Refer to section "2.3 Surface Deposits" on page 2-1 of this report.
- Gleyed soil** - Soil affected by gleyation.
- Gleyation** - A soil forming process, operating under poor drainage conditions, which results in the reduction of iron and other elements, and in gray colors and mottles.
- Gravel** - Rock fragments between 2 mm and 7.5 cm in diameter.
- Grumic** - A fine-textured (clay or heavy clay) soil which cracks extensively when dry and forms angular blocky structures with grooved surfaces in subsoils due to the effect of swelling and shrinking during periods of wetting and drying. Surface horizons are massive, often with granular secondary structure under cultivation.
- Gully** - A channel caused by erosion from concentrated but intermittent flow of water during and immediately after heavy rains or snowmelt. It is deep enough to interfere with and not be removed by tillage operations.

Heavy soil - A soil having a high content of fine particles, particularly clay, or a soil having a high drawbar pull and therefore requiring more power to cultivate.

Horizon - Refer to section "2.1 The Soil Profile" on page 2-1 of this report.

Humic layer - A layer of organic material containing large amounts of highly decomposed organic material; only small amounts of fiber are present that can be identified as to their botanical origin. Fibers can be easily destroyed by rubbing.

Hummocky - Refer to section "2.4 Surface Forms" on page 2-2 of this report.

Humus - (i) The fraction of the soil organic matter that remains after most of the added plant and animal residues have decomposed. It is usually dark-colored. (ii) Humus is also used in a broader sense to designate the humus forms referred to as forest humus. (iii) All the dead organic material on and in the soil that undergoes continuous breakdown, change, and synthesis.

Hydraulic conductivity - The rate at which saturated soils transmit water.

Inclined - Refer to section "2.4 Surface Forms" on page 2-2 of this report.

Infiltration - The downward movement of water into the soil.

Kettle - Depression left after the melting of a detached mass of glacial ice that was buried in glacial debris.

Knob - A pronounced, rounded hill commonly found in knob and kettle topography in morainic areas.

Knoll - A small, subdued, rounded hill commonly found in knoll and depression topography in areas of till plains.

Lacustrine deposit - Refer to section "2.3 Surface Deposits" on page 2-1 of this report.

Landform - The various shapes of the land surface resulting from a variety of actions such as deposition (eskers, moraines) and erosion (gullies, valleys).

Leaching - The downward removal from the soil of materials in solution.

Level - Refer to section "2.4 Surface Forms" on page 2-2 of this report.

Light soil - A soil having a high content of coarser particles, particularly sand, or a soil having a low drawbar pull and therefore easy to cultivate.

Loess - Material transported and deposited by wind and consisting of predominantly silt-sized particles.

Mesic layer - A layer of organic material in an intermediate stage of decomposition; intermediate amounts of fiber are present that can be identified as to their botanical origin.

Microrelief - Small scale, local differences in topography, including mounds, swales, or pits that are usually < 1 m in diameter and with elevation differences of up to 2 m.

Mineral soil - A soil consisting predominantly of mineral matter. It contains less than 17% organic carbon except for an organic surface layer that may be up to 40 cm thick if formed of mixed peat or 60 cm if formed of fibric peat.

Moraine - Distinct accumulations of glacial material, mainly till, deposited directly by glaciers.

Mottles - Spots or blotches of different color or shades of color interspersed with the dominant color.

Neutral soil - A soil having a pH of 7.0.

Organic matter, soil - The organic fraction of the soil; includes plant and animal residues at various stages of decomposition, cells and tissues of soil organisms, and substances synthesized by the soil population.

Parent material - The unconsolidated and more or less chemically unweathered mineral or organic matter from which the solum of a soil has developed.

Pedology - Those aspects of soil science involving the constitution, distribution, genesis, classification and mapping of soils.

Permeability, soil - The ease with which gases and liquids penetrate or pass through a bulk mass of soil or a layer of soil.

pH, soil - The negative logarithm of the hydrogen ion activity of a soil. The degree of acidity (or alkalinity) of a soil as determined by an electrode or indicator at a specified soil-water ratio, and expressed in terms of the pH scale.

Ridged - Refer to section "2.4 Surface Forms" on page 2-2 of this report.

Rolling - Refer to section "2.4 Surface Forms" on page 2-2 of this report.

Runoff - That portion of the total precipitation on an area that flows away through stream channels and that does not enter the soil.

Runway - The channel of a stream.

Saline soil - A soil that contains enough soluble salts to interfere with the growth of most crops. The amount of salts, as measured by the electrical conductivity of the saturation extract, is greater than 4 mS/cm. Very sensitive crops may be affected at electrical conductivities of 2 mS/cm.

Sand - (i) A soil particle between 0.05 and 2.0 mm in diameter. (ii) A soil textural class. See also "texture, soil".

very coarse sand - A soil particle between 1.0 and 2.0 mm in diameter.

coarse sand - (i) A soil particle between 0.5 and 1.0 mm in diameter. (ii) A soil textural class. See also "texture, soils".

medium sand - A soil particle between 0.25 and 0.5 mm in diameter.

fine sand - (i) A soil particle between 0.10 and 0.25 mm in diameter. (ii) A soil textural class. See also "texture, soil".

very fine sand - A soil particle between 0.05 and 0.10 mm in diameter.

Silt - (i) A soil particle between 0.002 and 0.05 mm in diameter. (ii) A soil textural class. See also "texture, soil".

Soil - (i) The unconsolidated mineral material on the immediate surface of the earth that serves as a natural medium

for the growth of land plants. (ii) The unconsolidated mineral matter on the surface of the earth that has been subjected to and influenced by genetic and environmental factors of: parent material, climate (including moisture and temperature effects), macro- and microorganisms and topography, all acting over a period of time and producing a product (soil) that differs from the material from which it is derived in many physical, chemical, biological and morphological properties and characteristics.

Solum - The upper horizons of a soil in which the parent material has been modified and in which most plant roots are contained. It usually consists of the A and B horizons.

Stratification - The arrangement of sediments in layers or strata marked by a change in color, texture, dimension of particles, and composition. Stratification usually means layers of sediments that separate readily along bedding planes because of different sizes and kinds of material or some interruption in deposition that permitted changes to take place before more material was deposited.

Structure, soil - The combination or arrangement of primary soil particles into aggregates of secondary soil particles, which are separated from each other by surfaces of weakness. These secondary particles may be, but usually are not, arranged in the profile in such a manner as to give a distinct characteristic pattern. The secondary particles are characterized and classified on the basis of size, shape, and degree of distinctness. The general shape types are structureless, plate-like, block-like and prism-like. The terms are:

structureless - Having no observable aggregation or no definite orderly arrangement around natural lines of weakness.

single grain - Loose, incoherent mass of individual particles as in sands.

massive - A coherent mass showing no evidence of any distinct arrangement of soil particles.

block-like - Soil particles are arranged around a point and bounded by flat or rounded sides.

blocky (angular blocky) - Having block-like structures with flat, rectangular faces and sharp, angular corners.

subangular blocky - Having block-like structures with rounded or flattened faces and rounded corners.

granular - Having block-like aggregates that appear as spheroids having curved surfaces which have slight or no accommodation to the faces of the surrounding aggregates.

plate-like - Soil particles are arranged around a horizontal plane and generally bounded by relatively flat, horizontal surfaces.

platy - Having thin, plate-like aggregates with faces mostly horizontal.

prism-like - Soil particles are arranged around a vertical axis and bounded by relatively flat, vertical surfaces.

prismatic - Having prism-like structures with vertical faces well-defined, and edges near the top sharp and somewhat angular.

columnar - Having column-like structures with vertical edges near the top of columns not sharp (columns may be flat-topped, round-topped or irregular).

Substrate modifier - A material of different origin that underlies material in which a soil is formed, at a depth of 1 m or less.

Superglacial - A glaciolacustrine or glaciofluvial deposit laid down in small ponds or lakes on the melting ice surface, which subsequently becomes mixed with or underlain by glacial till upon melting. It is not as uniform as typical lacustrine or fluvial deposits.

Terrace - A nearly level, usually narrow, plain bordering a river, lake, or sea. Rivers are sometimes bordered by a number of terraces at different levels.

Texture, soil - The relative proportions of the various soil particles (sand, silt or clay) in a soil as described by the classes of soil texture (see Figure 2 on the following page). The limits of the various classes and subclasses are:

sand - Soil material that contains 85% or more sand.

coarse sand - Soil material that contains 25% or more very coarse and coarse sand, and less than 50% of any other one grade of sand.

fine sand - Soil material that contains 50% or more fine sand or less than 25% very coarse, coarse, and medium sand and less than 50% very fine sand.

gravelly sand - Soil material which meets the requirements of a sand but also contains 20 to 50% by volume of coarse fragments from 2 mm to 75 mm in diameter.

loamy sand - Soil material that usually contains 70 to 85% sand but may contain as much as 90% sand depending upon the amount of clay present.

gravelly loamy sand - Soil material which meets the requirements of a loamy sand but also contains 20 to 50% by volume of coarse fragments from 2 mm to 75 mm in diameter.

sandy loam - Soil material that usually contains 52 to 70% sand but may contain as much as 85% or as little as 43% sand depending upon the amount of clay present.

fine sandy loam - Soil material that contains 30% or more fine sand and less than 30% very fine sand or between 15 and 30% very coarse, coarse, and medium sand.

gravelly sandy loam - Soil material which meets the requirements of a sandy loam but also contains 20 to 50% by volume of coarse fragments from 2 mm to 75 mm in diameter.

very fine sandy loam - Soil material that contains 30% or more very fine sand or more than 40% fine and very fine sand, at least half of which is very fine sand, and less than 15% very coarse, coarse, and medium sand.

loam - Soil material that contains 7 to 27% clay, 28 to 50% silt, and less than 52% sand.

gravelly loam - Soil material that meets the requirements of a loam but also contains 20 to 50% by volume of coarse fragments from 2 mm to 75 mm in diameter.

silt loam - Soil material that contains 50% or more silt and 12 to 27% clay, or 50 to 80% silt and less than 12% clay.

silt - Soil material that contains 80% or more silt and less than 12% clay.

sandy clay loam - Soil material that contains 20 to 35% clay, less than 28% silt, and 45% or more sand.

clay loam - Soil material that contains 27 to 40% clay and 20 to 45% sand.

silty clay loam - Soil material that contains 27 to 40% clay and less than 20% sand.

silty clay - Soil material that contains 40% or more silt and more than 40% clay.

clay - Soil material that contains 40% or more clay, less than 45% sand, and less than 40% silt.

heavy clay - Soil material that contains more than 60% clay.

Till - Unstratified glacial drift, deposited directly by the ice, consisting of a mixture of clay, sand, silt, gravel, and boulders.

eroded till - Glacial till that has been subjected to water erosion subsequent to deposition, often leaving a dense stone lag or stony, gravelly lense on the surface.

water-modified till - Glacial till that has had significant amounts of water-sorted materials incorporated, usually during deposition, that results in less stony than normal glacial till with more sandy, silty or clayey textures than unstratified glacial till.

Tilth - The physical condition of soil as related to its ease of tillage, fitness as a seedbed, and impedance to seedling emergence and root penetration.

Topography - The physical features of a district or region, taken collectively; especially, the relief and contours of the land.

Undulating - Refer to section "2.4 Surface Forms" on page 2-2 of this report.

Veneer - Unconsolidated materials too thin to mask the minor irregularities of the underlying unit surface. A veneer will range from 10 cm to 1 m in thickness and will possess no form typical of the material's genesis. An example of this is shallow lacustrine deposits overlying glacial till.

Water, soil - Water occupying the pore spaces in the soil.

Water table - The upper surface of groundwater or that level in the ground where the water is at atmospheric pressure.

Wilting point - The moisture content of a soil at which plants wilt and fail to recover their turgidity when placed in a dark, humid atmosphere.

Zone, soil - An area in which the dominant soils reflect the zonal influence of climate and vegetation, and form a natural land pattern with other soils that exhibit the zonal influence only weakly or not at all. In Saskatchewan soils, there is a gradual increase in the organic matter content of the surface horizons as one moves from the southwest to the northeast, as reflected by their surface color. This forms the basis of soil zonal separations in the province, namely Brown, Dark Brown, Black, Dark Gray, and Gray.

8. FURTHER INFORMATION

For more information about the data contained in this report or for more information about the Saskatchewan Soil Survey, contact:

**Saskatchewan Soil Survey
Room 5C26 Agriculture Building
University of Saskatchewan Campus
51 Campus Drive
SASKATOON SK S7N 5A8**

**Telephone:
(306) 975-4060**

**FAX:
(306)966-4226**

If more copies of this report are required, please indicate the name of the Rural Municipality and the number of copies required.

KEY MAP

